

2016

Geriatric medicine and heat shock gene therapy in global populations

Ian J. Martins
Edith Cowan University

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworkspost2013>



Part of the [Medicine and Health Sciences Commons](#)

Martins, I. J. (2016). Geriatric Medicine and Heat Shock Gene Therapy in Global Populations. *Current Updates in Gerontology, 1*, 1-5.

This Editorial is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/3422>

Geriatric Medicine and Heat Shock Gene Therapy in Global Populations

Martins IJ^{1,2,3*}

¹Centre of Excellence in Alzheimer's Disease Research and Care, School of Medical Sciences, Edith Cowan University, Australia

²School of Psychiatry and Clinical Neurosciences, The University of Western Australia, Australia

³McCusker Alzheimer's Research Foundation, Hollywood Medical Centre, Australia

***Corresponding author:** Ian Martins, School of Medical Sciences, Edith Cowan University, 270 Joondalup Drive, Joondalup, Western Australia 6027, Australia, Tel: +61863042574; Email: i.martins@ecu.edu.au

Copyright: © 2016 Ian Martins. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source.

Original Submission

Received: December 06, 2016

Accepted: December 15, 2016

Published: December 22, 2016

Open Peer Review Status: Editorials, news items, analysis articles, and features do not undergo external peer review.

How to cite this article: Ian Martins. Geriatric Medicine and Heat Shock Gene Therapy in Global Populations. *Curr Updates Gerontol.* (2016) 1: 2.1

Current Updates in Gerontology

Keywords

Diabetes; Thermoregulation Therapy; Mitochondria; Heat Shock Gene; Geriatric; NAFLD; Fat Diet; Mitochondria; Caffeine; Bacterial Lipopolysaccharides

Editorial

In the United States and Europe the geriatric population (> 65 years) is expected to double by the year 2060 with the death rate in the European Union in the geriatric population to be greater than 80% when compared with individuals < 65 years [1,2]. Geriatrics are susceptible to the global increase in chronic diseases with diabetes and neurodegenerative disease predicted to effect and determine the increased death rate of the geriatric population in the next 40 years. A defect in a single gene versus multi gene effects may be responsible for accelerated aging connected to mitochondrial apoptosis [3] and programmed cell death with relevance to insulin resistance and the increased death rate in geriatrics.




Figure 1: In the United States and the European Union the increasing geriatric population by the year 2060 has raised concern for the increased death rates in 30% of these geriatric communities (30%) and connected to the defective heat shock gene Sirt 1 (human knockout). Major interests in diet and lifestyle has escalated that may activate Sirt 1 relevant to neuron survival and stabilization of chronic diseases with possible reversal in 30 % of geriatric individuals. The global epidemic for chronic diseases in the developing world identifies 40 % of geriatrics as the major population that are at increased risk for programmed cell death with multiple organ diseases linked to Sirt 1 defects.

The United States population is composed of white Americans, black/African Americans, native Americans, Alaska natives, Asian Americans, native Hawaiians/pacific islanders and Hispanic/Latino Americans [4]. The defect in a gene that may cause mitochondrial apoptosis in all these individuals and connected to insulin resistance identifies the heat shock gene Sirtuin 1 (Sirt 1) to be defective in these populations [5]. Sirt 1 is a nicotinamide adenine dinucleotide dependent class III histone deacetylase that targets various transcription factors involved with insulin resistance, metabolic activity and inflammation [6]. Recent interests in the role of Sirt 1 as a heat shock gene in-

dicates that thermoregulation disorders as the inducing factor with increased risk to geriatrics for the development of Type 3 diabetes, stroke, cardiovascular disease, non alcoholic fatty liver disease (NAFLD), hypertension and blood brain barrier disease [7,8]. Geriatrics in 30% of global communities (Figure 1) that are induced with the various chronic diseases now indicate the heat shock gene Sirt 1 to be defective in these various communities irrespective of racial origin.

Major interests in geriatric medicine has accelerated with diet and lifestyles changes that may stabilize mitochondrial apoptosis [9] and organ diseases in these communities. In the developing world increased plasma LPS levels have raised alarm with relevance to thermoregulation disorders [7] connected to mitochondrial apoptosis relevant to NAFLD, myocardial infarction and various organ diseases [6]. Sirt 1 and the heat shock response involve the transcription factor p53/PGC1 alpha, various heat shock proteins and the heat shock transcription factor 1 important to neuron survival and insulin receptor pathways [7]. Sirt 1 regulation of HSF1 is via PGC1 alpha that is a direct transcription repressor of HSF1 [3,10,11]. Geriatrics and brain temperature regulation may be defective with relevance to Sirt 1 gene repression by LPS [12] and induction of NAFLD with increased transport to LPS to the brain [5].

Diets that contain fat may be metabolized rapidly in individuals (< 65 years) compared with geriatrics with thermoregulation defects and defective fat metabolism [8,13]. Consumption of fats such as palm oil (palmitic acid rich) and virgin coconut oil (saturated fatty acids) [7] that are solid (20-24C) may be sensitive to abnormal body temperature dysregulation with the induction of NAFLD (Figure 1) versus the consumption of olive oil (monounsaturated) that is liquid at a temperature (4C). Dietary fat restriction reduce LPS absorption with relevance to Sirt 1/p53 interactions that are essential for uncoupling protein 1 (UCP1) expression [14] with Sirt 1 activators important to activation of thermogenesis related genes (PTEN, UCP1) [15].

In geriatrics the discovery of the heat shock gene Sirt 1 [3] has become important with relevance to the use of thermoregulation drugs that maintain the thermoregulatory set points in geriatrics [16]. Other drugs for depression and psychosis [17,18] may be inactivated with relevance to thermoregulation disorders with increased transport to the brain relevant to defective insulin therapy [19,20] that determines social interaction and behaviour. In geriatrics Sirt 1 is responsible for appetite regulation [6] and its loss from the suprachiasmatic nucleus in the hypothalamus of geriatrics result not only in central neural thermoregulatory dysregulation, circadian rhythm abnormalities but also loss appetite control (anorexia nervosa) [21]. Caffeine consumption [22] in geriatrics without NAFLD needs to be carefully determined to activate hepatic mitochondrial function that may improve post-prandial lipid metabolism (Figure 1) after consumption of meals that contain fat. However global Type 3 diabetes and neurodegenerative diseases may be irreversible with relevance to long term caffeine consumption

Current Updates in Gerontology

in geriatrics. Caffeine may be relevant to accelerated neurodegeneration and critical changes to diet, lifestyles and thermoregulation are required to prevent caffeine induced neurodegeneration. Geriatrics and links to NAFLD [8] now indicate defective caffeine metabolism [23]. Caffeine has been used to improve mitochondrial thermogenesis [24,25] but with NAFLD defective caffeine metabolism over years increases CNS caffeine transport with relevance to p53 mediated mitochondrial death relevant to neuron apoptosis [26,27].

Conclusion

The global geriatric population by the year 2060 is expected to markedly increase and global death rate in geriatric individuals is predicted to rise sharply and associated with mitochondrial apoptosis in geriatric individuals with Type 3/Type 2 diabetes, NAFLD and neurodegenerative disease. The heat shock gene Sirt 1 is critical to geriatric medicine with relevance to appetite regulation, thermoregulation disorders and defective post-prandial lipid metabolism. Fat consumption such as palm oil/coconut oil should be carefully evaluated before consumption in geriatric individuals (thermoregulation disorders) with relevance to delayed metabolism of these fats (solid at body temperature) and the induction of NAFLD. Diet, drug therapy and lifestyle changes are a critical component for thermoregulatory adaptations that allow reversal of accelerated aging in geriatric individuals with the prevention of programmed cell death.

Acknowledgements

This work was supported by grants from Edith Cowan University, the McCusker Alzheimer's Research Foundation and the National Health and Medical Research Council.

References

1. http://ec.europa.eu/eurostat/statisticsexplained/index.php/Causes_of_death_statistics_-_people_over_65
2. Administration on Aging (AoA). (http://www.aoa.acl.gov/aging_statistics/index.aspx)
3. Martins IJ. Heat Shock Gene Sirtuin 1 Regulates Post-Prandial Lipid Metabolism with Relevance to Nutrition and Appetite Regulation in Diabetes. *Int J Diabetes Clin Diagn*. 2016; 3: 120. 1-3.
4. https://en.wikipedia.org/wiki/Race_and_ethnicity_in_the_United_States Race and ethnicity in the United States - Wikipedia
5. Martins IJ. Unhealthy Nutrigenomic Diets Accelerate NAFLD and Adiposity in Global communities. *J Mol Genet Med*. 2015; 9: 1-11.
6. Martins IJ. Diet and Nutrition reverse Type 3 Diabetes and Accelerated Aging linked to Global chronic diseases. *J Diab Res Ther*. 2016; 2: 1-6.
7. Martins IJ. Type 3 diabetes with links to NAFLD and Other Chronic Diseases in the Western World. *International Journal of Diabetes*. 2016; 1:1-5.
8. Bertolotti M, Lonardo A, Mussi C, Baldelli E, Pellegrini E, Ballestri S et al. Nonalcoholic fatty liver disease and aging: Epidemiology to management. *World J Gastroenterol*. 2014; 20: 14185–14204.
9. Martins IJ. Appetite Control with Relevance to Mitochondrial Biogenesis and Activation of Post- Prandial Lipid Metabolism in Obesity Linked Diabetes. *Ann Obes Disord*. 2016; 1: 1012.
10. Raynes R, Pombier KM, Nguyen K, Brunquell J, Mendez JE, et al. The SIRT1 Modulators AROS and DBC1 Regulate HSF1 Activity and the Heat Shock Response. *PLoS One*. 2013; 8: e54364.
11. Minsky N, Roeder RG. Direct link between metabolic regulation and the heat-shock response through the transcriptional regulator PGC-1 α . *Proc Natl Acad Sci U S A*. 2015; 112: E5669-5678.
12. Martins IJ. Anti-Aging Genes Improve Appetite Regulation and Reverse Cell Senescence and Apoptosis in Global Populations. *Advances in Aging Research*, 2016; 5:9-26.
13. Tokizawa K, Uchida Y, Nagashima K. Thermoregulation in the cold changes depending on the time of day and feeding condition: physiological and anatomical analyses of involved circadian mechanisms. *Neuroscience*. 2009; 164:1377-1386.
14. Hallenborg P, Fjære E, Liaset B, Petersen RK, Murano I, et al. p53 regulates expression of uncoupling protein 1 through binding and repression of PPAR γ coactivator-1 α . *Am J Physiol Endocrinol Metab*. 2016; 310: E116-128.
15. Andrade JM, Frade AC, Guimarães JB, Freitas KM, Lopes MT, et al. Resveratrol increases brown adipose tissue thermogenesis markers by increasing SIRT1 and energy expenditure and decreasing fat accumulation in adipose tissue of mice fed a standard diet. *Eur J Nutr*. 2014; 53: 1503-1510.
16. Cuddy ML. The effects of drugs on thermoregulation. *AACN Clin Issues*. 2004; 15: 238-253.
17. Frank C. Pharmacologic treatment of depression in the elderly. *Can Fam Physician*. 2014; 60: 121–126.
18. Targum SD. Treating Psychotic Symptoms in Elderly Patients. *Prim Care Companion J Clin Psychiatry*. 2001; 3: 156–163.
19. Morris JK, Burns JM. Insulin: an emerging treatment for Alzheimer's disease dementia? *Curr Neurol Neuro-*

Current Updates in Gerontology

roschi Rep. 2012; 12: 520-527.

20. Kim B, Feldman EL. Insulin resistance as a key link for the increased risk of cognitive impairment in the metabolic syndrome. *Exp Mol Med*. 2015; 47: e149.
21. Eating Disorders Recovery Today Anorexia Nervosa in The Elderly. (www.eatingdisordersrecoverytoday.com/anorexia-nervosa-in-the-elderly)
22. Martins IJ. Food intake and caffeine determine amyloid beta metabolism with relevance to mitophagy in brain aging and chronic disease. *European Journal of Food Science and Technology*. 2016; 4: 11-17.
23. Martins IJ. Caffeine consumption with Relevance to Type 3 diabetes and accelerated brain aging. *Research and Reveiws: Neuroscience*. 2016; 1: 1-5.
24. Riedel A, Pignitter M, Hochkogler CM, Rohm B, Walker J, et al. Caffeine dose-dependently induces thermogenesis but restores ATP in HepG2 cells in culture. *Food Funct*. 2012; 3: 955-964.
25. Astrup A, Toubro S, Cannon S, Hein P, Breum L, et al. Caffeine: a double-blind, placebo-controlled study of its thermogenic, metabolic, and cardiovascular effects in healthy volunteers. *Am J Clin Nutr*. 1990; 5: 759-767.
26. Dubrez L, Coll JL, Hurbin A, Solary E, Favrot MC. Caffeine sensitizes human H358 cell line to p53-mediated apoptosis by inducing mitochondrial translocation and conformational change of BAX protein. *J Biol Chem*. 2001; 276: 38980-38987.
27. Morrison RS, Kinoshita Y. The role of p53 in neuronal cell death. *Cell Death Differ*. 2000; 7: 868-879.