

2011

Ecological Restoration of Novel Lake Districts: New Approaches for New Landscapes

Clinton D. McCullough
Edith Cowan University

Eddie J. Van Etten
Edith Cowan University

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworks2011>



Part of the [Environmental Sciences Commons](#)

[10.1007/s10230-011-0161-5](https://ro.ecu.edu.au/ecuworks2011/211)

This is an Author's Accepted Manuscript of: McCullough, C. D., & Van Etten, E. J. (2011). Ecological Restoration of Novel Lake Districts: New Approaches for New Landscapes. *Mine Water and the Environment*, 30(4), 312-319. *The final publication is available at link.springer.com here*

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworks2011/211>

1 **Ecological engineering of a novel lake district: new approaches for new landscapes**

2 Clint. D. McCullough^{1,2} & Eddie J. B. van Etten¹

3 ¹Mine Water and Environment Research Centre (MiWER), Edith Cowan University, 270
4 Joondalup Dr, Perth, WA 6027, Australia

5 ²Golder Associates, 1 Havelock St., Perth, WA 6005, Australia

6 e-mail: c.mccullough@ecu.edu.au

7

8 **Abstract**

9 Open-cut mining presents mine rehabilitation challenges which are generally well-understood for
10 terrestrial ecosystems. Mine void pit lakes often form when water fills the empty pits and these
11 are frequently of poor water quality with potential for environmental harm that may dwarf other
12 mine closure environmental issues in terms of severity, scope and longevity. This is particularly
13 so when many pit lakes occur close together to form a new landscape as a 'lake district'. Pit
14 lakes may provide opportunities where lake or wetland ecosystems are developed as beneficial
15 end uses to fulfil mining industry commitments to sustainability. As for terrestrial ecosystems, a
16 clearly articulated restoration goal and strategic plan are necessary to ensure pit lake restoration
17 toward a new, yet regionally-relevant, aquatic ecosystem, which may facilitate achievement of
18 sustainability as out-of-kind environmental offsets. Such an approach must also consider
19 obstacles to development of a self-sustaining aquatic ecosystem such as water quality and
20 ecological requirements. We recommend integration of pit lakes into their catchments as a
21 landscape restoration planning exercise with clearly-identified roles and objectives for each new
22 lake habitat and its surrounds.

23

24 **Keywords:** restoration, rehabilitation, mining, pit lake, Germany, Australia

25

1 **Introduction**

2 Increasingly frequent, and of growing scale, open-cut/cast mining has left a legacy of many
3 thousands of mining pit voids worldwide (Klapper and Geller 2002; Castendyk and Eary 2009).
4 Where backfill of pits is not an economic or feasible option and the pit extends into the water
5 table, then pit lakes ranging from very deep (e.g., hard rock mining pits >250 m deep) to shallow
6 (e.g., dredge ponds <10 m) may form (Castro and Moore 2000).

7 There is also a growing demand on many natural water resources from nearby communities.
8 Many regions have seen reduced regional recharge through increased demand or climate change
9 and decrease in quality through pollution leading to damage or complete loss of aquatic habitats
10 as a result (Pyke 2004). This demand has been simultaneous to increasing development of
11 mining activities and may even sometimes be a direct result of this activity. These pressures
12 continue to contribute to an international loss of aquatic habitat types ranging from seasonal
13 wetlands to entire lake systems.

14 Rehabilitation of post-mining terrestrial landforms to provide restored ecosystems has now
15 become a well-researched (and generally successful) practice that borrows from both disciplines
16 of ecology and engineering. Indeed, post-mining rehabilitated ecosystems are a significant
17 landscape feature in many regions with mining history. However, this landscape restoration
18 typically ceases at the edge of open-cut/cast pits, unless backfill and/or landscaping can directly
19 incorporate the pit back into the surrounding terrestrial ecosystem. Instead, pit lakes are
20 generally left unconsidered with no clear regulation or restoration aims and techniques; as
21 ‘elephants in the mine closure room’. Geochemical weathering processes such as acid and
22 metalliferous drainage (AMD) may then lead to poor water quality resulting in lake waters toxic
23 to aquatic life (McCullough 2008). Such water quality impaired pit lakes typically have few
24 environmental values and may even detrimentally affect regional water bodies through
25 contamination of surface and groundwater sources (McCullough and Lund 2006). As a result, pit
26 lakes are often a social and environmental liability to the surrounding region (Doupé and
27 Lymbery 2005), frequently underestimated in terms of their scope and magnitude of
28 environmental impacts. Indeed, of all mine closure legacies, pit lakes are frequently the most
29 severe environmental impacts to a mining region and may continue to present even after the
30 mine is closed and the greater catchment is rehabilitated (Younger 2002).

1 Notwithstanding the significant environment and community problems that can be caused by the
2 new pit lake landscape features, a number of pit 'lake districts' have formed over the past few
3 decades, or are currently being formed for closure and return to state governments over the next
4 few decades (Table 1). Through improvements in scale of mineral extraction technology, these
5 more recent pit lakes are generally deeper and of greater volume than historically. Although it is
6 often assumed that pit lakes will follow an evolution from young to mature lakes resulting in
7 lakes with a well-developed ecosystem (Kalin and Geller 1998), there are many examples of pit
8 lakes formed soon after open cut mining technologies became commonplace that have not
9 improved in environmental quality or in biological measures such as biodiversity and ecological
10 function many decades after forming (McCullough et al. 2009b). Instead, many pit lakes present
11 continued risks to surrounding natural ecosystems and it is likely that many of these new habitats
12 may continue to display degraded ecosystems relative to natural systems for many hundreds of
13 years following lake filling (Castendyk in press).

14 In contrast to these risks which pit lakes may represent to adjacent and regional environments,
15 they may nevertheless also represent significant opportunities. There are many potential benefits,
16 most of which are untapped in the pursuit of mine closure planning by mining companies and
17 regulators concentrating on terrestrial restoration outcomes. Nonetheless, if appropriate
18 restoration can be achieved these large pit lake water bodies represent potentially valuable
19 environmental and social resources (McCullough and Lund 2006; McCullough et al. 2009a)
20 particularly in the face of global aquatic ecosystem losses to man regions (Sklenička and
21 Kašparová 2008). Such post-mining use of an industry legacy would help advance expectations
22 of best-practice mining environmental sustainability when pit lakes are final landforms.

23 This paper explores options for restoration that are rarely applied to pit lakes even within their
24 restored mining landscapes. We identify both opportunity, and constraint, within a contemporary
25 mine closure and restoration context and recommend regional planning strategies to best realise a
26 restored pit lake ecosystem of significant environmental value and successfully integrated into its
27 broader ecological landscape.

28

1 **Historical and Current Practice**

2 Traditionally pit lakes and even the pit void structure itself have rarely been considered in mine
3 rehabilitation plans aside from geotechnical health and safety aspects. As such, rehabilitation has
4 specifically addressed human and animal safety risks and have generally been achieved through
5 simple structures such as earthen bunds and fences, e.g. DMP/EPA (DMP/EPA 2011). Some
6 engineering technologies even take advantage of this isolationist perception to use pit voids as
7 reservoirs for tailings storage or as sacrificial sumps for AMD or erosion products from over-
8 burden and other disturbed landforms (Loch and Vacher 2006; McCullough and Lund 2006).

9 As a result, there are very few examples of restored pit lakes internationally where pit lakes and
10 their immediate surrounds have been rehabilitated to restore ecosystem values (regional or
11 otherwise). Where restoration has been achieved it has sometimes been incidental, e.g. some of
12 the former Eastern Germany lakes that were treated as waste dumps for sewage (Charles 1998)
13 and/or some ecosystem properties and processes developing naturally but only many decades or
14 even following filling. As a result, there are no demonstrative examples of pit lake restoration
15 success for most regions and mining types. This has often been because rehabilitation of pit
16 lakes, let alone restoration of a sustainable ecosystem therein, has not often been a focus for mine
17 closure planning.

18 The use of water quality as the most common, and often sole, criteria chosen by regulators may
19 be because most countries have well developed water quality guidelines that lend themselves to
20 this application (Jones and McCullough in press). Consequently, in some instances, pit lakes
21 have been relinquished to the state with restoration requirements, or at least consideration, to
22 state or national water quality guidelines. For example, stock water drinking guidelines tend to
23 be applied by regulators and as rehabilitation goals if the regional economy is predominantly
24 agricultural, and environmental guidelines used if this is an explicit state or federal requirement
25 E.g., Axler et al. (1998) or if there is risk of discharge to other regional water bodies. Given that
26 environmental water quality guidelines typically surpass guidelines for other water body uses
27 (e.g. industrial, agricultural use), relinquishing a pit lake with environmental water quality
28 standards may allow for many other end uses as well. However, although the use of water quality
29 guidelines for environmental standards may represent this ‘gold standard’ of restoration, other,

1 equally-important, ecological variables are generally not considered (McCullough et al. 2009a;
2 Lund and McCullough in press).

3

4 **Pit lakes as Out-of-Kind Environmental Offsets**

5 Rehabilitation practices mitigating environmental impact during operations and then
6 rehabilitating remaining disturbed or reformed terrestrial habitat will undoubtedly reduce overall
7 environmental impact and biodiversity losses from the post-mining landscape. However, it is
8 difficult to see how the goal of achieving no net biodiversity loss in their operations, or even a
9 “net positive impact” (NPI) on biodiversity, often proposed by many 'blue chip' mining
10 companies, e.g., Rio Tinto Plc (2008), can be achieved when a significant proportion of the mine
11 footprint becomes inundated at completion of mining and consequent cessation of dewatering.
12 Instead, developing aquatic ecosystems in and around a pit lake may be a means of helping to
13 achieve this biodiversity protection through a form of out-of-kind offset (Figure 1). Such offsets
14 are now well-established in regulatory policy in many countries, including USA, Australia and
15 South Africa (McKenney and Kiesecker 2010) where concerns of mining and, in particular, mine
16 water issues such as pit lakes, are growing (Newmont Golden Ridge Limited 2009). In this
17 closure planning model, all mining activities ranging from the direct impacts of mining through
18 to access corridors and other peripheral disturbances away from the mine footprint result in net
19 loss of terrestrial ecosystems. Many contemporary restoration strategies may redress this loss
20 through mitigation of potential impacts, to rehabilitation of impacted sites as mine closure. Still,
21 excavation of a vast open pit that floods to form a lake will result in irreversible net loss of
22 terrestrial ecosystems. Recognition of the value of a developing aquatic ecosystem in the
23 developing pit lake, and deliberate and targeted restoration of this ecosystem toward a regionally
24 relevant aquatic ecosystem of value may be a suitable offset which redresses this net terrestrial
25 ecosystem loss. Overall, there may even be a net ecosystem value (e.g., biodiversity) or gain
26 following mine closure when the pit lake ecosystem is included in mine site rehabilitation
27 accounting. As with terrestrial ecosystems, this gain is likely to significantly develop further
28 ecological value over time as the lake fills and a burgeoning aquatic ecosystem develops.

29

1 **Applying Restoration Theory to Pit Lake Districts**

2 Restoration theory and practice guidelines are generally well developed for terrestrial ecosystems
3 and typically seek to restore the disturbed landscape towards a regional “analogue system”. This
4 analogue system may represent either the pre-mining ecosystem type that was lost or
5 alternatively a local reference ecosystem. For example, in the case of a forest lost due to mining,
6 a reasonable analogue ecosystem in the first instance would be the pre-mined exact forest type,
7 but if this was not possible then a regionally-representative forest could be selected. However,
8 there is often a gross dichotomy between mine closure criteria for terrestrial and aquatic
9 communities on rehabilitated mining leases. This difference of expectations for ecological goals
10 at mine closure extends even to the edge of the pit lake, where riparian vegetation is seldom
11 either representative of the region or self-sustaining (e.g., Figure 2). Some of this difference may
12 be because pit lake formation presents a situation where the previous terrestrial habitat now
13 forming the lakes has been absolutely lost by coverage of significant volumes of permanent and
14 often deep pit lake waters.

15 Where the degree of environmental modification is often severe, such as typically follows
16 mining, achieving a pre-mining, or even analogue ecosystem is rarely achievable. Instead, it is a
17 common scenario that post-mining landscapes are modified to such an extent that the terrestrial
18 component is no longer available for restoration to a pre-mining landscape. Similarly terrestrial
19 goals for the areas now occupied by the lakes need to be abandoned and alternative restoration
20 goals must then be sought. Pit lakes and their terrestrial surrounds are often seen as classic
21 examples of novel ecosystems, with combinations of species and environmental conditions not
22 previously found (Hobbs et al. 2009). However, this needs not lead to a complete abandonment
23 of restoring ecological values; significant areas above water that will form lake riparian and
24 catchment could, and should be, clearly identified and restored to integrate the lakes into the
25 broader regional landscape as a first goal. Obtaining at least some properties and/or values of
26 regional reference aquatic ecosystems may even be a preferred goal, especially where such other
27 amphibious ecosystems are regionally rare (Brewer and Menzel 2009). The process of
28 determining and defining appropriate goals and end point criteria for completion, as well as
29 monitoring to ensure restoration is on the right trajectory to meet these goals (Society for
30 Ecological Restoration International 2004), are therefore integral components of ecological
31 restoration relevant to developing pit lake ecosystems.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Environmental Restoration Goals for Pit Lakes

Ecological sustainability is paramount to the regional value of these new lakes and their collective lake district. As with all restoration goals, although significant management intervention may be required during periods of physical and ecological development, the objective of management should be to restore an independently self-sustaining ecosystem for both terrestrial and aquatic habitats that are integrated into the new landscape. The first step in development of a pit lake ecosystem of environmental value is to identify an ‘Identifiable Desired State’(c.f. Grant 2006) as a restoration goal. Desired environmental values may come from a number of different, and often complimentary, end points. They may include the pit lake and its catchment providing habitat for charismatic species, typically demonstrated through waterfowl and mammal species (Santoul et al. 2004). Simultaneously, the pit lakes may even provide seasonal habitat for migratory bird species protected by international (e.g., RAMSAR) or other treaties. Although it is unlikely that the inherently artificial nature of the pit lake landscape will provide for many rare species with their often specific and narrow habitat and food requirements, such as aquatic macroinvertebrates (Kumar et al. in press), some endangered species may still be able to utilise pit lake districts as long-term refugia where the catchment-scale landscape approximates that of a natural lake district (sensu Brewer and Menzel 2009). Importantly, for the pit lake and its catchment to contribute value to the regional environment, this should be achieved by having a restoration target for aquatic and amphibious littoral and riparian (lake edge shallows and immediate terrestrial margin), through to terrestrial upper ecosystems, that are considered of ecological value and are regionally representative (Van Etten in press). A caution must be made that, in order to contribute to regional biodiversity, species that are found in the lake should not be those that are already common elsewhere so that there is no net loss in biodiversity. A similar caveat may hold for the genetic diversity within species that occupy the new lake ecosystems and their catchments through artificial translocation or natural migrations. Lake district ecosystems dominated by limited gene pool or demes will likely be less genetically diverse and resilient than natural lake districts that have developed over many thousands of years (Shwartz and May 2008).

1 **Compromised Ecosystems**

2 Some pit lakes and their catchments may be so disturbed, such as through extensive and
3 inappropriate (e.g., steep and eroding) terrestrial catchment and lake morphologies, or through
4 ongoing chemical processes such as AMD, that they will present long-term legacies of
5 compromised ecosystems. Such lakes that present no environmental value, or even
6 environmental risk, will be unavoidable even with pro-active restoration strategies in place.
7 Other pit lakes may be deemed ‘unmanageable’ or ‘un-fixable’ once formed as environmental
8 legacies and liabilities are substantial, however this is not always due to limited treatment or
9 remediation knowledge, but rather lack of financial or community will. Such lakes should then
10 be regarded as impaired ecosystems. An outcome for these lakes has been proposed as 'novel'
11 ecosystems that may contribute to scientific understanding through provision of 'natural
12 experiments', c.f. Hobbs et al. (2006). Indeed, there have even been proposals to maintain
13 especially acidic lakes such as these as valuable extreme and unique ecosystems warranting
14 protection under legislation (Nixdorf et al. 2005). It is unclear, however, how a regional ecology
15 could ever benefit from the presence of such potential risk in the landscape. A more preferable
16 stance may be that restoration endeavours for pit lakes and their districts need look past
17 traditional measures of restoration success such as approximation of regional physico-chemical
18 quality and biotic diversity and assemblages and instead to focus on what fundamental ecological
19 processes that have potential to be restored (Hobbs et al. 2009). Such basic processes include
20 development of nutrient cycling, functional feeding groups and/or trophic structure that might
21 satisfactorily compare to those of regional reference aquatic systems.

22 Similarly, measuring pit lake values in terms of recovery or development of ecosystem structure
23 (sensu Bell 2001) and services, such as habitat complexity, forms of carbon storage (e.g.,
24 through net respiration:production ratios) and other measures, may help identify contributions of
25 significant ecological values to a region's natural landscape even for highly 'impaired' pit lake
26 ecosystems . Whether the lake district is natural or anthropogenic in origin may be entirely
27 academic to the provision of these ecosystem services. Indeed, such new constructs containing
28 common or even alien species may present greater opportunity for ecosystem services than their
29 natural counterparts in the landscape (Lugo 1992; Ewel and Putz 2004).

30

1 **Restoring Ecosystem Values to Impaired Pit Lakes**

2 Some impaired pit lake systems may naturally restore to ecosystems of environmental value over
3 time though natural, albeit slow, remediative processes such as succession driven along a
4 restorative trajectory, e.g., water quality remediation by primary production and sulfate reduction
5 (King et al. 1974). However, these processes may occur at too slow a rate or may be inhibited by
6 negative feedback loops presenting as degraded local stable-states, c.f. Suding & Hobbs (2009)
7 (Figure 3). Such alternative stable state models are now a popular way to describe change in
8 disturbed environments (Hobbs and Suding 2009). The ecological successions of pit lakes in
9 these instances will need to be mediated by management interventions. Due to lack of examples
10 of long-term studies, it is largely unknown to what degree pit lakes, often described as examples
11 of primary succession, (Kalin et al. 2001) follow classic succession models which presume
12 gradual, predictable recoveries.

13 Prior to mining, a landscape dominated by terrestrial ecosystems has ecological values which are
14 definable by measures such as biodiversity, presence of rare species, productivity and other
15 ecosystem services (Figure 3). During the mining period the ecosystems affected by mining may
16 face a substantial decrease in their terrestrial ecosystem value as mining operations impose pulse
17 pressures of vegetation clearance and topsoil removal, and then excavation of over-burdens and
18 actual ore extraction activities forming an open mining pit. Longer lasting pressures of vehicle
19 disturbances of dust and noise and loss of habitat connectivity around the pit void will extend
20 this phase of decreasing ecosystem value. Formation of the pit void and then flooding when
21 dewatering ceases and the pit void forms a lake, will mean significant loss of terrestrial habitat.
22 Following rehabilitation of remaining terrestrial habitats, some terrestrial ecosystem of
23 ecological value will be regained. However, terrestrial habitat of the often extensive pit area will
24 have been submerged and converted to aquatic ecosystem habitat. Terrestrial habitat is lost and
25 cannot be rehabilitated and realised as terrestrial habitat ever again.

26 Through natural ecological succession processes, this evolving lake system may develop
27 increased ecosystem values over time as some primary production begins both within and on
28 lake banks and as fauna and flora colonise (Figure 3). However, fundamental physico-chemical
29 conditions may limit ecological development of the lake below a successional threshold, even at
30 this early stage, such as through AMD toxicity or other water quality issues. The ecological

1 consequences of AMD often include low species' diversity caused by pH stress and the exposure
2 to high concentrations of heavy metals (Nixdorf et al. 2001; Lee and Kim 2007), low trophic
3 states, low nutrient concentrations and low rates of primary production. Water quality is a master
4 threshold factor for almost all pit lake ecological processes and especially for those species of
5 lower levels of biological organisation. For example, pit lake water quality frequently displays
6 chemically-driven alternative stable states as stable, albeit poor water quality (sensu Sim et al.
7 Sim et al. 2009). This may be through abiotic processes as the only determinant for that
8 particular lake e.g., ongoing and irreversible increases of salinity in lake district regions of low
9 net precipitation. Local stable states of poor water quality may also be due to biotic remediation
10 processes present but weaker than their opposite and concurrent abiotic processes, for example
11 catchment and internal formation of acidity occurring at greater rates than external and internal
12 microbial driven-alkalinity generation processes. This state of aquatic ecosystem development
13 may be very stable, largely driven by geochemical processes. For example, development of a
14 basic self-sustaining food chain with phytoplankton algae in the lake is an initial challenge,
15 largely dependent upon water toxicity and nutrient concentrations. In this example, a
16 management intervention to improve water quality, such as by active or passive remediation of
17 AMD or similar issues, may be required before ecosystem development can continue to a high
18 level of ecological complexity (Figure 3). Such restoration efforts would then use a management
19 intervention to elevate the ecological succession path above this water quality threshold so that
20 the pit lake ecosystem may continue to develop and achieve greater ecological value (Klapper
21 and Geller 2002).

22 A pit lake ecosystem with high rates of primary production may be desirable in that it contributes
23 to the ecological value of a pit lake in many ways. Algal primary producers play an important
24 role in natural lakes, providing the dominant allochthonous energy sources that are the basis of
25 lake-ecosystem food webs (Bott 1996). Primary producers can facilitate sulphate production by
26 providing a carbon source for sulfate reducing bacteria (SRB) which increase alkalinity and pH
27 in AMD impaired lakes (Lund and McCullough 2009) and also chelate metals directly causing
28 toxicity or sorbing phosphorus and overcome carbon limitation (Nixdorf and Kapfer 1998).
29 Primary producers may also accelerate development of a natural food chain. Conversely, AMD
30 may lead to low pH and high acidity, increased metal and/or other contaminant concentrations
31 and a paucity of the macro-nutrients carbon and phosphorus that all limit primary production

1 rates and primary producer biomass. These limitations may then cascade as bottom-up controls
2 on higher trophic levels and reduced abundances of taxa such as fishes and waterfowl
3 (McCullough et al. 2009b). Pit lake restoration efforts in this first instance might identify the
4 biotic processes needing assistance from abiotic factors that buffer ecological development. For
5 example, AMD with low pH and elevated metal concentrations, or other issues with water
6 quality that limit ecological succession such as low nutrient levels (e.g., phosphorus, carbon)
7 (Tittel and Kamjunke 2004).

8 Adequate and appropriate revegetation within catchments is also important in developing
9 functional lake riparian vegetation which, in turn, may play a key role in many pit lake
10 ecological processes. Even with good pit lake water quality, many pit lakes fail to attain bank
11 vegetation of any description, even after many years (Figure 3). Riparian vegetation is also
12 important to integrate pit lakes into their greater catchments to form connected and functioning
13 landscapes. There may also be interactions between terrestrial and aquatic ecosystem
14 components remediating physico-chemical water quality issues and also providing ecological
15 habitat. This interaction shows the need to clearly identify how individual ecological components
16 must be considered in the context of the overall ecosystem in pit lake ecosystem development.
17 The contribution of organic carbon from riparian and catchment vegetation was recognised many
18 years ago as a primary causative factor in water quality improvements in AMD pit lakes
19 (Campbell and Lind 1969). Riparian vegetation may also contribute to bank stabilisation,
20 facilitating further littoral and riparian establishment. The development of sustainable pit lake
21 communities finfish and large crustacea will require such an environmental suite that is more
22 holistic than just water quality; one that also includes habitat such as fallen logs and bank
23 overhangs, as well as food resources (McCullough et al. 2009b; Van Etten in press).

24

25 **Conclusions and Recommendations**

26 Aquatic habitats are increasingly diminished in their frequency and quality through both local
27 and global anthropogenic activities. Concurrently, the growing activities of open-cut mining are
28 contributing pit lake aquatic habitats to post-mining landscapes. These pit lakes environments
29 often display depauperate ecologies of little representation and value to a regional reference
30 environment and may even present an environmental risk to nearby natural water bodies because

1 of long-term ecological development inhibition due to poor water quality and/or other ecological
2 factors. There is often no or little planning for a functioning pit lake of targeted ecological value
3 with pit lakes often overlooked in rehabilitation efforts because aquatic habitats were not present
4 previously in many of these disturbed mining locales. Nonetheless, pit lakes represent significant
5 landscape restoration opportunities for replacement (or offset) of lost terrestrial habitat values
6 with the alternative habitat values of an aquatic landscape as entire lake districts.

7 Fundamental restoration theory directs mine closure planning of post-mining landscapes that will
8 contain pit lakes, to first identify end use values. These are often environmental values as
9 specific endpoints; or as endpoints that still provide for alternative uses such as for recreation or
10 aquaculture/agriculture.

11 How do we 'restore' pit lakes as ecosystems then? Achieving a desirable pit lake ecosystem will
12 involve more than just attaining good water quality. Water quality guidelines are only the
13 beginning. Recognition of limiting factors to development of a self-sustaining ecology of
14 regional values are essential. It must also be recognised that there will be much more limited
15 scope for management manipulation of the pit lake after filling; therefore, any obstacles to
16 ecosystem development should be identified and remedied as much as possible prior to filling,
17 starting with water quality. Obtaining environmental values at higher levels than simply
18 improving water quality must also be achieved through ecological approaches, a goal which is
19 frequently ignored by restoration managers and regulators (McCullough et al. 2009b; Lund and
20 McCullough in press). Such ecological approaches to develop pit lake ecosystems may assist in
21 clearly articulating targets for the long term sustainability of pit lake districts. Such ecological
22 versus physical/chemical-driven approaches also recognise mine water-affected landscapes such
23 as pit lakes as more than a geochemical environment, with consequent further (and often simple)
24 requirements for fundamental limnological and ecological processes also needing to be
25 addressed if restoration to a representative functional ecosystem is to be successful.

26 Although it is likely that their broad environmental requirements for food and habitat will be
27 very similar to those in natural systems, pit lake biota and their ecological requirements remain
28 rarely studied and poorly understood. As such, there remains a pressing need for catchment-scale
29 rehabilitation attempts of pit lakes to move towards development of aquatic ecosystems as a best
30 practice. These restoration attempts are likely to initially fall-short of attaining satisfactory

1 ecosystem values due to a lack of knowledge of both general pit lake formation and ecological
2 processes, as well as intrinsic site-specific considerations. However, monitoring and ad hoc
3 investigation studies of combined physico-chemical and ecological characteristics of these early
4 attempts will provide fertile insight for future restoration attempts.

5 In conclusion, we hope that this paper serves to develop the field of mining closure planning by
6 both considering pit lake ecosystems as desirable and valid restoration goals. Considering mine
7 waters legacies in the context of their catchments, and vice versa, will also lead to realisation of
8 more holistic environmental benefit to post-mining landscapes. We trust that the transdisciplinary
9 perspective offered by this study will translate into improved community and regulatory
10 involvement in mine closure planning, as well as providing an example to the mining industry of
11 further opportunities with which to effectively achieve environmental sustainability targets when
12 presented with these new landscape challenges.

13

14 **Acknowledgements**

15 Thanks to Will Stock and Mark Lund (Centre for Ecosystem Management, Edith Cowan
16 University), Mark Tibbett (Centre for Land Rehabilitation, University of Western Australia) and
17 Steven Dickinson (Golder Associates) for fruitful discussion that helped develop these ideas.

18

19 **References**

- 20 Axler R, Yokom S, Tikkanen C, McDonald M, Runke H, Wilcox D & Cady B (1998)
21 Restoration of a mine pit lake from aquacultural nutrient enrichment. *Restoration*
22 *Ecology* 6: 1-19
- 23 Bell LC (2001) Establishment of native ecosystems after mining —Australian experience across
24 diverse biogeographic zones. *Ecological Engineering* 17: 179-186
- 25 Bott TL (1996) Primary production and metabolism. In, *Methods in Stream Ecology*, Hauer, F.
26 R. & Lamberti, G. A. (eds.) Academic Press, San Diego, 533-556pp.
- 27 Brewer JS, Menzel T (2009) A method for evaluating outcomes of restoration when no reference
28 sites exist. *Restoration Ecology* 17: 4-11

- 1 Campbell RS, Lind OT (1969) Water quality and aging of lakes. *Journal of the the Water*
2 *Pollution Control Federation* 41: 1943-1955
- 3 Castendyk D (in press) Lessons Learned from Pit Lake Planning and Development. In, *Mine Pit*
4 *lake Closure and Management*, McCullough, C. D. (ed.) Australian Centre for
5 Geomechanics, Perth, Australia,
- 6 Castendyk D, Eary T (2009) The nature and global distribution of pit lakes. In, *Mine Pit Lakes:*
7 *Characteristics, Predictive Modeling, and Sustainability* Castendyk, D.; Eary, T. & Park,
8 B. (eds.) Society for Mining Engineering (SME), Kentucky, USA, 1-11pp. 978-0-87335-
9 305-2.
- 10 Castro JM, Moore JN (2000) Pit lakes: their characteristics and the potential for their
11 remediation. *Environmental Geology* 39: 254-260
- 12 Charette T, Wylynko D (in press) Generating Regional Guidance for Pit lake Reclamation. In,
13 *Mine Pit lake Closure and Management*, McCullough, C. D. (ed.) Australian Centre for
14 Geomechanics, Perth, Australia,
- 15 Charles D (1998) Wasteworld. *New Scientist* 157: 32
- 16 DMP/EPA (2011) Guidelines for preparing mine closure plans. Western Australian Department
17 of Mines and Petroleum (DMP), Environmental Protection Authority of Western
18 Australia (EPA), Perth, Australia. 70pp.
- 19 Doupé RG, Lymbery AJ (2005) Environmental risks associated with beneficial end uses of mine
20 lakes in southwestern Australia. *Mine Water and the Environment* 24: 134-138
- 21 Ewel JJ, Putz FE (2004) A place for alien species in ecosystem restoration. *Frontiers in Ecology*
22 *and the Environment* 2: 354–360
- 23 Grant C (2006) State-and-transition successional model for bauxite mining rehabilitation in the
24 jarrah forest of Western Australia. *Restoration Ecology* 14: 28-37
- 25 Hobbs RJ, Arico S, Aronson J, Baron JS, Bridgewater P, Cramer VA, Epstein PR, Ewel JJ, Klink
26 CA, Lugo AE, Norton D, Ojima D, Richardson DM, Sanderson EW, Valladares F, Vilà
27 M, Zamora R & Zobel M (2006) Novel ecosystems: theoretical and management aspects
28 of the new ecological world order. *Glob Ecol Biogeogr* 15: 1-7
- 29 Hobbs RJ, Higgs E & Harris JA (2009) Novel ecosystems: implications for conservation and
30 restoration *Trends Ecol Evol* 24: 599-605
- 31 Hobbs RJ, Suding KN (2009) Hobbs RJ, Suding KN (eds.) *New Models for Ecosystem*
32 *Dynamics and Restoration* Island Press. 352pp.
- 33 Jones H, McCullough CD (in press) Regulator guidance and legislation relevant to pit lakes. In,
34 *Mine Pit lake Closure and Management*, McCullough, C. D. (ed.) Australian Centre for
35 Geomechanics, Perth, Australia,

- 1 Kalin M, Cao Y, Smith M & Olaveson MM (2001) Development of the phytoplankton
2 community in a pit-lake in relation to water quality changes. *Water Research* 35: 3215-
3 3225
- 4 Kalin M, Geller W (1998) Limnological fundamentals of acid mining lakes. In, *Acidic Mining*
5 *Lakes*, Geller, W.; Klapper, H. & Salomons, W. (eds.) Springer, Berlin, 423-425pp.
- 6 King DL, Simmler JJ, Decker CS & Ogg CW (1974) Acid strip mine lake recovery. *Journal of*
7 *the Water Pollution Control Federation* 46: 2301-2315
- 8 Klapper H, Geller W (2002) Water quality management of mining lakes – a new field of applied
9 hydrobiology. *Acta Hydrochimica et Hydrobiologica* 29: 363-374
- 10 Kumar RN, McCullough CD & Lund MA (in press) Pit lakes in Australia. In, *Acidic Pit Lakes -*
11 *Legacies of surface mining on coal and metal ores*, Geller, W. & Schultze, M. (eds.)
12 Springer, Berlin, Germany,
- 13 Lee J, Kim Y (2007) A quantitative estimation of the factors affecting pH changes using simple
14 geochemical data from acid mine drainage. *Environmental Geology* 55: 65-75
- 15 Loch R, Vacher C (2006) Assessing and managing erosion risk for constructed landforms on
16 minesites. *Proceedings of the Goldfields Environmental Management Group Workshop*
17 *on Environmental Management 2006*. Kalgoorlie, Australia. 24th - 26th May, 121-127pp.
- 18 Lugo AE (1992) Comparison of tropical tree plantations with secondary forests of similar age.
19 *Ecological Monographs* 62: 2-41
- 20 Lund MA, McCullough CD (2009) Biological remediation of low sulphate acidic pit lake waters
21 with limestone pH neutralisation and amended nutrients *Proceedings of the International*
22 *Mine Water Conference*. Pretoria, South Africa. 19-23 October, International Mine Water
23 Association, 8pp.
- 24 Lund MA, McCullough CD (in press) Restoring pit lakes: factoring in the biology. In, *Mine Pit*
25 *lakes: Closure and Management*, McCullough, C. D. (ed.) Australian Centre for
26 Geomechanics, Perth, Australia,
- 27 McCullough CD (2008) Approaches to remediation of acid mine drainage water in pit lakes.
28 *International Journal of Mining, Reclamation and Environment* 22: 105-119
- 29 McCullough CD, Hunt D & Evans LH (2009a) Sustainable development of open pit mines:
30 creating beneficial end uses for pit lakes. In, *Mine Pit Lakes: Characteristics, Predictive*
31 *Modeling, and Sustainability* Castendyk, D.; Eary, T. & Park, B. (eds.) Society for
32 *Mining Engineering (SME)*, Kentucky, USA, 249-268pp. 978-0-87335-305-2.
- 33 McCullough CD, Lund MA (2006) Opportunities for sustainable mining pit lakes in Australia.
34 *Mine Water and the Environment* 25: 220-226


- 1 McCullough CD, Steenbergen J, te Beest C & Lund MA (2009b) More than water quality:
2 environmental limitations to a fishery in acid pit lakes of Collie, south-west Australia.
3 Proceedings of the International Mine Water Conference. Pretoria, South Africa. 19-23
4 October, International Mine Water Association, 507-511pp.
- 5 McKenney BA, Kiesecker JM (2010) Policy development for biodiversity offsets: a review of
6 offset frameworks. *J Environ Manag* 45: 165-176
- 7 Newmont Golden Ridge Limited (2009) BBOP Pilot Project Case Study. Akyem Gold Mining
8 Project, Eastern Region, Ghana. Accra, Ghana. Available from [www.forest-](http://www.forest-trends.org/biodiversityoffsetprogram/guidelines/newmont-casestudy.pdf)
9 [trends.org/biodiversityoffsetprogram/guidelines/newmont-casestudy.pdf](http://www.forest-trends.org/biodiversityoffsetprogram/guidelines/newmont-casestudy.pdf),
- 10 Nixdorf B, Fyson A & Krumbeck H (2001) Review: plant life in extremely acidic waters.
11 *Environmental and Experimental Biology* 46: 203-211
- 12 Nixdorf B, Kapfer M (1998) Stimulation of phototrophic pelagic and benthic metabolism close
13 to sediments in acidic mining lakes. *Water, Air, and Soil Pollution* 108: 317-330
- 14 Nixdorf B, Lessmann D & Deneke R (2005) Mining lakes in a disturbed landscape: application
15 of the EU Water Framework Directive and future management strategies. *Ecological*
16 *Engineering* 24: 67-73
- 17 NSW EPA (2002) Green offsets for sustainable development, a concept paper. New South Wales
18 Environmental Protection Authority (NSW EPA), Sydney.
- 19 Otahel'ová H, Ot'ahel' J (2006) Distribution of aquatic macrophytes in pit lakes in relation to the
20 environment (Borská Nížina lowland, Slovakia). *Eklógia* 25: 398-411
- 21 Pyke CR (2004) Habitat loss confounds climate change impacts. *Frontiers in Ecology and the*
22 *Environment* 2: 178-182
- 23 Rio Tinto (2008) Rio Tinto's Biodiversity Strategy. Rio Tinto London, UK. 38pp.
- 24 Sánchez-Espanã J, Lo´pez-Pamo E, Santofimia E & Diez-Ercilla M (2008) The acidic mine pit
25 lakes of the Iberian Pyrite Belt: an approach to their physical limnology and
26 hydrogeochemistry. *Applied Geochemistry* 23: 1,260–1,287
- 27 Santoul F, Figuerola J & Green A (2004) Importance of gravel pits for the conservation of
28 waterbirds in the Garonne River floodplain (southwest France). *Biodivers Conserv* 13:
29 1,231-1,243
- 30 Schultze M, Geller W, Benthaus FC & Jolas P (in press) Filling and management of pit lakes
31 with diverted river water and with mine water – German experiences. In, *Mining Pit*
32 *Lakes: Closure and Management*, McCullough, C. D. (ed.) Australian Centre for
33 Geomechanics, Perth, Australia,

- 1 Shwartz RS, May B (2008) Genetic evaluation of isolated populations for use in reintroductions
2 reveals significant genetic bottlenecks in potential stocks of Sacramento Perch. *Trans Am*
3 *Fish Soc* 137: 1,764-1,777
- 4 Sim LL, Davis JA & Chambers JM (2009) Development of conceptual models for ecological
5 regime change in temporary Australian wetlands. In: Hobbs, R. J. & Suding, K. N.,
6 (Hobbs, R. J. & Suding, K. N.(Hobbs, R. J. & Suding, K. N.s Hobbs, R. J. & Suding, K.
7 N.). *New Models for Ecosystem Dynamics and Restoration* Island Press.
- 8 Sklenička P, Kašparová I (2008) Restoration of visual values in a post-mining landscape. *Journal*
9 *of Landscape Studies* 1: 1-10
- 10 Society for Ecological Restoration International (2004) *The SER International Primer on*
11 *Ecological Restoration*. http://www.ser.org/content/ecological_restoration_primer.asp. .
- 12 Suding KN, Hobbs RJ (2009) Threshold models in restoration and conservation: a developing
13 framework. *Trends Ecol Evol* 24: 271-279
- 14 Tittel J, Kamjunke N (2004) Metabolism of dissolved organic carbon by planktonic bacteria and
15 mixotrophic algae in lake neutralisation experiments. *Freshwater Biology* 49: 1062-1071
- 16 Van Etten EJB (in press) Riparian vegetation considerations for pit lakes. In, *Pit lake Mine*
17 *Closure and Management*, McCullough, C. D. (ed.) Australian Centre for Geomechanics,
18 Perth, Australia,
- 19 Younger PL (2002) Mine waste or mine voids: which is the most important long-term source of
20 polluted mine drainage?, www.mineralresourcesforum.org/docs/pdfs/younger1102.pdf,
21 Accessed: 22nd February 2006.
- 22
- 23

1 Table 1. Examples of pit lake districts internationally.

Lake District	Country	Number of lakes in District	Reference
Athabaskan Oil Sands region	Canada	0 current (26 proposed)	(Charette and Wylynko in press)
Borská Nížina lowlands	Slovakia	11 current	(Otahel'ová and O'tahel' 2006)
Central German and Lusatian districts; Rhenish district	Germany	370 current; 205 current	(Schultze et al. in press)
Collie Lake District	Australia	13 current (more proposed)	(Kumar et al. in press)
Iberian	Spain	22 current	(Sánchez-Espanã et al. 2008)


2



1

1





1

1 **Figure 1.** Hierarchy of increasing biodiversity achievements through standard-practice terrestrial
2 rehabilitation and then inclusion of pit lake aquatic ecosystem in post mining landscape
3 restoration efforts. After NSW EPA (2002).

4

5 **Figure 2.** A typical 'bathtub' ring effect showing failure of a functional riparian vegetation
6 community to develop. WO3 lake (50 years old), Collie Lake District, Australia.

7

8 **Figure 3.** Successional development of a pit lake ecosystem from low ecological value
9 immediately following mining to attainment of prior ecological value, albeit now dominated by
10 aquatic ecosystems. Local-stables states demonstrate fundamental ecological thresholds
11 management restoration activities must overcome to realise a self-sustaining aquatic ecosystem
12 of value. After Grant (2006).