

2015

The preparation and culture of washed human sperm: a comparison of a suite of protein-free media with media containing human serum albumin

Kelli L. Peirce
Edith Cowan University

Peter Roberts
Edith Cowan University, p.roberts@ecu.edu.au

Jaffar Ali

Phillip Matson
Edith Cowan University

[10.1016/j.apjr.2015.06.006](https://ro.ecu.edu.au/ecuworkspost2013/2430)

Originally published as: Peirce, K.L., Roberts, P., Ali, J., Matson, P.L. (2015). The preparation and culture of washed human sperm: A comparison of a suite of protein-free media with media containing human serum albumin, *Asian Pacific Journal of Reproduction*. 4(3) 222-227.

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/2430>



HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Reproduction

journal homepage: www.apjr.netOriginal research <http://dx.doi.org/10.1016/j.apjr.2015.06.006>

The preparation and culture of washed human sperm: A comparison of a suite of protein-free media with media containing human serum albumin

Kelli L. Peirce^{1,2*}, Peter Roberts¹, Jaffar Ali³, Phillip Matson^{1,2}¹Edith Cowan University, School of Medical Sciences, Joondalup, Western Australia, 6027, Australia²Fertility North, Suite 30, Level 2 Specialist Medical Centre East, Joondalup Health Campus, Shenton Avenue, Joondalup, Western Australia, 6027, Australia³Department of Obstetrics and Gynaecology, Women's & Children's Health Complex, University of Malaya Medical Centre, Faculty of Medicine, University of Malaya, 59100, Kuala Lumpur, Malaysia

ARTICLE INFO

Article history:

Received 27 Mar 2015

Received in revised form 20 May 2015

Accepted 24 May 2015

Available online 24 July 2015

Keywords:

Sperm

Sperm preparation

HSA

Methylcellulose

ABSTRACT

Objective: To compare two suites of culture media (one with HSA and one protein-free (PF) supplemented with methylcellulose) for washing human sperm in IVF.**Methods:** Semen samples ($n = 41$) underwent parallel density gradient preparation using PF or HSA-supplemented culture medium and subsequent yield, survival, morphology and motility were compared.**Results:** The PF medium resulted in a significantly higher sperm yield ($P < 0.0001$), but similar sperm morphology ($P = 0.822$) and 24-h survival ($P = 0.11$). There was, however, a lower percentage of progressively motile sperm ($P < 0.0001$) and a higher proportion of sperm demonstrating non-progressive motility ($P < 0.0001$) in the PF medium when observed on a Makler Chamber, apparently an artefact as a similar sperm motility index was measured using a Sperm Quality Analyser ($P = 0.83$). Attachment of sperm in PF medium to the glass chamber reduced with time and any differences had disappeared after 6 min on the counting chamber.**Conclusion:** These results support the use of PF media supplemented with methylcellulose as an alternative to HSA, although a modification to the manufacturer's protocol of 6-min pre-incubation before assessing sperm motility must be used. Further studies should investigate the function of such sperm prepared in PF medium.

1. Introduction

The desire to use serum-free medium exists in a number of industries, ranging from the pharmaceutical industry where there is a need to produce biopharmaceutical products free of infectious agents [1,2], to the area of toxicity testing, in which novel protein biomarkers of cells in culture need to be monitored with minimal background contamination [3]. But with over 70 000 treatment cycles and 13 000 babies born each year in Australia alone following the culture of human embryos *in vitro* [4], assisted reproductive technology (ART) is a major user of medium which, when used to culture preimplantation embryos for less than one week after fertilisation, can have

profound effects on the health and characteristics of the foetus and children later in life [5–9].

IVF medium must be safe and effective [10], and over the years has contained protein supplemented in the form of patient serum through to human serum albumin (HSA) as a means of improving the growth of the embryos [11]. Nevertheless, the various protein supplements have each presented their own problems regarding safety with pooled patient serum having resulted in the transmission of hepatitis B to recipients [12] and a batch of HSA prepared from donated blood having been withdrawn because of a donor having been found to be infected with Creutzfeldt–Jakob disease [13]. Whilst albumin can have significant batch-to-batch variability [14], HSA is beneficial in improving fertilisation, cleavage and pregnancy rates [15] but even current day practices of HSA purification still results in the presence of a large variety of non-declared proteins of unknown consequences to the health of patients and future offspring [16,17]. Given the aforementioned

*Corresponding author: K.L. Peirce, Edith Cowan University, School of Medical Sciences, Joondalup, Western Australia, 6027, Australia.

E-mail: kelli.peirce@fertilitynorth.com.au

Peer review under responsibility of Hainan Medical College.

limitations and risks of the use of human serum albumin in ART, some interest has been shown in the identification and/or development of more stable and safer alternatives. There are several macromolecules derived from non-animal sources that have been investigated for use as a supplement for ART culture media [18–20], though many of these alternatives are yet to be incorporated into commercially available culture media and adopted into routine clinical use. One exception, however, is methylcellulose: a non-toxic, non-allergenic, non-carcinogenic, non-infectious, plant-derived macromolecule with antioxidant and osmolytic properties [21]. A formulation of IVF medium which contains methylcellulose and is truly protein-free has shown excellent results in preliminary trials [22–24], although experience with a commercially-available version has not been so encouraging [25].

Whilst a commercially-available protein-free ART medium would have merit, the ability of such a medium to support the many steps involving gametes and embryos in an IVF procedure must be demonstrated, and given the apparent lack of investigations into same, a systematic appraisal of the various individual steps is warranted. The aim of the present study was, therefore, to assess the effectiveness of a commercially-available protein-free medium and its associated density gradient system when applied to sperm preparation in terms of sperm yield, motility, morphology and motility maintenance in culture, and to compare this to a commercially-available suite of culture media supplemented with HSA as they might be used in a clinical setting.

2. Materials and methods

2.1. Study design

The present study involved the analysis and parallel preparation of semen samples from men attending Fertility North for fertility assessment between August and December 2013. The study design ensured that each sample was prepared in an identical fashion using both media types to allow direct comparison, using only one sample per man to avoid pseudo-replication. The study was approved by the Joondalup Health Campus Human Research Ethics Committee and the Ethics Committee of Edith Cowan University, and all participants provided informed consent for their samples to be utilised in the research.

2.2. Collection of semen

Semen samples were obtained over the full period of the study from a total of 41 men (average \pm SEM age of 35.4 ± 1.0 years). Each man was instructed to abstain from ejaculation for 2–5 days and then produce a semen sample by masturbation into a wide-mouthed, 60 mL sterile specimen container (TechnoPlas, Australia). After production, each sample was held in a warming oven (Memmert, Germany) at 37 °C for a maximum of 60 min until analysis and processing.

2.3. Semen and sperm assessment

All semen samples underwent routine semen analysis following protocols employed routinely by the clinic, which are based upon the methods described in the World Health Organisation Guidelines [26]. Washed sperm were assessed using the same methodology. To reflect clinical practice, sperm

concentration was determined for each sample at a dilution of 1:20 using an Improved Neubauer Haemocytometer. A 50 μ L aliquot of neat semen or the final sperm suspension was added to 950 μ L of water and mixed well to ensure the suspension was evenly distributed. Then 10 μ L of this diluted suspension was pipetted under the cover slip of each chamber of the haemocytometer and allowed to settle for 5 min in a humidified chamber, before the sperm were counted. Sperm morphology was evaluated by making duplicate air-dried smears and staining with DiffQuick[®] (Bactolaboratories, Australia). A minimum of 200 sperm were assessed at $\times 400$ magnification under brightfield illumination, and the proportion of sperm with normal morphology expressed as a percentage of the total number of sperm assessed. For the assessment of sperm motility, 10 μ L of neat semen or the final sperm suspension was aliquoted onto the surface of a Makler chamber (Sefi Medical Instruments Ltd, Israel) and the cover slip put in place. At least 200 sperm were observed at 200 \times using phase-contrast microscopy, and their motility categorised as rapid progressive (“A”), slow progressive (“B”), non-progressive (“C”) or immotile (“D”). In addition, sperm motility was assessed in some samples using an automated Sperm Quality Analyser (SQA IIB, Medical Electronic Systems, Israel). This form of assessment uses deep chambers and the instrument detects movement of sperm suspended in medium by measuring light scatter and calculating a sperm motility index (SMI) [27].

2.4. Sperm preparation and culture

Two suites of culture media were used, one supplemented with HSA (Quinn's Advantage HSA [#3001] and Quinn's Advantage Medium with Hepes [#1023], Sage Biopharma, USA; Puresperm[®] #PS100-250, Nidacon, Sweden) and the other being protein-free (CellCura[®] PF Protein Free GradiART[™] Upper and lower Layer Medium [#GP21050 and #GP31050], PF Protein Free Gamete Handling Medium with HEPES and Gentamicin [#GP700500], CellCura, Norway). All preparations were performed “blind” so the scientist doing the work was unaware which preparation was HSA or PF.

Following routine semen analysis, all 41 semen samples underwent parallel density gradient preparation in each suite of media. Depending on the volume of semen available, between 0.5 and 1.0 mL semen was applied to discontinuous gradients of equal volumes (1 mL each) of 95% and 50% Puresperm or Upper and Lower CellCura gradients in conical centrifuge tubes (#352095, Falcon Corning, USA). The same volume of semen was applied to each of the two types of gradients to avoid discrepancies. Each gradient tube was centrifuged (#C28A, Boeco, Germany) at 350 g for 15 min. The resulting pellet was removed and resuspended in a clean Falcon tube (352095, Falcon Corning, USA) in 2 mL of the corresponding culture media: Sage (HSA) or CellCura (PF). Each suspension was washed by centrifugation, at 500 g for 5 min. The supernatant was removed by aspiration with a clean Pasteur pipette (PPB-150-100-PL, Hunter, Australia) and resuspended with another 2 mL of the corresponding culture media: HSA or PF. Each suspension then underwent a final wash, at 500 g for 5 min. The supernatant was removed and the pellet resuspended in 0.3 mL of HSA or PF media and the suspension was examined microscopically at 200 \times (phase-contrast, 20 \times objective, 10 \times ocular) (#BX50F, Olympus, Australia) and resulting sperm motility and concentration then manually assessed.

Following the trial preparation described above, 20 of the duplicate preparations were cultured in Falcon tubes (#352095, Falcon Corning, USA) overnight at room temperature. Twenty-four hours later, sperm motility and the calculated motility maintenance (=final progressive motility*100% initial progressive motility) of the sperm in each culture media were manually assessed.

2.5. Exposure to anti-static measures

Following trial preparation, the HSA and PF suspensions of 11 samples were manually assessed for sperm motility on microscope slides (Knittel, Germany) that had been subjected to antistatic treatments in the 5 min preceding the motility assessment, namely (i) wiped slides had been treated with an anti-static Superfine Fibre Cloth (GGs Photographic Equipment Company, China), and (ii) slides had been treated with Distinct Anti-Static Screen Cleaning fluid (E3! Style Product Specialists, Brisbane, Australia).

2.6. Statistical analysis

The differences between the experimental and control groups were compared for statistical significance with the Paired Student's *t*-test using the StatistXL (StatistXL, Western Australia, 2009) add-in to Excel (Microsoft Office, Microsoft Corporation, 2010). Differences were considered significant if $P < 0.05$.

3. Results

3.1. Initial density gradient preparations sperm culture

The overall results of the semen preparation using HSA-supplemented and PF media for all 41 samples are shown in Table 1. Analysis of the total number of sperm recovered from each media suite showed that PF preparation resulted in significantly higher numbers of sperm compared with the HSA suite ($P < 0.05$) and consequently a higher yield from the semen ($P < 0.05$). A slight but significant improvement in sperm morphology was noted for both gradients compared to the original semen (both $P < 0.05$) although they did not differ between themselves. Compared to the semen, both gradients resulted in higher proportions of sperm with progressive (A + B)

Table 1

The preparation of washed sperm by discontinuous gradients with suites of media either containing human serum albumin (HSA media; Medicult medium and Puresperm) or being protein-free (PF media; CellCura media and gradient). Values are mean \pm sem for 41 samples from 41 men.

Sperm parameter	Original semen	Final preparation	
		HSA media	PF media
Sperm concentration ($\times 10^6$ /mL)	87.8 \pm 10.2	–	–
Total sperm loaded/recovered ($\times 10^6$)	77.2 \pm 9.0	14.9 \pm 2.1	23.1 \pm 2.8*
Normal forms (%)	10.7 \pm 1.3	13.4 \pm 2.7	13.6 \pm 2.3
Motility (%)	57.2 \pm 2.2	83.9 \pm 2.6	71.3 \pm 2.4*
A + B	7.6 \pm 0.7	8.1 \pm 1.3	24.1 \pm 2.2*
C	34.3 \pm 1.9	8.4 \pm 2.3	5.1 \pm 1.2
D	–	20.5 \pm 2.6	31.7 \pm 4.0*
Yield (%)	–	–	–

* $P < 0.05$ between the HSA and protein-free medium.

Table 2

The motility and motility maintenance (mean \pm SEM) of washed sperm cultured for 24 h following preparation in HSA-containing and protein-free media. A total of 20 samples from 20 men were processed, and sperm survival in culture was defined as the final motility \times 100/initial motility.

Sperm parameter	HSA media		PF media	
	0 h	24 h	0 h	24 h
Motility: A + B (%)	83.9 \pm 2.6	57.3 \pm 4.9*	71.3 \pm 2.4	39.6 \pm 5.9*
C (%)	8.1 \pm 1.3	16.6 \pm 1.9*	24.1 \pm 2.2	35.5 \pm 4.5*
D (%)	8.4 \pm 2.3	25.8 \pm 4.1*	5.1 \pm 2.5	26.6 \pm 5.5*
Survival (%) ^a	–	68.0 \pm 5.3	–	57.7 \pm 7.5

*Results for the same preparation method after 24 h in culture are significantly different ($P < 0.05$).

^a Survival = final motility \times 100/initial progressive motility (motility maintenance).

motility and fewer immotile (D) sperm (both $P < 0.05$). However, there did appear to be a difference in the motility profiles of the sperm recovered from the two different media suites, with PF suspensions demonstrating a lower proportion of sperm with “A + B” type motility and a greater proportion of “C” type motile sperm. The proportion of immotile sperm was not significantly different between the two preparation types.

A subset of 20 prepared pairs of samples were cultured at room temperature overnight and the motility and percentage sperm motility maintenance were assessed as shown in Table 2. The proportion of immotile sperm was the same for both systems, but once again the proportion of “A + B” motile sperm at 24 h was greater in those suspensions prepared using the HSA suite of media whilst there was more “C” type motility observed in the PF suspensions. Whilst the survival of sperm seemed to be superior in the HSA suspensions, this difference was not statistically significant.

3.2. Factors affecting sperm motility profile

The increased proportion of sperm showing “C” grade motility in the PF medium, as described above, was eventually noticed to be associated with sperm sticking to the glass of the Makler chamber. In an attempt to explain this, sperm motility was further assessed in a sub-set of 9 sperm suspensions using the SQA II automated analyser as shown in Table 3. Despite the samples processed using the PF gradient again showing the same pattern of lower progressive and higher non-progressive motility than the HSA gradients preparations when manually assessed on the Makler chamber, the SQA demonstrated no significant difference in the SMI between the two media suites ($P = 0.83$).

Table 3

The motility (mean \pm SEM) of sperm prepared using HSA-containing and protein-free media when assessed both manually and with the Sperm Quality Analyser ($n = 9$ semen samples from 9 men).

Motility assessment (%)		Media suite	
		HSA	Protein-free
Manual	A + B	90.8 \pm 2.9	73.1 \pm 4.4*
	C	4.9 \pm 1.9	23.2 \pm 3.3*
	D	4.2 \pm 1.9	3.8 \pm 1.6
SQA analyser	Sperm motility index	452.4 \pm 43.9	470.1 \pm 36.8

* $P < 0.05$ between the HSA and protein-free medium.

Table 4

Motility of sperm (mean \pm SEM) prepared using HSA-containing and protein-free media on slides when assessed on microscope slides pre-treated with anti-static spray or wipes ($n = 11$).

Motility grade	Spray pre-treatment		Wipe pre-treatment	
	HSA	Protein-free	HSA	Protein-free
A + B (%)	53.1 \pm 3.4	41.0 \pm 4.2*	55.3 \pm 3.6	43.0 \pm 4.4
C (%)	36.0 \pm 2.9	45.8 \pm 4.4	36.0 \pm 3.1	49.8 \pm 3.8*
D (%)	9.8 \pm 2.1	11.1 \pm 3.1	9.6 \pm 1.2	7.3 \pm 1.6

*Significant $P < 0.05$.

To endeavour to determine whether static electricity may have caused adherence of sperm to the glass microscope slide during motility assessment, and hence an apparent increase in “C” type motility, the slides used to assess 11 pairs of washed preparations were exposed to 2 antistatic measures as shown in Table 4. The results indicate that the antistatic spray and wipes both increased the proportion of sperm with “C” type motility in the HSA-containing medium, and did not improve the situation in the PF medium.

Sperm motility in PF medium was opportunistically noticed to improve in PF medium when viewed on a microscope slide left on the bench. In a final attempt to ensure the difference in motility patterns between the two different media was not a simple artefact, motility was manually assessed at 0, 3 and 6 min after loading of the glass slides in a subset of 11 samples as shown in Table 5. While statistically significant differences between patterns of sperm motility are apparent in the different media types at the early time intervals, these differences appear to resolve over time. Samples therefore processed on the two sets of gradients and assessed 6 min after loading on to the Makler chamber, as shown in Figure 1, were similar for both media types.

4. Discussion

The use of serum to supplement the culture of mammalian embryos dates back to the 1960s when it was discovered that the addition of rabbit or bovine serum to culture media significantly increased the proportion of blastocysts formed [28], and the early work on human IVF used patient serum as a supplement to the medium [29]. Serum is thought to help create an environment that more closely resembles that of the female reproductive tract that may improve the rate of cleavage, encourage activation of the embryonic genome and enhance the embryos' implantation potential [30], and because of its complex composition, human serum added to culture media can act as a buffer, helping to minimise fluctuations in pH, whilst the amino acids contained within it may supplement the activity of those amino acids

Table 5

The assessment of motility (mean \pm SEM) at different times after loading on to a microscope slide. Sperm from 11 ejaculates produced by 11 men were prepared and cultured in both HSA-containing and protein-free media.

Time (mins)	HSA media			PF media		
	A + B	C	D	A + B	C	D
0	84.7 \pm 3.7	7.5 \pm 2.4	7.9 \pm 3.5	74.9 \pm 2.7	21.1 \pm 3.0*	3.6 \pm 1.0
3	85.6 \pm 4.1	8.1 \pm 2.6	7.1 \pm 3.7	71.0 \pm 3.4*	25.4 \pm 3.6*	3.6 \pm 1.2
6	85.2 \pm 3.5	7.0 \pm 2.1	8.3 \pm 3.2	86.8 \pm 2.0	9.4 \pm 1.9	3.4 \pm 0.9

* $P < 0.05$ comparing HSA-containing and protein-free media.




Figure 1. The motility of sperm from 11 semen samples prepared using density gradients and resuspended in medium either containing human serum albumin (HSA) or being protein-free (PF). Sperm motility was assessed 6 min after loading on to the microscope slide.

already separately added to culture media and it may also help bind any heavy metals [31]. In addition to its effect on embryos, the presence of serum appears to facilitate the process of capacitation and/or the acrosome reaction of sperm, which is essential to enable them to fertilise an oocyte *in vitro* [32,33]. Serum in culture also appears to reduce the incidence of DNA damage in sperm and increase the rate of cleavage in human embryos [34]. At a practical level, serum also prevents gametes and embryos from becoming sticky and difficult to handle in culture, thus facilitating their manipulation in culture and preventing them from adhering to the plastic and glassware used [35].

The use of serum has been criticised because, being a biological solution, its composition is difficult to define, can be subject to significant variation, and may include components that impact on embryonic development such as hormones, vitamins, fatty acids and ions [14,36]. When used as a supplement in assisted reproductive technology, there is the compounding problem that the maternal serum from some women, such as those with endometriosis, may contain elements that are embryotoxic [37]. Perhaps more importantly, the use of serum, both from animal and human origins, has also been criticised because of the risk of disease transmission. In spite of donor screening processes, in 1991 in the Netherlands, 79 IVF patients contracted Hepatitis B following the culture of their embryos in culture media containing pooled serum that had been contaminated by the Hepatitis B virus [12] and, more recently, it was revealed that a commercially-available HSA product had potentially been contaminated following the death of an albumin donor due to the infectious Creutzfeldt–Jakob Disease (CJD) prion [13].

In the quest to remove serum from IVF culture medium, a version of medium without serum has been used [38]. However, human serum albumin in one form or another, but still derived from blood, is predominantly used to supplement ART culture media in the hope that it provides a better defined medium

and elicits improved results [15,30,39–41], although there is some concern that serum supplementation yields better results than protein supplementation alone and the latter may be suitable only in conjunction with additional components [42]. Recombinant albumin has been used in cell culture because of the concerns over transmissible spongiform encephalopathies [2], and it has also been used in a general IVF setting [43] as well as the special case of Jehovah's witnesses [44]. Nevertheless, the use of non-protein macromolecules has been investigated for use as a supplement for ART culture media in an attempt to avoid the problems associated with blood products, such as the addition of polyvinyl alcohol (PVA) [45], polyvinylpyrrolidone (PVP) [46] and dextran [18] to aid with the potential handling differences when HSA is not incorporated into culture media. However, the teratological safety of the likes of PVA and PVP is uncertain [47].

Methylcellulose has been shown to be a suitable macromolecule to support gametes and embryos [22,23] and following patenting of a formulation [21] this has now been placed in to commercial production and marketed by CellCura®. The availability of such a medium would appear to be a step forward but a recent clinical trial had to be cancelled due to poor performance of the medium [25]. It was unclear whether this was due to issues with the formulation, manufacture or shipment, but the satisfactory performance of the formulation reported earlier [22–24] cannot be ignored. The usefulness of the medium is best confirmed by a systematic evaluation of the methylcellulose-supplemented medium in the key steps of the ART process, rather than just an overview of the overall process. The aim of the present study, therefore, was to independently assess the effectiveness of the medium using density gradients, a routine sperm preparation technique employed by many clinics worldwide, in terms of sperm yield, motility, survival and morphology. One suite of media used was supplemented traditionally with HSA; the other with the plant-derived macromolecule, methylcellulose.

The results of the present study confirmed that the yield of sperm using the PF suite of gradients and medium was satisfactory. However, the main limitation was the artefact in assessing sperm motility. There was an apparent increase in the proportion of non-progressively motile sperm but this was not apparent with the deep chamber used with the SQA sperm analyser, and resolved itself on glass slides by simply waiting for 6 min and allowing the sperm to detach themselves from the glass surface. As mentioned above, the “stickiness” of cells is relieved by the inclusion of protein [35], but equally a revised protocol using the PF medium was effective. However, regulatory authorities (e.g. the Therapeutic Goods Administration, TGA, in Australia) require such a modification to be documented in the manufacturer's Instructions for Use (see <https://www.tga.gov.au/publication/australian-regulatory-guidelines-medical-devices-argmd>) and this was not provided. The use of the suite of PF media under sub-optimal conditions will bias the findings of laboratories and may result in erroneous conclusions on the suitability of the media for clinical use. These findings raise the question of whether modifications are required at other stages of gamete culture, and so further systematic studies on different aspects of use of the media with sperm are warranted.

In summary, the present study has confirmed that the use of a commercially-available protein-free suite of media can result in good preparations of washed sperm, but that a modified protocol

must be used to assess sperm motility in order to avoid the artefact of sperm sticking to the microscope observation chamber. While these results support the use of methylcellulose as an alternative to HSA, albeit it with a modified protocol, further studies on the functional capacity of the sperm are required.

Declaration of interest statement

The authors KP, PR, and PM declare no competing interests. JA is a scientific consultant to CellCura and the inventor of the PF medium.

Acknowledgements

We gratefully acknowledge the assistance of Kailin Yap, Melissa Stemp, Katie Feenan, Clare Deuchar, Yanhe Liu and Bridget Blackwell in the recruitment of patients to this study and blinding of samples.

References

- [1] Castle P, Robertson JS. Animal Sera, Animal Sera derivatives and substitutes used in the manufacture of pharmaceuticals. *Biologicals* 1998; **26**(4): 365-368.
- [2] Keenan J, Pearson D, Clynes M. The role of recombinant proteins in the development of serum-free media. *Cytotechnology* 2006; **50**(1–3): 49-56.
- [3] Groebe K, Hayess K, Klemm-Manns M, Schwall G, Wozny W, Steemans M, et al. Protein biomarkers for *in vitro* testing of embryotoxicity. *J Proteome Res* 2010; **9**(11): 5727-5738.
- [4] Macaldowie A, Wang YA, Ghughtai A, Chambers GM. *Assisted reproductive technology in Australia and New Zealand 2012*. Sydney: National Perinatal Epidemiology and Statistics Unit, the University of New South Wales; 2014.
- [5] Green M, Mouat F, Miles H, Pleasants A, Henderson H, Hofman P, et al. The phenotype of an IVF child is associated with periconception measures of follicular characteristics and embryo quality. *Hum Reprod* 2014; **29**(11): 2583-2591.
- [6] Green MP, Mouat F, Miles HL, Hopkins SA, Derraik JGB, Hofman PL, et al. Phenotypic differences in children conceived from fresh and thawed embryos in *in vitro* fertilization compared with naturally conceived children. *Fertil Steril* 2013; **99**(7): 1898-1904.
- [7] Dumoulin JC, Land JA, Van Montfoort AP, Nelissen EC, Coonen E, Derhaag JG, et al. Effect of *in vitro* culture of human embryos on birthweight of newborns. *Hum Reprod* 2010; **25**(3): 605-612.
- [8] Nelissen ECM, Van Montfoort APA, Smits LJM, Menheere PPCA, Evers JLH, Coonen E, et al. IVF culture medium affects human intrauterine growth as early as the second trimester of pregnancy. *Hum Reprod* 2013; **28**(8): 2067-2074.
- [9] Chen M, Wu L, Zhao J, Wu F, Davies MJ, Wittert GA, et al. Altered glucose metabolism in mouse and humans conceived by *in vitro* fertilization (IVF). *Diabetes* 2014; **63**(10): 3189-3198.
- [10] Chronopoulou E, Harper JC. IVF culture media: past, present and future. *Hum Reprod Update* 2014; **21**(1): 39-55; <http://dx.doi.org/10.1093/humupd/dmu040>.
- [11] Blake D, Svalander P, Jin M, Silversand C, Hamberger L. Protein supplementation of human IVF culture media. *J Assist Reprod Genet* 2002; **19**(3): 137-143.
- [12] van Os HC, Drogendijk AC, Fetter WP, Heijntink RA, Zeilmaker GH. The influence of contamination of culture medium with hepatitis B virus on the outcome of *in vitro* fertilization pregnancies. *Am J Obstet Gynecol* 1991; **165**(1): 152-159.
- [13] Kemmann E. Creutzfeldt-Jakob disease (CJD) and assisted reproductive technology (ART). Quantification of risks as part of informed consent. *Hum Reprod* 1998; **13**(7): 1777.

- [14] Barnes D, Sato G. Methods for growth of cultured cells in serum-free medium. *Anal Biochem* 1980; **102**(2): 255-270.
- [15] Laverge H, De Sutter P, Desmet R, Van der Elst J, Dhont M. Prospective randomized study comparing human serum albumin with fetal cord serum as protein supplement in culture medium for in-vitro fertilization. *Hum Reprod* 1997; **12**(10): 2263-2266.
- [16] Dyrlund TF, Kirkegaard K, Poulsen ET, Sanggaard KW, Hindkjær JJ, Kjems J, et al. Unconditioned commercial embryo culture media contain a large variety of non-declared proteins: a comprehensive proteomics analysis. *Hum Reprod* 2014; **29**(11): 2421-2430.
- [17] Morbeck D, Paczkowski M, Fredrickson J, Krisher R, Hoff H, Baumann N, et al. Composition of protein supplements used for human embryo culture. *J Assist Reprod Genet* 2014; **31**(12): 1703-1711.
- [18] Matson P, Tardif S. A preliminary search for alternatives to albumin as a medium supplement for the culture of human sperm. *Reprod Biol* 2012; **12**(3): 329-331.
- [19] Bavister B. Substitution of a synthetic polymer for protein in a mammalian gamete culture system. *Dev Biol* 1981; **217**(1): 45-51.
- [20] Bavister BD. Culture of preimplantation embryos: facts and artifacts. *Hum Reprod Update* 1995; **1**(2): 91-148.
- [21] Ali Bin, Abudallah Jaffar M, Cellcura, Inc., assignee. Protein-free gamete and embryo handling and culture media products. 13/799400 (patent) 2013.
- [22] Ali J. Generation of viable human embryos in a protein-free embryo culture (ART-7b) medium enhances clinical pregnancy rate and prevents disease transmission in assisted reproduction. *Middle East Fertil Soc J* 2004; **9**(2): 118-127.
- [23] Ali J, Shahata MA, Al-Natsha SD. Formulation of a protein-free medium for human assisted reproduction. *Hum Reprod* 2000; **15**(1): 145-156.
- [24] Ali J. Investigation into the nutrient requirement of human embryos: successful formulation and clinical trial of a novel protein-free embryo culture medium. *Emir Med J* 2000; **18**: 195-202.
- [25] Devroe J, Debrock S, Afschrift H, D'Hooghe T, Spiessens C. Protein-free culture medium has a negative effect on embryo quality: data analysis of a cancelled randomized trial. *Hum Reprod* 2014; **29**(Supp 1): i3.
- [26] WHO. *World Health Organisation laboratory manual for the examination of human semen and sperm-cervical mucus interaction*. Cambridge: Cambridge University Press; 1999.
- [27] Matson PL, Webb S, Kappelle W, Clarke BR. The benefits and limitations of the sperm quality analyzer IIB in assessing porcine sperm motility. *Arch Androl* 2007; **53**(6): 359-365.
- [28] Onuma H, Maurer RR, Foote RH. In-vitro culture of rabbit ova from early cleavage stages to the blastocyst stage. *J Reprod Fertil* 1968; **16**(3): 491-493.
- [29] Edwards RG, Steptoe PC. Current status of in-vitro fertilisation and implantation of human embryos. *Lancet* 1983; **322**(8362): 1265-1269.
- [30] Desai NN, Sheean LA, Martin D, Gindlesperger V, Austin CM, Lisbonna H, et al. Clinical experience with synthetic serum substitute as a protein supplement in IVF culture media: a retrospective study. *J Assist Reprod Genet* 1996; **13**(1): 23-31.
- [31] Trounson A, Gardner DK. *Handbook of in vitro fertilization*. Boca Raton, FL, USA: CRC Press; 1993.
- [32] Dow MP, Bavister BD. Direct contact is required between serum albumin and hamster spermatozoa for capacitation in vitro. *Gamete Res* 1989; **23**(2): 171-180.
- [33] Fraser LR. Albumin is required to support the acrosome reaction but not capacitation in mouse spermatozoa in vitro. *J Reprod Fertil* 1985; **74**(1): 185-196.
- [34] Aitken RJ, Clarkson JS. Significance of reactive oxygen species and antioxidants in defining the efficacy of sperm preparation techniques. *J Androl* 1988; **9**(6): 367-376.
- [35] Jones R, Mann T, Sherins R. Peroxidative breakdown of phospholipids in human spermatozoa, spermicidal properties of fatty acid peroxides, and protective action of seminal plasma. *Fertil Steril* 1979; **31**(5): 531-537.
- [36] Barnes D, Sato G. Serum-free cell culture: a unifying approach. *Cell* 1980; **22**(3): 649-655.
- [37] Ito F, Fujino Y, Ogita S. Serum from endometriosis patients impairs the development of mouse embryos in vitro—comparison with serum from tubal obstruction patient and plasmanate. *Acta Obstet Gynecol Scand* 1996; **75**(10): 877-880.
- [38] Caro CM, Trounson A. Successful fertilization, embryo development, and pregnancy in human in vitro fertilization (IVF) using a chemically defined culture medium containing no protein. *J In Vitro Fert Embryo Transf* 1986; **3**(4): 215-217.
- [39] Adler A, Reing AM, Bedford JM, Alikani M, Cohen J. Plasmanate as a medium supplement for in vitro fertilization. *J Assist Reprod Genet* 1993; **10**(1): 67-71.
- [40] Ashwood-Smith MJ, Hollands P, Edwards RG. The use of albumin 5 as a medium supplement in clinical IVF. *Hum Reprod* 1989; **4**(6): 702-705.
- [41] Weathersbee PS, Pool TB, Ord T. Synthetic serum substitute (SSS): a globulin-enriched protein supplement for human embryo culture. *J Assist Reprod Genet* 1995; **12**(6): 354-360.
- [42] Hargreaves C, Rahman F, Cowan D, Santis M, Keefe T, Howell R, et al. Serum is more effective than albumin in promoting human embryo development and implantation. *Fertil Steril* 1995; **64**(6): 1162-1166.
- [43] Bungum M, Humaidan P, Bungum L. Recombinant human albumin as protein source in culture media used for IVF: a prospective randomized study. *Reprod Biomed Online* 2002; **4**(3): 233-236.
- [44] Grubb J, Muramoto O, Matson P. Issues arising during the treatment of Jehovah's Witnesses by in vitro fertilisation. *Hum Fertil* 2011; **14**(1): 35-40.
- [45] Biggers JD, Summers MC, McGinnis LK. Polyvinyl alcohol and amino acids as substitutes for bovine serum albumin in culture media for mouse preimplantation embryos. *Hum Reprod Update* 1997; **3**(2): 125-135.
- [46] Cholewa JA, Whitten WK. Development of two-cell mouse embryos in the absence of a fixed-nitrogen source. *J Reprod Fertil* 1970; **22**(3): 553-555.
- [47] Gardner D, Lane M. Culture of viable human blastocysts in defined sequential serum-free media. *Hum Reprod* 1998; **13**(Suppl 3): 148-159.