

1-1-2023

Sex differences in skeletal muscle fiber types: A meta-analysis

James L. Nuzzo
Edith Cowan University

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworks2022-2026>



Part of the [Medicine and Health Sciences Commons](#)

[10.1002/ca.24091](https://doi.org/10.1002/ca.24091)

Nuzzo, J. L. (2023). Sex differences in skeletal muscle fiber types: A meta-analysis. *Clinical Anatomy*, 37(1), 81-91.

<https://doi.org/10.1002/ca.24091>

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworks2022-2026/2809>

REVIEW

CLINICAL
ANATOMY

WILEY

Sex differences in skeletal muscle fiber types: A meta-analysis

James L. Nuzzo School of Medical and Health Sciences, Edith
Cowan University, Joondalup, Australia

Correspondence

James L. Nuzzo, School of Medical and Health
Sciences, Edith Cowan University,
270 Joondalup Drive, Joondalup WA 6027,
Australia.
Email: j.nuzzo@ecu.edu.au

Abstract

Biopsies have been acquired from living men and women to determine proportions of Type I (slow-twitch) and II (fast-twitch) skeletal muscle fibers since the 1970s. Sex differences have been assumed but the literature has not been submitted to meta-analysis. Here, the aim was to generate effect sizes of sex differences in muscle fiber cross-sectional areas, distribution percentages, and area percentages. Data from 2875 men and 2452 women, who participated in 110 studies, were analyzed. Myofibrillar adenosine triphosphatase histochemistry was used in 71.8% of studies to classify fibers as Type I, II, IIA, and/or IIX; immunohistochemistry, immunofluorescence, or sodium dodecyl sulfate-polyacrylamide gel electrophoresis were used in 35.4% of studies to similarly classify myosin heavy chain (MHC) isoform content. Most studies involved biopsies from vastus lateralis (79.1%) in healthy individuals (92.7%) between 18 and 59 years old (80.9%). Men exhibited greater cross-sectional areas for all fiber types ($g = 0.40$ – 1.68); greater distribution percentages for Type II, MHC II, IIA, IIX fibers ($g = 0.26$ – 0.34); greater area percentages for Type II, IIA, MHC IIA, IIX fibers ($g = 0.39$ – 0.93); greater Type II/I and Type IIA/I fiber area ratios ($g = 0.63$, 0.94). Women exhibited greater Type I and MHC I distribution percentages ($g = -0.13$, -0.44); greater Type I and MHC I area percentages ($g = -0.53$, -0.69); greater Type I/II fiber area ratios ($g = -1.24$). These data, which represent the largest repository of comparative muscle fiber type data from living men and women, can inform discussions about biological sex and its impact on pathologies and sports performance (e.g., explaining sex differences in muscle strength and muscle endurance).

KEYWORDS

anatomy, biopsy, men, muscles, physiology, quadriceps muscle, sex, women

1 | INTRODUCTION

Researchers have biopsied skeletal muscles from living men and women to determine muscle fiber types since the 1970s. In early studies, myofibrillar adenosine triphosphatase (mATPase) histochemistry was used to classify fibers as Type I ("slow twitch") or Type II ("fast twitch"), with additional subtypes established for Type II fibers

(e.g., Type IIA, Type IIX [called Type IIB in some papers]). Later, immunohistochemistry and sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) were used to examine myosin heavy chain (MHC) isoform content, with strong correlations between proportional representations of MHC I, MHC IIA, and MHC IIX isoforms and proportional representations of Type I, Type IIA, and Type IIX fibers, respectively (Chalmers & Row, 2011; Damer et al., 2022; Fry

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Clinical Anatomy* published by Wiley Periodicals LLC on behalf of American Association of Clinical Anatomists and British Association of Clinical Anatomists.

et al., 1994; Scott et al., 2001; Staron, 1991). Because Type II fibers shorten at faster velocities and produce more force and power than Type I fibers, and because Type I fibers possess a greater capacity for oxidative energy production and are more enduring than Type II fibers (Bottinelli et al., 1996; 1999; Essén et al., 1975; Frontera et al., 2000; Garnett et al., 1979; Krivickas et al., 2001; 2006; Larsson et al., 1997; Stienen et al., 1996; Trappe et al., 2003; Widrick et al., 2002), a sex difference in muscle fiber type has implications for discussions about biological sex and its impacts on neuromuscular physiology, sport performance, and disease pathology. Differences in muscle fiber types might help to explain how men produce more force than women, as greater muscle mass in men than women does not appear to fully explain the sex difference in muscle strength (Nuzzo, 2023). A sex difference in muscle fiber types might also help to explain why women are more resistant to muscle fatigue than men are in certain exercise tasks (Hunter, 2016a; Nuzzo, 2023). Biological causes for sex differences in neuromuscular performance then have implications for discussions about transgender women (i.e., biological men) competing against biological women in the female category of sport (Hilton & Lundberg, 2021; Nokoff et al., 2023). A study of sex differences in muscle fiber types might also enhance understanding of potential sex differences in susceptibilities to the effects of aging (Doherty, 2001; Lexell, 1995), disease (Ciciliot et al., 2013; Talbot & Maves, 2016), and exercise participation (Straight et al., 2020).

In a recent narrative review, Nuzzo (2023) collated data from 34 studies that reported on skeletal muscle fiber types in men and women. Nuzzo (2023) concluded that men and women have similar proportional numbers (i.e., distribution percentages) of Type I and Type II fibers but that Type II fibers occupy a greater proportional area (i.e., area percentage) of muscle in men than women. Moreover, Type I muscle fibers occupy a greater area percentage of muscle in women, leading to higher Type II/I fiber area ratios in men than women (Nuzzo, 2023). Hunter (2016a) reached a similar conclusion after examining results from 13 studies. Nevertheless, the reviews by Nuzzo (2023) and Hunter (2016a) were narrative, and they were not concentrated on the topic of sex differences in muscle fiber types. Consequently, the literature searches were not exhaustive, and only mean values, without consideration of measurement variability and effect sizes, were presented (Hunter, 2016a; Nuzzo, 2023).

Therefore, the purpose of the current study was to use meta-analysis to generate effect sizes of sex differences in cross-sectional areas, distribution percentages, and area percentages of various skeletal muscle fiber types as determined by biopsy in living humans. Such information has potential to clarify differences in neuromuscular anatomy and physiology between men and women and thus inform ongoing discussions about biological sex and its impacts on neuromuscular physiology, sport performance, and disease pathology (Hilton & Lundberg, 2021; Hunter, 2014; 2016a; 2016b; Jones et al., 2021; Nokoff et al., 2023; Nuzzo, 2023; Roberts et al., 2020; Vinciguerra et al., 2023). Such information also meets calls for more sex-segregated data in biomedical research (Schilaty et al., 2018; Smith, McKay, Ackerman, et al., 2022; Smith, McKay, Kuikman, et al., 2022).

2 | METHODS


2.1 | Literature search

Thirty-four papers with relevant data were already known to the author based on a previous narrative review (Nuzzo, 2023). However, an updated and more exhaustive literature search was performed in March of 2023. The search involved a mixed approach that was similar to that described by Greenhalgh and Peacock (2005). It relied on the investigator's personal knowledge from previous research on the topic (Nuzzo, 2023), checking of personal digital files, relevant keyword searches in PubMed and Google Scholar, and "snowballing" strategies (i.e., reference and citation tracking). Keyword searches in PubMed and Google Scholar included various combinations of words such as, but not limited to, "biopsy," "fast twitch," "female," "fiber type," "male," "men," "muscle fiber," "sex," "slow twitch," and "women." No date restrictions were placed on the search. Tables of muscle fiber type data presented in other papers were also a focus of the search strategy (Cagnie et al., 2015; Chalmers & Row, 2011; Damer et al., 2022; Gouzi et al., 2013; Hunter, 2016a; Naruse et al., 2023; Ng et al., 1998; Nuzzo, 2023; Staron et al., 2000; Straight et al., 2020).

2.2 | Eligibility

A study was eligible for inclusion into the meta-analysis if the following conditions were met: (a) data were published in a journal paper; (b) the paper was published in English; (c) muscle biopsies were performed on living humans; (d) muscle fiber percent distribution or percent area data were reported; and (e) mean values of muscle fiber percent distribution or percent area were reported for both men and women. Common reasons for exclusion included: data acquired at autopsy; data not segregated by sex or acquired in only men or only women; or no data on muscle fiber percent distribution or percent area reported. Also, the following papers were excluded due to duplication of data: (a) the paper by Bell and Jacobs (1989) was excluded because the authors presented the same data elsewhere (Bell & Jacobs, 1990) (included); (b) the paper by Froese and Houston (1987) was excluded because the authors presented the same data elsewhere (Froese & Houston, 1985) (included); (c) a paper by Jaworowski et al. (2002) was excluded because the authors presented the same data elsewhere (Holmbäck et al., 2003) (included). A paper by Evertsen et al. (1999) was also excluded, because, although the authors reported no sex difference in muscle fiber distribution percent of the vastus lateralis muscle in cross-country skiers, the authors did not present the sex-specific means. Papers that included means but no standard deviations (SD) or standard errors for the muscle fiber type outcomes of interest were still included in the research. The absence of SDs or standard errors prevented such data from contributing to the computed effect sizes. However, these data still contributed to the sex-specific weighted means for muscle fiber type outcomes. A

FIGURE 1 Flow diagram of search strategy and exclusion process.



flow diagram of the search strategy and exclusion process is presented in Figure 1.

2.3 | Data extraction

Data that were extracted from eligible papers included author name, study year, country, sample size, age, health status (i.e., healthy or patient group), muscle biopsied, method used to analyze the biopsies, and means and SDs for fiber cross-sectional area, fiber distribution percent, fiber area percent, and fiber area ratios. For papers that presented muscle fiber type data before and after interventions, only measurements from baseline were extracted. When authors reported fiber size as diameter rather than cross-sectional area, the data were converted to area using an online calculator (<https://www.omnicalculator.com/math/area-of-a-circle>) ($\text{area} = \pi \times r^2$). When

authors reported variance as standard error, the data were converted to SD by multiplying the standard error by the square root of the sample size. When authors presented data in figures, data were extracted using a graph digitizer (WebPlotDigitizer, <https://apps.automeris.io/wpd/>).

Classification of muscle fiber types has evolved over time (Chalmers & Row, 2011; Damer et al., 2022; Fry et al., 1994; Scott et al., 2001; Staron, 1991). Thus, because researchers have reported muscle fiber type data using different taxonomies, a decision was made by the author to extract data of the fiber type classifications reported most commonly across the entirety of the relevant literature. The extracted and analyzed muscle fiber type classifications were: Type I, Type II, Type IIA, Type IIX, MHC I, MHC II, MHC IIA, and MHC IIX. Of note, results from mATPase histochemistry and MHC isoform methods correlate (e.g., Type I vs MHC I) (Fry et al., 1994; Staron, 1991). Thus, in the current paper, Type I and MHC I fibers can

be considered Type I slow twitch fibers, Type IIX and MHC IIX fibers can be considered Type II fast twitch fibers, and Type IIA and MHC IIA can be considered Type II intermediate fast twitch fibers. The Type I and MHC data that corresponded to each other (i.e., Type I and MHC I) were not combined in the current analysis, because, although their correlations are strong (Fry et al., 1994; Staron, 1991), they not perfect, and it was clear that enough data were available from both types of outcomes that they could be analyzed separately.

2.4 | Statistical analysis

Version 29 of the Statistical Software Package for the Social Sciences (SPSS, Armonk, US) was used to analyze the data and create plots for the meta-analysis. Random effects meta-analyses were used to generate effect sizes (Hedges *g*) with 95% confidence intervals (CI) and prediction intervals (Inthout et al., 2016) for sex differences in muscle fiber types. Separate meta-analyses were performed for cross-sectional areas, distribution percentages, area percentages for Type I, Type II, Type IIA, Type IIX, MHC I, MHC II, MHC IIA, and MHC IIX muscle fibers. Meta-analyses were also performed for Type I/II, Type II/I, and Type IIA/I fiber area ratios. Forest plots were used to graphically display the results from the meta-analyses. Each Forest plot displayed the effect sizes of individuals studies and the overall effect size and 95% CI. Effect sizes equal to 0.2, 0.5, and 0.8 are often considered small, moderate, and large, respectively; however, such benchmarks are arbitrary and should not be interpreted rigidly (Lakens, 2013). Confidence intervals not crossing zero indicate overall effects that are statistically significant (i.e., $p \leq 0.05$) (Cumming, 2009). Also, in the Forest plots, when applicable, data were separated by age group (< 18, 18–59, and > 59 years). The age group-specific effect sizes were presented along with the overall effect size across all individuals. The overall effect size was the main statistic of interest, with age group-specific effect sizes presented as preliminary results. Funnel plots were generated to explore potential publication and other bias. Tests of heterogeneity were also conducted. Sex-specific weighted means for fiber cross sectional areas, distribution percentages, area percentages, and fiber area ratios were also computed. All of the data and results associated with the research are available either in the text, Appendix S1, or at the Open Science Framework (<https://osf.io/m97z3/>).

3 | RESULTS

3.1 | Study characteristics

A total of 110 studies, which ranged in publication year from 1976 to 2022, were included in the analysis (see Appendix S1). Four studies (Costill et al., 1976; Nygaard et al., 1983; Rolf et al., 1997; Sjøgaard, 1982) did not contribute effects for any of the outcomes because the authors reported only means and no SDs or standard errors. Thus, 106 studies contributed to one or more of the effect

sizes in the meta-analysis. Some studies included multiple groups of men and women and thus contributed multiple effects to the analysis. All 110 studies contributed to one or more of the weighted means. The 110 studies included data from 2875 men and 2452 women, who comprised 157 study groups. The means \pm SD of the sample sizes for men and women were 26.1 ± 32.9 (minimum = 4; maximum = 215) and 22.3 ± 27.5 (minimum = 4; maximum = 203), respectively.

Table 1 presents the number of studies by country, age group, health status, and muscle biopsied. Most studies were conducted in Sweden ($n = 27$, 24.5%) and the United States ($n = 27$, 24.5%). The vast majority of studies included participants who were healthy ($n = 102$, 92.7%) and 18–59 years old ($n = 89$, 80.9%). Athletes were participants in 11.8% of studies. The vastus lateralis was the muscle most frequently biopsied ($n = 87$, 79.1%). mATPase was the method used most frequently to determine muscle fiber type ($n = 79$, 71.8%).

3.2 | Muscle fiber type variables

All Forrest plots from the meta-analysis are provided in the supplementary figures (Figures S1–S26). Table 2 provides a summary of the findings from the meta-analyses for each muscle fiber type outcome (i.e., number of effects, Hedges *g*, 95% CIs, and 95% predictions intervals). Table 2 also displays the weighted means each of the muscle fiber type outcomes. Muscle-specific weighted means for Type I distribution percentages are presented in Table 3. For vastus lateralis, weighted means for other muscle fiber type outcomes are also presented in Table 3.

Men exhibited larger cross-sectional areas for Type I, Type II, Type IIA, Type IIX, MHC I, MHC II, MHC IIA, and MHC IIX muscle fibers (Figures S1–S8). Men exhibited greater distribution percentages for Type II, MHC II, MHC IIA, and MHC IIX muscle fibers (Figures S10, S14–S16). Women exhibited greater distribution percentages for Type I and MHC I muscle fibers (Figures S9, S13) and there were no sex differences in distribution percentages for Type IIA and Type IIX muscle fibers (Figures S11, S12). Men exhibited greater area percentages for Type II, Type IIA, MHC IIA, and MHC IIX muscle fibers (Figures S18, S19, S22, S23). Women exhibited greater area percentages for Type I and MHC I muscle fibers (Figures S17, S21) and there was no sex difference in area percentage for Type IIX muscle fibers (Figure S20). Men exhibited greater Type II/I and Type IIA/I fiber area ratios (Figures S25, S26). Women exhibited greater Type I/II fiber area ratios (Figure S24).

The figures in Appendix S1 also display effect sizes by age group. The results from this preliminary analysis appear to show that age and sex interact to impact the size of the sex difference for some muscle fiber type outcomes, most notably Type I distribution percent (girls < boys; women > men in middle age; women = men in older age) and Type IIA and Type IIX distribution percent (girls > boys; women < men in middle age; women = men in older age). When data from youth and older adults are excluded, a statistically significant sex difference is present for Type IIA distribution percentage (men > women, Figure S11).

TABLE 1 Characteristics of study ($n = 110$) methods and samples.

Category	<i>n</i>	% study total
Country		
Australia	3	2.7
Canada	15	13.6
Croatia	1	0.9
Denmark	9	8.2
Finland	4	3.6
France	2	1.8
Greece	2	1.8
Japan	2	1.8
South Korea	1	0.9
Netherlands	2	1.8
Norway	1	0.9
Russia	2	1.8
Spain	3	2.7
Sweden	27	24.5
Switzerland	4	3.6
United Kingdom	4	3.6
United States of America	27	24.5
Venezuela	1	0.9
Muscle		
Biceps brachii	6	5.5
Gastrocnemius medialis	2	1.8
Gastrocnemius lateralis	5	4.5
Longissimus/ES	7	6.4
Multifidus	5	4.5
Rectus femoris	2	1.8
Semitendinosus	1	0.9
Soleus	1	0.9
Tibialis anterior	2	1.8
Triceps brachii	2	1.8
Vastus lateralis	87	79.1
Method for fiber typing		
mATPase	79	71.8
MHC immunohistochemistry	16	14.5
MHC SDS-PAGE	21	19.1
MHC immunofluorescence	2	1.8
Age group		
Youth (<18 years)	7	6.4
Adult (18–59 years)	89	80.9
Older adult (≥60 years)	24	21.8
Health status		
Mixed group	1	0.9
Healthy	102	92.7
Non-athlete	89	80.9
Athlete	13	11.8
Bodybuilders	2	1.8

(Continues)

TABLE 1 (Continued)

Category	<i>n</i>	% study total
Dancers	2	1.8
Jumpers	1	0.9
Mixed group	1	0.9
Orienteers	1	0.9
Powerlifters	3	2.7
Runners—long distance	1	0.9
Runners—middle distance	2	1.8
Runners—sprinters	2	1.8
Speed skaters	1	0.9
Throwers—javelin	1	0.9
Throwers—shot, discus	1	0.9
Patients	20	18.2
Anorexia nervosa	1	0.9
Chronic heart failure	1	0.9
COPD	2	1.8
Diabetes	3	2.7
Low back pain	8	7.3
Multiple sclerosis	2	1.8
Obesity	3	2.7
Peripheral artery disease	1	0.9

Abbreviations: COPD, chronic obstructive pulmonary disease; mATPase, myofibrillar adenosine triphosphatase; MHC, myosin heavy chain; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

4 | DISCUSSION

The purpose of the current study was to use meta-analysis to generate effect sizes of sex differences in cross-sectional areas, distribution percentages, and area percentages of various skeletal muscle fiber types in living men and women. Data from 2875 men and 2452 in 110 studies were analyzed. The results reveal that men exhibit greater cross-sectional areas for all muscle fiber types and greater area percentages for Type II muscle fibers. Women exhibit greater distribution and area percentages for Type I fibers. Consequently, women exhibit greater Type I/II fiber area ratios than men, whereas men exhibit greater Type II/I fiber ratios than women.

Similar to conclusions reached in narrative reviews that considered fewer studies (Hunter, 2016a; Nuzzo, 2023), the current meta-analysis confirmed that Type II muscle fibers comprise a larger area of male than female skeletal muscle, whereas Type I muscle fibers comprise a larger area of female than male skeletal muscle. Consequently, Type II/I fiber area ratios are greater in men than women. However, unlike previous reviews that concluded there may be no difference in fiber distribution percentages between men and women (Hunter, 2016a, Nuzzo, 2023), the current meta-analysis revealed that distribution percentages of Type I muscle fibers are greater in women than men, whereas distribution percentages of Type II muscle fibers tend to be greater in men than women, though the effects are small-to-moderate in size.

TABLE 2 Sex differences in muscle fiber type characteristics as determined by random-effects model meta-analyses.

Fiber type characteristic	No. effects	Effect size			Weighted means					
		Hedges <i>g</i>	95% CI	95% PI	Men			Women		
			LB, UB	LB, UB	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD
CSA (μm ²)										
Type I	63	0.75*	0.58, 0.92	−0.26, 1.76	1403	4440.0	951.5	1279	3775.2	883.3
Type II	23	1.68*	1.26, 2.09	−0.12, 3.47	396	4671.4	2067.4	377	3127.3	1447.2
Type IIA	48	1.42*	1.18, 1.66	0.00, 2.84	1117	4923.0	1166.8	1025	3507.4	793.9
Type IIX	38	1.40*	1.05, 1.76	−0.62, 3.43	996	4310.1	1024.9	905	2844.1	739.0
MHC I	21	0.40*	0.21, 0.59	−0.07, 0.87	305	4862.6	841.1	280	4259.5	793.4
MHC II	11	1.08*	0.73, 1.44	0.02, 2.15	178	5234.4	1387.9	149	3669.2	1177.4
MHC IIA	12	0.81*	[0.45, 1.18	−0.31, 1.94	153	4643.0	700.5	152	3769.2	937.9
MHC IIX	10	1.37*	1.11, 1.64	1.06, 1.67	141	3730.6	492.3	138	2191.4	572.6
Distribution %										
Type I	97	−0.13*	−0.24, −0.03	−0.79, 0.53	2149	50.6	9.3	1774	53.2	9.4
Type II	21	0.34*	0.03, 0.64	−0.78, 1.46	520	47.5	13.2	419	45.5	13.4
Type IIA	76	0.10	0.00, 0.19	−0.25, 0.44	1732	33.6	7.1	1420	32.4	6.4
Type IIX	71	0.07	−0.05, 0.18	−0.54, 0.67	1718	15.2	5.1	1404	14.3	6.0
MHC I	51	−0.44*	−0.59, −0.30	−1.09, 0.20	905	41.0	8.9	872	46.8	8.6
MHC II	10	0.26*	0.05, 0.47	0.01, 0.51	230	53.2	9.2	150	50.9	9.6
MHC IIA	44	0.33*	0.11, 0.55	−0.84, 1.49	777	43.5	8.3	764	39.5	9.5
MHC IIX	32	0.33*	0.16, 0.49	−0.21, 0.86	655	18.4	7.3	651	14.8	7.4
Area %										
Type I	31	−0.53*	−0.76, −0.30	−1.55, 0.49	597	49.3	12.4	514	54.8	14.2
Type II	8	0.93*	0.60, 1.26	0.52, 1.34	129	59.2	13.0	132	52.9	12.3
Type IIA	25	0.40*	0.20, 0.59	−0.21, 1.01	450	33.4	8.5	368	29.6	9.3
Type IIX	25	0.19	−0.04, 0.41	−0.63, 1.00	450	15.0	5.2	368	13.4	6.2
MHC I	13	−0.69*	−0.92, −0.47	−0.95, −0.44	202	37.5	4.9	157	47.2	7.4
MHC IIA	12	0.39*	0.10, 0.68	−0.32, 1.10	186	50.3	4.9	142	45.2	6.7
MHC IIX	10	0.50*	0.26, 0.75	0.21, 0.78	150	14.9	4.0	131	9.8	4.2
Area ratio										
Type I/II	7	−1.24*	−1.85, −0.62	−3.16, 0.69	134	1.05	0.12	117	1.36	0.28
Type II/I	9	0.94*	0.53, 1.35	0.00, 1.88	88	1.27	0.33	79	1.00	0.30
Type IIA/I	10	0.63*	0.29, 0.96	−0.04, 1.29	110	1.17	0.21	89	1.02	0.14

Abbreviations: CI, confidence interval; CSA, cross-sectional area; LB, lower bound; M, men; MHC, myosin heavy chain; PI, prediction interval; UB, upper bound; W, women.

*Statistically significant effects ($p \leq 0.05$).

The cause of these sex differences in muscle fiber types is unknown. Esbjörnsson et al. (2021) have suggested that sex differences in physical activity patterns, interacting with sex differences in hormones (Haizlip et al., 2015), could contribute to sex differences in muscle fiber type. For example, men participate in physical activity and sport at higher rates than women do (Deaner et al., 2012; Gerovasili et al., 2015; Nuzzo, 2023; Van Tuyckom et al., 2010). Men are also more likely than women to prefer exercise at high intensities (Nuzzo, 2023). Thus, sex differences in physical activity preferences and participation might cause men to more regularly use fast-twitch muscle fibers in their daily lives.

The results might help to explain sex differences in exercise and sports performance. For example, as muscle *fiber* cross-sectional areas are greater in men than women, this helps to explain how cross-sectional area of *whole muscle* in women is 60%–70% of that of men (Nuzzo, 2023). Moreover, the percentage of whole muscle area occupied by Type II muscle fibers is greater in men than women. Thus, as Type II fibers produce more force and power than Type I fibers (Bottinelli et al., 1996; Bottinelli et al., 1999; Frontera et al., 2000; Krivickas et al., 2001; Krivickas et al., 2006; Larsson et al., 1997; Stienen et al., 1996; Trappe et al., 2003; Widrick et al., 2002), larger muscle sizes and greater Type II fiber distribution and area

TABLE 3 Weighted means of type I muscle fiber distribution percentages in various muscles, with weighted means for other muscle fiber types provided for vastus lateralis.

Muscle	Fiber type	Weighted means					
		Men			Women		
		n	Mean	SD	n	Mean	SD
Bicep brachii	Type I	57	47.7	8.3	45	58.1	18.5
Gastrocnemius lateralis	Type I	77	54.8	8.8	108	54.8	5.9
Gastrocnemius medialis	Type I	22	61.8	7.8	28	47.8	14.1
Longissimus/ES	Type I	150	62.7	6.2	129	67.6	7.1
Multifidus	Type I	123	59.5	8.5	97	58.6	7.8
Tibialis anterior	Type I	30	75.1	2.7	30	75.8	1.1
Triceps brachii	Type I	17	47.2	3.9	13	44.1	7.8
Vastus lateralis	Type I	1654	48.0	7.3	1303	50.6	6.9
Vastus lateralis	Type IIA	1732	33.6	7.1	1420	32.4	6.4
Vastus lateralis	Type IIX	1718	15.2	5.2	1404	14.3	6.0
Vastus lateralis	MHC I	905	41.0	8.9	872	46.8	8.6
Vastus lateralis	MHC IIA	777	43.5	8.3	764	39.5	9.5
Vastus lateralis	MHC IIX	655	18.4	7.3	651	14.8	7.4

Abbreviations: ES, erector spinae; MHC, myosin heavy chain.

percentages among men than women help to explain how muscle strength in women is ~60% of muscle strength of men (Nuzzo, 2023). Conversely, as Type I fibers possess a greater capacity for oxidative energy production than Type II fibers (Essén et al., 1975; Garnett et al., 1979), and women exhibit greater distribution and area percentages of Type I fibers than men, this might help to explain how women are less fatigable than men during some resistance exercise tasks (Hunter, 2014; 2016a; Nuzzo, 2023).

The extent to which sex differences in muscle fibers type might differently impact responses to resistance exercise between men and women also warrants consideration, though evidence of a differential impact is generally lacking. For example, a bout of eccentric resistance exercise damages Type II muscles fibers more than Type I fibers (Fridén et al., 1983; Jones et al., 1986; Lieber et al., 1991; Macaluso et al., 2012; Magal et al., 2010). Thus, as men have greater distribution and area percentages of Type II fibers, one might expect men to experience more muscle damage from eccentric resistance exercise than women. This, however, does not appear to be the case. Two reviews on the topic have concluded that men and women experience similar magnitudes of muscle soreness and strength loss after a bout of eccentric resistance exercise (Morawetz et al., 2020; Nuzzo, 2023). Whether sex differences in muscle fiber type might impact muscle fiber adaptations to weeks of resistance exercise also warrants consideration. Regarding hypertrophy of specific muscle fibers, Straight et al. (2020) reviewed results from 35 studies and found that men and women over the age of 55 experience similar increases in sizes of MHC I and MHC II muscle fibers after weeks of resistance exercise. Regarding muscle fiber type “transitions,” the most consistent finding is that Type IIX and hybrid muscle fibers transition to Type IIA fibers after weeks of resistance exercise, but evidence of a sex difference in such transitions is lacking (Bamman et al., 2003; Fry et al., 1994; Guadalupe-Grau et al., 2009; Häkkinen et al., 2002; Kosek

et al., 2006; McGuigan et al., 2001; Sharman et al., 2001; Staron et al., 1994; Williamson et al., 2001). Thus, how differences in muscle fiber types between men and women might differently impact adaptations to a given exercise intervention remains to be understood.

Sex differences in muscle fiber types can also be considered in relation to aging. Aging atrophies Type II muscle fibers more than Type I fibers, causing fast-to-slow twitch muscle fiber transition in older adults (Doherty, 2001; Lexell, 1995). Men exhibit greater proportions of Type II fibers than women, thus one might expect men would experience a greater proportional decrease in muscle strength with aging. However, mixed results have been reported regarding whether there is a sex difference in relative strength loss with aging (Charlier et al., 2015; Hughes et al., 2001; Oksuzyan et al., 2010). Preliminary analysis of an interaction between sex and age on muscle fiber type in the current research appears to show that the magnitude of the sex difference in Type I distribution percentage is impacted by age. Type I distribution percentage appears to be less in girls than in boys, then becomes greater in women than men in middle age, and then no sex difference exists between women and men older than 59 years of age. Distribution percentages for Type IIA and Type IIX demonstrate the opposite pattern. Type IIA and Type IIX distribution percentages appear to be greater in girls than in boys, then become greater in men than in women in middle age, and then no sex difference exists between women and men older than 59 years of age.

The extents to which sex differences in muscle fiber types might also associate with disease prevalence and severity can also be considered. Men have greater distribution and area percentages of Type II skeletal muscle fibers, and conditions such as Duchenne muscular dystrophy, which is more prevalent among males than females (Centers for Disease Control and Prevention, 2009; Romitti et al., 2015), impact Type II muscle fibers more than Type I fibers (Ciciliot et al., 2013; Talbot & Maves, 2016). Conversely, other

conditions impact Type I skeletal muscle fiber more than Type II fibers, causing slow-to-fast muscle fiber transitions. Examples include spinal cord injury, diabetes, heart disease, and chronic obstructive pulmonary disease (Ausín et al., 2017; Ciciliot et al., 2013; Mador & Bozkanat, 2001; Mårin et al., 1994; Talbot & Maves, 2016; Torres et al., 2011). In the current research, too little data were available from patient groups to explore potential interactive effects of disease type with sex on muscle fiber types. Nevertheless, the current research illustrates that in future studies on the impact of disease on muscle fiber types participant sex should be controlled for and/or male and female data should be presented and analyzed separately (Schilatý et al., 2018; Smith, McKay, Ackerman, et al., 2022; Smith, McKay, Kuikman, et al., 2022).

Results from the current study confirm differences in fiber type distribution percentages across muscle groups, though this was not the primary aim of this research. The muscle-specific weighted means in Table 3 illustrate that Type I distribution percentages are higher in tibialis anterior (~75%), longissimus (~65), and multifidus (~59%) and lower in vastus lateralis and biceps and triceps brachii (~45–50%). Nevertheless, differences in Type I distribution percentages across muscle groups appear to be generally similar between men and women. As the majority of muscle biopsies in the current analysis were of vastus lateralis, a potential aim for future research could be to generate more precise sex-specific estimates of fiber type characteristics of skeletal muscles other than vastus lateralis during development, aging, and disease progression. However, practical issues of acquiring muscle biopsies from willing participants are acknowledged.

Finally, the current meta-analysis is not without limitations. First, the current research did not abide by procedures for literature searches that are often used in systematic reviews nor did the current research include assessments of study quality. Consequently, results from the analysis should be considered with some caution. Nevertheless, the search was thorough. It led to the discovery of 110 studies with sex-segregated muscle fiber type data. This is significantly more than the number of studies summarized in previous narrative reviews (Hunter, 2016a; Nuzzo, 2023), and narrative reviews would be more prone to investigator bias than the current meta-analysis. Second, in the current research, data from participants of all ages, health statuses, and exercise backgrounds were combined into one large analysis. If sex interacts with age, health status, and exercise background to impact muscle fiber types then the current approach might not be ideal. The preliminary analysis of the data partitioned by age suggests that inclusion of muscle fiber data from youth (<18 years) and older adults (>59 years) decreases or underestimates the size of sex differences for some outcomes (e.g., Type I distribution percent, see Figure S9). Thus, in studying sex differences in muscle fiber types, combining data across age groups is a conservative approach that favors the null hypothesis (i.e., no sex difference in muscle fibers in middle-aged adults). These preliminary results can be tested more formally in future meta-analyses after more muscle fiber type data are presented in youth and older adult samples.

5 | CONCLUSION

Results from the current meta-analysis reveal that sex differences in muscle fiber types exist. Men exhibit greater cross-sectional areas for all muscle fiber types, greater distribution percentages for Type II, MHC II, MHC IIA, and MHC IIX fibers, and greater area percentages for Type II, Type IIA, MHC II, MHC IIA, and MHC IIX fibers. Women exhibit greater distribution and area percentages for Type I and MHC I fiber types. Sex differences in muscle fiber types might help to explain greater muscle strength and power among men than women and greater muscle endurance among women than men for some resistance exercise tasks. These data, which are believed to be the largest repository of comparative muscle fiber type data in living men and women, can inform ongoing discussions about biological sex and its impact on neuromuscular physiology, disease pathology, and sports performance.

ACKNOWLEDGMENT

Open access publishing facilitated by Edith Cowan University, as part of the Wiley - Edith Cowan University agreement via the Council of Australian University Librarians.

ORCID

James L. Nuzzo  <https://orcid.org/0000-0001-9081-0522>

REFERENCES

- Ausín, P., Martínez-Llorens, J., Sabaté-Bresco, M., Casadevall, C., Barreiro, E., & Gea, J. (2017). Sex differences in function and structure of the quadriceps muscle in chronic obstructive pulmonary disease patients. *Chronic Respiratory Disease*, 14, 127–139.
- Bamman, M. M., Hill, V. J., Adams, G. R., Haddad, F., Wetzstein, C. J., Gower, B. A., Ahmed, A., & Hunter, G. R. (2003). Gender differences in resistance-training-induced myofiber hypertrophy among older adults. *The Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*, 58, 108–116.
- Bell, D. G., & Jacobs, I. (1989). Muscle fiber-specific glycogen utilization in strength-trained males and females. *Medicine and Science in Sports and Exercise*, 21, 649–654.
- Bell, D. G., & Jacobs, I. (1990). Muscle fibre area, fibre type, and capillarization in male and female body builders. *Canadian Journal of Sport Science*, 15, 115–119.
- Bottinelli, R., Canepari, M., Pellegrino, M. A., & Reggiani, C. (1996). Force-velocity properties of human skeletal muscle fibres: Myosin heavy chain isoform and temperature dependence. *Journal of Physiology*, 495(Pt 2), 573–586.
- Bottinelli, R., Pellegrino, M. A., Canepari, M., Rossi, R., & Reggiani, C. (1999). Specific contributions of various muscle fibre types to human muscle performance: An in vitro study. *Journal of Electromyography and Kinesiology*, 9, 87–95.
- Cagnie, B., Dhooze, F., Schumacher, C., De Meulemeester, K., Petrovic, M., VAN Oosterwijck, J., & Danneels, L. (2015). Fiber typing of the erector spinae and multifidus muscles in healthy controls and back pain patients: A systematic literature review. *Journal of Manipulative and Physiological Therapeutics*, 38, 653–663.
- Centers for Disease Control and Prevention. (2009). Prevalence of Duchenne/Becker muscular dystrophy among males aged 5–24 years—four states, 2007. *Morbidity and Mortality Weekly Report*, 58, 1119–1122.

- Chalmers, G. R., & Row, B. S. (2011). Common errors in textbook descriptions of muscle fiber size in nontrained humans. *Sports Biomechanics*, 10, 254–268.
- Charlier, R., Mertens, E., Lefevre, J., & Thomis, M. (2015). Muscle mass and muscle function over the adult life span: A cross-sectional study in Flemish adults. *Archives of Gerontology and Geriatrics*, 61, 161–167.
- Ciciliot, S., Rossi, A. C., Dyar, K. A., Blaauw, B., & Schiaffino, S. (2013). Muscle type and fiber type specificity in muscle wasting. *International Journal of Biochemistry and Cell Biology*, 45, 2191–2199.
- Costill, D. L., Daniels, J., Evans, W., Fink, W., Krahenbuhl, G., & Saltin, B. (1976). Skeletal muscle enzymes and fiber composition in male and female track athletes. *Journal of Applied Physiology*, 40, 149–154.
- Cumming, G. (2009). Inference by eye: Reading the overlap of independent confidence intervals. *Statistics in Medicine*, 28, 205–220.
- Damer, A., El Meniawy, S., Mcpherson, R., Wells, G., Harper, M. E., & Dent, R. (2022). Association of muscle fiber type with measures of obesity: A systematic review. *Obesity Reviews*, 23, e13444.
- Deaner, R. O., Geary, D. C., Puts, D. A., Ham, S. A., Kruger, J., Fles, E., Winegard, B., & Grandis, T. (2012). A sex difference in the predisposition for physical competition: Males play sports much more than females even in the contemporary U.S. *PLoS One*, 7, e49168.
- Doherty, T. J. (2001). The influence of aging and sex on skeletal muscle mass and strength. *Current Opinion in Clinical Nutrition and Metabolic Care*, 4, 503–508.
- Esbjörnsson, M. E., Dahlström, M. S., Gierup, J. W., & Jansson, E. C. (2021). Muscle fiber size in healthy children and adults in relation to sex and fiber types. *Muscle & Nerve*, 63, 586–592.
- Essén, B., Jansson, E., Henriksson, J., Taylor, A. W., & Saltin, B. (1975). Metabolic characteristics of fibre types in human skeletal muscle. *Acta Physiologica Scandinavica*, 95, 153–165.
- Evertsen, F., Medbo, J. I., Jebens, E., & Gjøvaag, T. F. (1999). Effect of training on the activity of five muscle enzymes studied on elite cross-country skiers. *Acta Physiologica Scandinavica*, 167, 247–257.
- Fridén, J., Sjöström, M., & Ekblom, B. (1983). Myofibrillar damage following intense eccentric exercise in man. *International Journal of Sports Medicine*, 4, 170–176.
- Froese, E. A., & Houston, M. E. (1985). Torque-velocity characteristics and muscle fiber type in human vastus lateralis. *Journal of Applied Physiology*, 59, 309–314.
- Froese, E. A., & Houston, M. E. (1987). Performance during the Wingate anaerobic test and muscle morphology in males and females. *International Journal of Sports Medicine*, 8, 35–39.
- Frontera, W. R., Suh, D., Krivickas, L. S., Hughes, V. A., Goldstein, R., & Roubenoff, R. (2000). Skeletal muscle fiber quality in older men and women. *American Journal of Physiology. Cell Physiology*, 279, C611–C618.
- Fry, A. C., Allemeier, C. A., & Staron, R. S. (1994). Correlation between percentage fiber type area and myosin heavy chain content in human skeletal muscle. *European Journal of Applied Physiology and Occupational Physiology*, 68, 246–251.
- Garnett, R. A., O'donovan, M. J., Stephens, J. A., & Taylor, A. (1979). Motor unit organization of human medial gastrocnemius. *Journal of Physiology*, 287, 33–43.
- Gerovasili, V., Agaku, I. T., Vardavas, C. I., & Filippidis, F. T. (2015). Levels of physical activity among adults 18–64 years old in 28 European countries. *Preventive Medicine*, 81, 87–91.
- Gouzi, F., Maury, J., Molinari, N., Pomiès, P., Mercier, J., Préfaut, C., & Hayot, M. (2013). Reference values for vastus lateralis fiber size and type in healthy subjects over 40 years old: A systematic review and metaanalysis. *Journal of Applied Physiology*, 115, 346–354.
- Greenhalgh, T., & Peacock, R. (2005). Effectiveness and efficiency of search methods in systematic reviews of complex evidence: Audit of primary sources. *BMJ*, 331, 1064–1065.
- Guadalupe-Grau, A., Perez-Gomez, J., Olmedillas, H., Chavarren, J., Dorado, C., Santana, A., Serrano-Sanchez, J. A., & Calbet, J. A. (2009). Strength training combined with plyometric jumps in adults: Sex differences in fat-bone axis adaptations. *Journal of Applied Physiology*, 106, 1100–1111.
- Haizlip, K. M., Harrison, B. C., & Leinwand, L. A. (2015). Sex-based differences in skeletal muscle kinetics and fiber-type composition. *Physiology*, 30, 30–39.
- Häkkinen, K., Kraemer, W. J., Pakarinen, A., Triplett-Mcbride, T., McBride, J. M., Häkkinen, A., Alen, M., McGuigan, M. R., Bronks, R., & Newton, R. U. (2002). Effects of heavy resistance/power training on maximal strength, muscle morphology, and hormonal response patterns in 60–75-year-old men and women. *Canadian Journal of Applied Physiology*, 27, 213–231.
- Hilton, E. N., & Lundberg, T. R. (2021). Transgender women in the female category of sport: Perspectives on testosterone suppression and performance advantage. *Sports Medicine*, 51, 199–214.
- Holmbäck, A. M., Porter, M. M., Downham, D., Andersen, J. L., & Lexell, J. (2003). Structure and function of the ankle dorsiflexor muscles in young and moderately active men and women. *Journal of Applied Physiology*, 95, 2416–2424.
- Hughes, V. A., Frontera, W. R., Wood, M., Evans, W. J., Dallal, G. E., Roubenoff, R., & Fiatarone Singh, M. A. (2001). Longitudinal muscle strength changes in older adults: Influence of muscle mass, physical activity, and health. *The Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*, 56, B209–B217.
- Hunter, S. K. (2014). Sex differences in human fatigability: Mechanisms and insight to physiological responses. *Acta Physiologica*, 210, 768–789.
- Hunter, S. K. (2016a). The relevance of sex differences in performance fatigability. *Medicine and Science in Sports and Exercise*, 48, 2247–2256.
- Hunter, S. K. (2016b). Sex differences in fatigability of dynamic contractions. *Experimental Physiology*, 101, 250–255.
- Inthout, J., Ioannidis, J. P., Rovers, M. M., & Goeman, J. J. (2016). Plea for routinely presenting prediction intervals in meta-analysis. *BMJ Open*, 6, e010247.
- Jaworowski, A., Porter, M. M., Holmbäck, A. M., Downham, D., & Lexell, J. (2002). Enzyme activities in the tibialis anterior muscle of young moderately active men and women: Relationship with body composition, muscle cross-sectional area and fibre type composition. *Acta Physiologica Scandinavica*, 176, 215–225.
- Jones, D. A., Newham, D. J., Round, J. M., & Tolfree, S. E. (1986). Experimental human muscle damage: Morphological changes in relation to other indices of damage. *Journal of Physiology*, 375, 435–448.
- Jones, M. D., Wewege, M. A., Hackett, D. A., Keogh, J. W. L., & Hagstrom, A. D. (2021). Sex differences in adaptations in muscle strength and size following resistance training in older adults: A systematic review and meta-analysis. *Sports Medicine*, 51, 503–517.
- Kosek, D. J., Kim, J. S., Petrella, J. K., Cross, J. M., & Bamman, M. M. (2006). Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *Journal of Applied Physiology*, 101, 531–544.
- Krivickas, L. S., Fielding, R. A., Murray, A., Callahan, D., Johansson, A., Dor, D. J., & Frontera, W. R. (2006). Sex differences in single muscle fiber power in older adults. *Medicine and Science in Sports and Exercise*, 38, 57–63.
- Krivickas, L. S., Suh, D., Wilkins, J., Hughes, V. A., Roubenoff, R., & Frontera, W. R. (2001). Age- and gender-related differences in maximum shortening velocity of skeletal muscle fibers. *American Journal of Physical Medicine and Rehabilitation*, 80, 447–455.
- Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: A practical primer for t-tests and ANOVAs. *Frontiers in Psychology*, 4, 863.
- Larsson, L., Li, X., & Frontera, W. R. (1997). Effects of aging on shortening velocity and myosin isoform composition in single human skeletal muscle cells. *American Journal of Physiology*, 272, C638–C649.

- Lexell, J. (1995). Human aging, muscle mass, and fiber type composition. *The Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*, 50, 11–16.
- Lieber, R. L., Woodburn, T. M., & Fridén, J. (1991). Muscle damage induced by eccentric contractions of 25% strain. *Journal of Applied Physiology*, 70, 2498–2507.
- Macaluso, F., Isaacs, A. W., & Myburgh, K. H. (2012). Preferential type II muscle fiber damage from plyometric exercise. *Journal of Athletic Training*, 47, 414–420.
- Mador, M. J., & Bozkanat, E. (2001). Skeletal muscle dysfunction in chronic obstructive pulmonary disease. *Respiratory Research*, 2, 216–224.
- Magal, M., Dumke, C. L., Urbiztondo, Z. G., Cavill, M. J., Triplett, N. T., Quindry, J. C., McBride, J. M., & Epstein, Y. (2010). Relationship between serum creatine kinase activity following exercise-induced muscle damage and muscle fibre composition. *Journal of Sports Sciences*, 28, 257–266.
- Mårin, P., Andersson, B., Krotkiewski, M., & Björntorp, P. (1994). Muscle fiber composition and capillary density in women and men with NIDDM. *Diabetes Care*, 17, 382–386.
- Mcguigan, M. R., Bronks, R., Newton, R. U., Sharman, M. J., Graham, J. C., Cody, D. V., & Kraemer, W. J. (2001). Resistance training in patients with peripheral arterial disease: Effects on myosin isoforms, fiber type distribution, and capillary supply to skeletal muscle. *The Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*, 56, B302–B310.
- Morawetz, D., Blank, C., Koller, A., Arvandi, M., Siebert, U., & Schoberberger, W. (2020). Sex-related differences after a single bout of maximal eccentric exercise in response to acute effects: A systematic review and meta-analysis. *Journal of Strength and Conditioning Research*, 34, 2697–2707.
- Naruse, M., Trappe, S., & Trappe, T. A. (2023). Human skeletal muscle-specific atrophy with aging: A comprehensive review. *Journal of Applied Physiology*, 134, 900–914.
- Ng, J. K., Richardson, C. A., Kippers, V., & Parnianpour, M. (1998). Relationship between muscle fiber composition and functional capacity of back muscles in healthy subjects and patients with back pain. *Journal of Orthopaedic and Sports Physical Therapy*, 27, 389–402.
- Nokoff, N. J., Senefeld, J., Krausz, C., Hunter, S., & Joyner, M. (2023). Sex differences in athletic performance: Perspectives on transgender athletes. *Exercise and Sport Sciences Reviews*, 51, 85–95.
- Nuzzo, J. L. (2023). Narrative review of sex differences in muscle strength, endurance, activation, size, fiber type, and strength training participation rates, preferences, motivations, injuries, and neuromuscular adaptations. *Journal of Strength and Conditioning Research*, 37, 494–536.
- Nygaard, E., Houston, M., Suzuki, Y., Jorgensen, K., & Saltin, B. (1983). Morphology of the brachial biceps muscle and elbow flexion in man. *Acta Physiologica Scandinavica*, 117, 287–292.
- Oksuzyan, A., Crimmins, E., Saito, Y., O'rand, A., Vaupel, J. W., & Christensen, K. (2010). Cross-national comparison of sex differences in health and mortality in Denmark, Japan and the US. *European Journal of Epidemiology*, 25, 471–480.
- Roberts, B. M., Nuckols, G., & Krieger, J. W. (2020). Sex differences in resistance training: A systematic review and meta-analysis. *Journal of Strength and Conditioning Research*, 34, 1448–1460.
- Rolf, C., Andersson, G., Westblad, P., & Saltin, B. (1997). Aerobic and anaerobic work capacities and leg muscle characteristics in elite orienteers. *Scandinavian Journal of Medicine and Science in Sports*, 7, 20–24.
- Romitti, P. A., Zhu, Y., Puzhankara, S., James, K. A., Nabukera, S. K., Zamba, G. K., Cifaloni, E., Cuniff, C., Druschel, C. M., Mathews, K. D., Matthews, D. J., Meaney, F. J., Andrews, J. G., Conway, K. M., Fox, D. J., Street, N., Adams, M. M., & Bolen, J. (2015). Prevalence of Duchenne and Becker muscular dystrophies in the United States. *Pediatrics*, 135, 513–521.
- Schilaty, N. D., Bates, N. A., & Hewett, T. E. (2018). Relative dearth of 'sex differences' research in sports medicine. *Journal of Science and Medicine in Sport*, 21, 440–441.
- Scott, W., Stevens, J., & Binder-Macleod, S. A. (2001). Human skeletal muscle fiber type classifications. *Physical Therapy*, 81, 1810–1816.
- Sharman, M. J., Newton, R. U., Triplett-Mcbride, T., Mcguigan, M. R., McBride, J. M., Häkkinen, A., Häkkinen, K., & Kraemer, W. J. (2001). Changes in myosin heavy chain composition with heavy resistance training in 60- to 75-year-old men and women. *European Journal of Applied Physiology*, 84, 127–132.
- Sjøgaard, G. (1982). Capillary supply and cross-sectional area of slow and fast twitch muscle fibres in man. *Histochemistry*, 76, 547–555.
- Smith, E. S., McKay, A. K. A., Ackerman, K. E., Harris, R., Elliott-Sale, K. J., Stellingwerff, T., & Burke, L. M. (2022). Methodology review: A protocol to audit the representation of female athletes in sports science and sports medicine research. *International Journal of Sport Nutrition and Exercise Metabolism*, 32, 114–127.
- Smith, E. S., McKay, A. K. A., Kuikman, M., Ackerman, K. E., Harris, R., Elliott-Sale, K. J., Stellingwerff, T., & Burke, L. M. (2022). Auditing the representation of female versus male athletes in sports science and sports medicine research: Evidence-based performance supplements. *Nutrients*, 14, 3372.
- Staron, R. S. (1991). Correlation between myofibrillar ATPase activity and myosin heavy chain composition in single human muscle fibers. *Histochemistry*, 96, 21–24.
- Staron, R. S., Hagerman, F. C., Hikida, R. S., Murray, T. F., Hostler, D. P., Crill, M. T., Ragg, K. E., & Toma, K. (2000). Fiber type composition of the vastus lateralis muscle of young men and women. *Journal of Histochemistry and Cytochemistry*, 48, 623–629.
- Staron, R. S., Karapondo, D. L., Kraemer, W. J., Fry, A. C., Gordon, S. E., Falkel, J. E., Hagerman, F. C., & Hikida, R. S. (1994). Skeletal muscle adaptations during early phase of heavy-resistance training in men and women. *Journal of Applied Physiology*, 76, 1247–1255.
- Stienen, G. J., Kiers, J. L., Bottinelli, R., & Reggiani, C. (1996). Myofibrillar ATPase activity in skinned human skeletal muscle fibres: Fibre type and temperature dependence. *Journal of Physiology*, 493(Pt 2), 299–307.
- Straight, C. R., Fedewa, M. V., Toth, M. J., & Miller, M. S. (2020). Improvements in skeletal muscle fiber size with resistance training are age-dependent in older adults: A systematic review and meta-analysis. *Journal of Applied Physiology*, 129, 392–403.
- Talbot, J., & Maves, L. (2016). Skeletal muscle fiber type: Using insights from muscle developmental biology to dissect targets for susceptibility and resistance to muscle disease. *Wiley Interdisciplinary Reviews: Developmental Biology*, 5, 518–534.
- Torres, S. H., Montes DE Oca, M., Loeb, E., Mata, A., & Hernández, N. (2011). Gender and skeletal muscle characteristics in subjects with chronic obstructive pulmonary disease. *Respiratory Medicine*, 105, 88–94.
- Trappe, S., Gallagher, P., Harber, M., Carrithers, J., Fluckey, J., & Trappe, T. (2003). Single muscle fibre contractile properties in young and old men and women. *Journal of Physiology*, 552, 47–58.
- Van Tuyckom, C., Scheerder, J., & Bracke, P. (2010). Gender and age inequalities in regular sports participation: A cross-national study of 25 European countries. *Journal of Sports Sciences*, 28, 1077–1084.
- Vinciguerra, C., Iacono, S., Bevilacqua, L., Landolfi, A., Piscosquito, G., Ginanneschi, F., Schirò, G., Di Stefano, V., Brighina, F., Barone, P., & Balistreri, C. R. (2023). Sex differences in neuromuscular disorders. *Mechanisms of Ageing and Development*, 211, 111793.
- Widrick, J. J., Stelzer, J. E., Shoepe, T. C., & Garner, D. P. (2002). Functional properties of human muscle fibers after short-term resistance exercise training. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 283, R408–R416.

Williamson, D. L., Gallagher, P. M., Carroll, C. C., Raue, U., & Trappe, S. W. (2001). Reduction in hybrid single muscle fiber proportions with resistance training in humans. *Journal of Applied Physiology*, 91, 1955–1961.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Nuzzo, J. L. (2023). Sex differences in skeletal muscle fiber types: A meta-analysis. *Clinical Anatomy*, 1–11. <https://doi.org/10.1002/ca.24091>