

2023

Comparing clinico-demographics and neuropsychiatric symptoms for immigrant and non-immigrant aged care residents living with dementia: A retrospective cross-sectional study from an Australian dementia-specific support service

Pelden Chejor
Edith Cowan University

Mustafa Atee
Edith Cowan University

Patricia Cain
Edith Cowan University

Daniel Whiting

Thomas Morris

See next page for additional authors

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworks2022-2026>



Part of the [Geriatric Nursing Commons](#)

[10.1186/s12877-023-04447-3](https://doi.org/10.1186/s12877-023-04447-3)

Chejor, P., Atee, M., Cain, P., Whiting, D., Morris, T., & Porock, D. (2023). Comparing clinico-demographics and neuropsychiatric symptoms for immigrant and non-immigrant aged care residents living with dementia: A retrospective cross-sectional study from an Australian dementia-specific support service. *BMC Geriatrics*, 23, article 729. <https://doi.org/10.1186/s12877-023-04447-3>

This Journal Article is posted at Research Online.
<https://ro.ecu.edu.au/ecuworks2022-2026/3182>

Authors

Pelden Chejor, Mustafa Atee, Patricia Cain, Daniel Whiting, Thomas Morris, and Davina Porock

RESEARCH

Open Access



Comparing clinico-demographics and neuropsychiatric symptoms for immigrant and non-immigrant aged care residents living with dementia: a retrospective cross-sectional study from an Australian dementia-specific support service

Pelden Chejor^{1*}, Mustafa Atee^{1,2,3,4}, Patricia Cain¹, Daniel Whiting⁵, Thomas Morris⁵ and Davina Porock¹

Abstract

Background Neuropsychiatric symptoms of dementia such as agitation and aggression are common in people living with dementia. The presentation of neuropsychiatric symptoms is influenced by the cultural background of people living with dementia. Further, identifying factors contributing to neuropsychiatric symptoms may be complicated if people living with dementia are immigrants or from non-English-speaking backgrounds. Most of what is known about differences in neuropsychiatric symptoms between racial and ethnic groups living with dementia come from community-based samples. This study investigated differences in clinico-demographics and neuropsychiatric symptoms between immigrants and non-immigrants living with dementia in residential aged care homes who were referred to two Dementia Support Australia programs.

Methods This was a retrospective observational cross-sectional study from 2018 to 2022 using data extracted from the Dementia Support Australia database. Immigrant status was identified by documented country of birth. We conducted exploratory subgroup analyses for English-speaking or non-English-speaking immigrants in comparison to non-immigrants. Neuropsychiatric Inventory and PainChek[®] were used to assess neuropsychiatric symptoms of dementia and pain, respectively.

Results Of the 23,889 referrals, 36% were immigrants living with dementia. Immigrants were 0.8 years older than non-immigrants on average. Immigrants had a slightly higher prevalence of mixed dementia (9.5%) than non-immigrants (8.2%). Overall, the groups had no difference in the severity of neuropsychiatric symptoms and associated caregiver distress. However, there was a significant difference in the total number of neuropsychiatric inventory domains (Cohen's $d = -0.06$ [-0.09, -0.02], $p < .001$) between non-English-speaking immigrants and non-immigrants. Immigrants were more likely to present with agitation/aggression, while non-immigrants were more likely to present

*Correspondence:

Pelden Chejor

p.chejor@ecu.edu.au

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

with hallucinations. Factors contributing to neuropsychiatric symptoms were common between the groups, with language barriers and cultural considerations frequently endorsed for immigrants.

Conclusion This study reveals a mixed picture of neuropsychiatric symptoms between immigrants and non-immigrants. However, due to the exploratory nature of the hypotheses, our findings need to be replicated in future studies to confirm any conclusions. There is a need for increased awareness on the impact of culture and language on neuropsychiatric symptoms for people receiving residential care. Future studies investigating neuropsychiatric symptoms in different immigrant groups will help increase our understanding of neuropsychiatric symptoms for all people.

Keywords Immigrants, Neuropsychiatric symptoms, Dementia, NPS, Non-English-speaking, Caregiver distress, Pain

Introduction

In Australia, over 400,000 people are living with dementia, a number projected to double by 2058 [1]. At least 54% (132,000) of people living in residential aged care homes (RACHs) in 2019 – 2020 had a dementia diagnosis [1]. Over 31% of aged care residents were also born overseas (19.6% from non-English-speaking countries and 12.2% from other English-speaking countries), and 9.2% of people using aged care preferred a language other than English [2]. In 2019 – 2020, 21% of people living with dementia in RACHs were immigrants from non-English speaking countries [1]. International studies have reported that immigrants experience a higher prevalence of dementia due to differing life experiences including those related to trauma, low literacy, and socioeconomic status [3–5].

Caring for people living with dementia presents specific challenges, such as communication due to the loss of cognitive function, loss of language, neuropsychiatric symptoms (NPS), and the complex progressive nature of the condition [6]. Care complexity can increase when the recipient is an immigrant. As dementia progresses, it is common for people from non-English-speaking backgrounds to lose their ability to communicate in English and revert to using their first language to communicate [7, 8]. Resultant communication barriers where formal care providers do not share the same language may complicate the ability of care providers to understand and meet the needs of people living with dementia [9], particularly around expression of pain and NPS.

More than 95% of people living with dementia in RACHs experience at least one NPS such as agitation or aggression [10]. However, most of what is known about differences in NPS based on racial and ethnic groups living with dementia comes from community-based samples. For people living with Alzheimer's disease in the community, it has been reported that NPS such as hallucinations, night-time behaviour, and elation were more common for people with Hispanic backgrounds compared to people from non-Hispanic backgrounds and that the severity of NPS differed between the two groups [11]. A recent study found that NPS were more likely to occur

in non-Hispanic Black *[sic]* participants with dementia than they were in non-Hispanic White *[sic]* participants with dementia [12]. Similarly, a case-control study from a Dutch urban memory clinic (case, $n = 415$, control, $n = 428$) showed that older ethnic minority patients with dementia migrating from 47 countries had significantly more depression, change in appetite, agitation/aggression, disinhibition, and aberrant motor behaviour than native Dutch people with dementia [13].

NPS can impact both people living with dementia and their caregivers. For the individual, NPS are associated with faster disease progression [14, 15] and poor quality of life [16, 17]. For caregivers, research has focused mainly on the informal caregiver burden and less on the formal caregiver perspective. NPS are related to increased informal caregiver distress [18] and burden [19, 20] as well as poor caregiver mental health [21]. Cultural diversity may also impact caregiver burden, as shown in one study, where foreign-born informal caregivers of people living with dementia reported more stress associated with NPS than native Canadian caregivers [22]. In addition, a recent systematic review found that NPS increases the burden for caregivers of older immigrants living with dementia [23]. Fauth and Gibbons [24] also noted that NPS-associated caregiver distress could be higher for family or informal caregivers than formal caregivers who are routinely dealing with NPS.

Factors contributing to the development of NPS are numerous and are often categorised as factors related to people living with dementia (e.g., neurobiologically related disease factors, medical illness, unmet needs), caregiver factors (e.g., stress, burden, communication issues), and factors related to environment or care settings (over- or under-stimulation, unsafe environment) [25]. Undiagnosed medical illnesses of people living with dementia have previously been related to NPS [26]. Pain is also strongly linked to agitation/aggression [27] as people living with dementia often express pain and discomfort by displaying aggressive behaviours [28]. Caregiver factors such as increased stress and depression were related to more NPS in care recipients [29], while changes in daily routines (e.g., bathing and dressing) were

associated with stress for people living with dementia [30]. The complexity of NPS means understanding the underlying causes of NPS is important for tailoring support for people living with dementia and their caregivers. Pain, carer approach, loneliness/boredom, mood disorders, and communication difficulties were identified as the common contributing factors to NPS for people living with dementia in a national Australian sample referred to the Dementia Support Australia (DSA) programs [31]. Other factors including sleep disorders [32], sex, and education [33] were also attributed as potentially contributing to NPS. Further factors may also contribute to NPS and identifying these may be complicated in instances where people living with dementia are immigrants or from non-English-speaking backgrounds. This topic remains largely under-researched, particularly in Australia, where the older immigrant population living with dementia is projected to increase [34].

This study aimed to compare the clinico-demographic characteristics (e.g., age, sex, dementia subtype), prevalence and severity of NPS, and associated caregiver distress among people living with dementia (immigrants and non-immigrants) in RACHs referred to DSA, a national provider of dementia behaviour support. To address the objectives of this study, we set the following hypotheses:

- (1) there will be differences in demographic characteristics between immigrants (English-speaking or not) and non-immigrants.
- (2) there will be differences in the individual and overall prevalence and severity of NPS between immigrants (English-speaking or not) and non-immigrants.
- (3) there will be differences in overall caregiver distress reported by caregivers of immigrants (English-speaking or not) and non-immigrants.
- (4) there will be differences in the prevalence of contributing factors for NPS in immigrants (English-speaking or not) and non-immigrants.

Methods

Study design and setting

This observational cross-sectional study drew on retrospective demographic and clinical data (e.g., age, dementia subtype) from the DSA database from 1 June 2018 to 30 June 2022. DSA is a dementia-specific government-funded program in Australia that provides free-of-charge national support programs in the form of individualised psychosocial or non-pharmacological interventions for people living in various care settings with varying severity of NPS-related dementia [35]. Support from DSA includes the Dementia Behaviour Management Advisory Service (DBMAS) for mild to moderate NPS and the

Severe Behaviour Response Teams (SBRT) for moderate to severe NPS. Eligibility criteria for receiving support from DSA include: (1) having a confirmed or probable diagnosis of dementia; (2) people living with dementia exhibiting NPS related to dementia; (3) people living with dementia exhibiting NPS that impacts their care and well-being, or their caregivers; and (4) receiving consent from people living with dementia or their responsible caregiver [31, 36]. This study focused on support provided by DSA in RACHs.

Data extraction

Demographic and clinical data (age, sex, dementia subtypes, country of birth, Neuropsychiatric Inventory (NPI) scores, pain scores, and contributing factors) were extracted by a data custodian using the DSA database for RACHs referrals with NPS seeking support from DBMAS and SBRT programs [35].

Study instruments

The NPI and PainChek® are clinical instruments that DSA routinely administer to assess NPS and pain, respectively. The NPI is used during each consultation and the version of NPI used is determined by the type of DSA programs delivered. The Neuropsychiatric Inventory Questionnaire (NPI-Q) is administered for DBMAS referrals while the Neuropsychiatric Inventory Nursing Home version (NPI-NH) is administered for SBRT referrals [35].

The NPI-NH [36] and NPI-Q [37] are informant-based instruments that are valid, consistent, and reliable tools for assessing 12 NPI domains -aberrant motor behaviour, agitation/aggression, anxiety, apathy/indifference, appetite and eating, delusions, depression/dysphoria, disinhibition, elation/euphoria, hallucinations, irritability/labability, and night-time behaviour [36, 37]. The NPI-Q and NPI-NH rate each NPI domain on its presence (Yes = present, No = absent) and severity ('mild' (1), 'moderate' (2), 'severe' (3)) [36, 37]. The NPI-Q and NPI-NH also assess the caregiver distress/disruptiveness associated with each of the 12 NPI domains with ratings from ('not at all' (0), 'minimally' (1), 'mildly' (2), 'moderately' (3), 'severely' (4), 'extremely' (5)) [36, 37]. The total caregiver distress/disruptiveness scores can be calculated separately by adding the distress/disruptiveness scores of all 12 NPI domains. Higher scores on distress denote more distressing behaviours, while higher scores on disruptiveness denote more disruptive behaviours. For our scoring, scores from both NPI-Q and NPI-NH are referred to as total NPI scores, and the caregiver distress/disruptiveness is referred to as caregiver distress.

Contributing factors to NPS in DSA referrals

Contributing factors to NPS are identified by DSA consultants through a comprehensive onsite assessment of referrals. This includes consultation with the referral and their formal and/or informal caregivers and a review of the referral's clinical files such as medical records and social history. DSA consultants conduct clinical observations and interviews, collect proxy assessments from families and caregivers, and use validated instruments such as PainChek[®] and the NPI [35]. During an assessment, DSA consultants use a comprehensive list of factors contributing to NPS and assign those applicable to the referral. The list of potential contributing factors covers several broad causes of NPS from the categories of biological, environmental, psychological, and social factors [31].

Pain is routinely assessed by DSA consultants using the PainChek[®] Adult, a multimodal, psychometrically sound, artificial intelligence (AI)-based pain assessment tool in the form of a point-of-care app for non-verbal adults [39]. PainChek[®] is registered as a software medical device by Australia's Therapeutic Goods Administration, Health Canada, Singapore Health Sciences Authority, and European Conformity [38, 39]. The PainChek[®] pain assessment scale is a 42-item instrument assessing six domains [Face (9 items), Voice (9 items), Movement (7 items), Behaviour (7 items), Activity (4 items), and Body (6 items)]. Each item is provided with a clear operational definition to improve interrater consistency, and items are rated on a binary level (Yes = present, No = absent) [38]. The final pain score ranges from 0 (no pain) to 42 (severe pain). Pain is present if the final score is ≥ 7 and absent if the final score is <7 [39, 40].

Similar to Loi et al [31], we investigated in this study the prevalence of common contributing factors of NPS plus two additional factors (language barriers and cultural considerations), which are directly related to our study objectives.

Eligibility criteria of participants for data analysis

Data were included in the study if the data related to someone who was: (1) referred to DSA programs during the study period; (2) residing in RACH at the time of referral; and (3) aged 65 years or older. If the referrals accessed DSA services multiple times during the study period, the earliest referral was used. Referrals not meeting any of the above criteria were excluded.

Data analysis

For analysis, data were classified into two groups according to country of birth: (1) non-immigrant group for those referrals whose country of birth was documented as Australia; (2) immigrant group for those referrals

whose country of birth documented was other than Australia. The immigrant group was further classified into subgroups as English-speaking immigrants or non-English-speaking immigrants depending on whether they were born in countries where English is the primary language or not, using the World Population Review list of English-Speaking Countries 2022 [41].

We used R version 4.2.2 [42] for the statistical analyses. We considered a p -value of $<.001$ statistically significant to reduce the risk of false positives [43] as p -values would easily reach statistical significance in large sample sizes [44]. Demographic characteristics (age, gender, dementia subtypes) and NPS characteristics (total number of domains, severity, and caregiver distress) between groups were compared. Descriptive statistics (mean, standard deviation, frequency, and percentages) were used to describe demographic variables and assessment characteristics. Pearson's chi-square test with Cramer's V effect size was used to compare categorical demographic variables (Hypothesis 1 and 2). Cramer's V effect sizes were interpreted in relation to the associated degrees of freedom of chi-square test where small effect values are in the range of 0.04 (for five degrees of freedom) to 0.10 (for 1 degree of freedom) [45].

For Hypothesis 1, 2 and 3, continuous variables were compared using Welch's t -test and Cohen's d effect sizes, with a 95% confidence interval. Cohen's d effect sizes were interpreted as small ($d = 0.2$), medium ($d = 0.5$), and large ($d = 0.8$) [46]. A logistic regression model was used to predict immigrant status from the individual NPI domain severity scores (Hypothesis 2). Two linear regression models with immigrant status, age, and sex as predictors were also created, using total NPS severity and total caregiver distress as dependent variables, to investigate the relationship between immigrant status and total NPS severity and total caregiver distress scores. For all regression analyses, the reference group was non-immigrants living with dementia. The rates of most frequently identified contributing factors (in addition to language barriers and cultural considerations) for each group, were compared using a two-sample Z -test with a 95% confidence interval (Hypothesis 4).

Results

Descriptive characteristics

A total of 23,889 referrals to DSA were eligible for the study, of which 36% were immigrants living with dementia ($n = 8607$, 55.4% female). Table 1 shows the descriptive characteristics of non-immigrants and immigrants, with further categorisation for non-English-speaking immigrants and English-speaking immigrants. There was a significant difference in age, with immigrants being 0.8 years older than non-immigrants on average (Cohen's d

Table 1 Descriptive characteristics of non-immigrants, immigrants, NES immigrants, and ES immigrants

Group	Non-immigrants	Immigrants	Effect size [95% CI]	<i>p</i>	NES immigrants	ES immigrants
Sample size, <i>n</i> (%)	15,282 (64.0%)	8,607 (36.0%)	NA	NA	5,723 (23.9%)	2,884 (12.1%)
Age, years						
Mean (SD)	83.7 (7.6)	84.5 (7.0)	0.12 [0.09, 0.14]*	< .001	85.0 (6.9)	83.7 (7.2)
Sex, <i>n</i> (%)						
Female	8,770 (57.4%)	4,766 (55.4%)	0.02 [0.00, 0.03]**	.003	3,155 (55.1%)	1,611 (55.9%)
Male	6,482 (42.4%)	3,821 (44.4%)			2,554 (44.6%)	1,267 (43.9%)
Missing	18 (0.1%)	15 (0.2%)			10 (0.2%)	5 (0.2%)
Intersex or indeterminate	12 (0.1%)	5 (0.1%)			4 (0.1%)	1 (0.0%)
Dementia subtype, <i>n</i> (%)						
Alzheimer's disease	5,747 (37.6%)	3,251 (37.8%)	0.03 [0.01, 0.05]**	< .001	2,033 (35.5%)	1,218 (42.2%)
Dementia unspecified	4,412 (28.9%)	2,503 (29.1%)			1,734 (30.3%)	769 (26.7%)
Vascular dementia	1,910 (12.5%)	1,114 (12.9%)			767 (13.4%)	347 (12.0%)
Mixed dementia	1,260 (8.2%)	817 (9.5%)			598 (10.4%)	219 (7.6%)
Other dementias	775 (5.1%)	364 (4.2%)			232 (4.1%)	132 (4.6%)
Lewy body dementia	417 (2.7%)	220 (2.6%)			130 (2.3%)	90 (3.1%)
Frontal lobe dementia	380 (2.5%)	157 (1.8%)			107 (1.9%)	50 (1.7%)
Dementia in Parkinson's disease	238 (1.6%)	104 (1.2%)			70 (1.2%)	34 (1.2%)
Missing	143 (0.9%)	77 (0.9%)			52 (0.9%)	25 (0.9%)
NPI Totals, Mean (SD)						
Severity	10.8 (5.9)	10.8 (5.8)	0.00 [-0.03, 0.02]*	.746	10.6 (5.7)	11.2 (5.8)
Caregiver distress	14.6 (8.6)	14.6 (8.6)	0.00 [-0.03, 0.03]*	.923	14.3 (8.6)	15.0 (8.6)
Number of Domains	5.2 (2.2)	5.2 (2.2)	-0.02 [-0.05, 0.01]*	.269	5.1 (2.2)	5.3 (2.2)
Pain Present, <i>n</i> (%)						
PainChek® score ≥7 (Pain present)	5,039 (68.0%)	3,016 (68.4%)	0.00 [0.00, 0.02]**	.720	2,036 (69.9%)	980 (65.3%)
PainChek® score <7 (Pain absent)	2,367 (32.0%)	1,395 (31.6%)			875 (30.1%)	520 (34.7%)

Percentages for the NES and ES immigrants are calculated using the number of all immigrants

CI, *p*-value, and effect sizes are only applicable to non-immigrants and all immigrants

* Cohen's *d* effect size; **Cramer's *V* effect size, NES Non-English-speaking, ES English-speaking, NA Not applicable, NPI Neuropsychiatric inventory, CI Confidence interval, *p* probability value, SD Standard deviation, Other dementias: alcohol-related dementia, dementia in human immunodeficiency virus, dementia in Huntington's disease and dementia in other substance abuse

= 0.12 [0.09, 0.14], $p < .001$). Subgroup analyses revealed that non-English-speaking immigrants were 1.3 years older than non-immigrants on average (Cohen's $d = 0.18$ [0.15, 0.21], $p < .001$), but without significant difference in sex. No significant differences in age and sex were observed between English-speaking immigrants and non-immigrants. There was a significant difference in the prevalence of dementia subtypes between immigrants and non-immigrants but again with small effect sizes (Cramer's $V = 0.03$ [0.01, 0.05], $p < .001$). This effect was primarily driven by differences in mixed dementia and frontal lobe dementia, with immigrants being 1.3% higher in mixed dementia and 0.7% lower in frontal lobe dementia than non-immigrants. There were also significant differences in dementia subtypes in both subgroup comparisons, with mixed dementia being 2.2% higher in non-English speaking immigrants than non-immigrants, and Alzheimer's disease being 4.6% higher in English-speaking immigrants compared to non-Immigrants.

Neuropsychiatric symptoms

There were no differences in the total NPI severity scores and the total number of NPI domains between immigrants and non-immigrants (Table 1). None of the NPI total scores had significant differences between English-speaking immigrants and non-immigrants. However, there was a significant difference in the total number of NPI domains (Cohen's $d = -0.06$ [-0.09, -0.02], $p < .001$) between non-English-speaking immigrants and non-immigrants (See Additional file 1). Individual NPI domain severity and prevalence scores are displayed in Table 2 and Fig. 1, respectively. The most frequently presenting NPS for immigrants and non-immigrant groups were agitation/aggression (87.9% and 86.5%), irritability/liability (65.0% and 64.2%), depression/dysphoria (59.6% and 58.4%), and anxiety (58.1% and 59.5%) (Fig. 1). However, significant differences were observed only for the prevalence of hallucinations (16.0% and 18.3%). In comparing non-English-speaking immigrants and non-immigrants,

Table 2 Individual NPI domain severity scores for non-immigrants, immigrants, NES immigrants, and ES immigrants

Domain	Non-immigrants M(SD)	Immigrants M(SD)	NES immigrants M(SD)	ES immigrants M(SD)
Aberrant Motor Behavior	1.0 (1.2)	1.1 (1.2)	1.0 (1.2)	1.1 (1.2)
Agitation/Aggression	1.9 (1.0)	1.9 (1.0)	1.9 (1.0)	2.0 (1.0)
Anxiety	1.3 (1.2)	1.3 (1.2)	1.2 (1.2)	1.3 (1.2)
Apathy/Indifference	0.7 (1.0)	0.7 (1.0)	0.7 (1.0)	0.7 (1.0)
Appetite and Eating	0.5 (0.9)	0.5 (0.9)	0.5 (0.9)	0.5 (0.9)
Delusions	0.8 (1.1)	0.7 (1.1)	0.7 (1.1)	0.8 (1.1)
Depression/Dysphoria	1.2 (1.1)	1.2 (1.1)	1.2 (1.1)	1.2 (1.1)
Disinhibition	0.9 (1.1)	0.8 (1.1)	0.8 (1.1)	0.9 (1.1)
Elation/Euphoria	0.1 (0.4)	0.1 (0.4)	0.1 (0.4)	0.1 (0.4)
Hallucinations	0.4 (0.8)	0.3 (0.8)	0.3 (0.7)	0.3 (0.8)
Irritability/Lability	1.4 (1.2)	1.4 (1.2)	1.4 (1.2)	1.5 (1.2)
Night-time Behavior	0.8 (1.1)	0.9 (1.1)	0.9 (1.1)	0.9 (1.1)

NPI Neuropsychiatric inventory, NES Non-English-speaking, ES English-speaking, M Mean, SD Standard deviation




Fig. 1 Prevalence of each NPI domain for immigrants and non-immigrants. NPI: neuropsychiatric inventory. Differences between the groups at the .05 level and .001 level are marked with * and **, respectively

non-immigrants had a significantly higher prevalence of delusions, disinhibition, and hallucinations (See Additional file 2). There were no significant differences in individual NPI domain prevalence observed between English-speaking immigrants and non-immigrants (See Additional file 3). The prevalence of each domain for each group with the 95% confidence intervals is shown in Supplementary Table 2 (See Additional file 4).

A logistic regression, controlling for age and sex, examined the relationship between immigrant status and individual NPI domain severity scores (Table 3). There were significant effects for agitation/aggression and hallucinations for immigrant and non-immigrant groups. For every 1-point increase in agitation/aggression, it was 1.08 times more likely that a person was an immigrant. Conversely, for every 1-point increase in hallucinations, it

Table 3 Logistic regression model predicting immigrant status from NPS severity, controlling for age and sex

Term	Immigrants				NES immigrants				ES immigrants						
	B	SE	Z	OR [95% CI]	p	B	SE	Z	OR [95% CI]	p	B	SE	Z	OR [95% CI]	p
(intercept)	-2.27	0.18	-12.53	0.10 [0.07, 0.15]	<.001	-3.33	0.21	-15.68	0.04 [0.02, 0.05]	<.001	-2.15	0.26	-8.16	0.12 [0.07, 0.19]	<.001
Aberrant Motor Behavior	0.03	0.01	2.05	1.03 [1.00, 1.05]	.041	0.01	0.02	0.75	1.01 [0.98, 1.04]	.451	0.05	0.02	2.76	1.06 [1.02, 1.10]	.006
Agitation/Aggression	0.08	0.02	4.54	1.08 [1.05, 1.12]	<.001	0.09	0.02	4.37	1.09 [1.05, 1.14]	<.001	0.06	0.03	2.29	1.06 [1.01, 1.12]	.022
Anxiety	-0.03	0.01	-2.43	0.97 [0.94, 0.99]	.015	-0.06	0.02	-3.39	0.95 [0.92, 0.98]	<.001	0	0.02	0.23	1.00 [0.96, 1.05]	.814
Apathy/Indifference	-0.04	0.02	-2.78	0.96 [0.93, 0.99]	.005	-0.05	0.02	-2.6	0.95 [0.92, 0.99]	.009	-0.04	0.02	-1.58	0.96 [0.92, 1.01]	.114
Appetite and Eating	-0.02	0.02	-1.22	0.98 [0.95, 1.01]	.222	-0.03	0.02	-1.31	0.97 [0.93, 1.01]	.190	-0.01	0.03	-0.53	0.99 [0.94, 1.04]	.599
Delusions	-0.04	0.02	-2.36	0.96 [0.94, 0.99]	.018	-0.05	0.02	-2.9	0.95 [0.92, 0.98]	.004	-0.01	0.02	-0.29	0.99 [0.95, 1.04]	.773
Depression/Dysphoria	0.04	0.01	2.96	1.05 [1.01, 1.08]	.003	0.06	0.02	3.52	1.06 [1.03, 1.10]	<.001	0.01	0.02	0.64	1.01 [0.97, 1.06]	.523
Disinhibition	-0.05	0.01	-3.22	0.95 [0.93, 0.98]	.001	-0.07	0.02	-3.86	0.94 [0.91, 0.97]	<.001	-0.01	0.02	-0.5	0.99 [0.95, 1.03]	.615
Elation/Euphoria	0.03	0.04	0.8	1.03 [0.96, 1.11]	.425	0.01	0.04	0.32	1.01 [0.93, 1.10]	.752	0.05	0.05	0.93	1.05 [0.95, 1.16]	.350
Hallucinations	-0.07	0.02	-3.31	0.93 [0.90, 0.97]	<.001	-0.09	0.02	-3.67	0.91 [0.87, 0.96]	<.001	-0.03	0.03	-1.13	0.97 [0.91, 1.02]	.260
Irritability/Lability	0.02	0.01	1.33	1.02 [0.99, 1.05]	.185	0.01	0.02	0.75	1.01 [0.98, 1.05]	.454	0.03	0.02	1.43	1.03 [0.99, 1.08]	.152
Night-time Behavior	0.04	0.01	2.91	1.04 [1.01, 1.07]	.004	0.05	0.02	3.21	1.05 [1.02, 1.09]	.001	0.02	0.02	0.98	1.02 [0.98, 1.06]	.326
Age	0.02	0	8.78	1.02 [1.01, 1.02]	<.001	0.03	0	10.86	1.03 [1.02, 1.03]	<.001	0	0	1.02	1.00 [1.00, 1.01]	.309
Sex - Male	0.12	0.03	3.88	1.13 [1.06, 1.20]	<.001	0.13	0.04	3.72	1.14 [1.07, 1.23]	<.001	0.09	0.05	2.03	1.10 [1.00, 1.20]	.042

NPS Neuropsychiatric symptoms, NES Non-English-speaking, ES English-speaking, B: unstandardized coefficient, SE Standard error, Z z score for the Wald test, OR Odds ratio, CI Confidence interval, p: probability value. The reference group is non-immigrants

was 1.07 times less likely that a person was an immigrant. In comparing non-English-speaking immigrants and non-immigrants, agitation/aggression, anxiety, depression/dysphoria, disinhibition, and hallucinations showed significant effects (Table 3).

A linear regression model with immigrant status, age, and sex as predictors, was created using total NPI severity scores as the dependent variable. Across all the groups, immigrant status did not predict total NPI severity scores in the model when controlling for age and sex (See Additional file 5).

Caregiver distress

There were no significant differences in total caregiver distress scores between all groups (Table 1). A linear regression model with immigrant status, age, and sex as predictors was created, using total caregiver distress scores as dependent variables. For all groups, immigrant status did not predict total caregiver distress when controlling for age and sex. However, age and sex had significant effects, with caregivers of older people and male referrals having lower distress scores across all comparison groups (See Additional file 6).

Contributing factors to NPS

The prevalence of common contributing factors and two additional factors (language barriers and cultural considerations), which are directly related to our study objectives were compared across groups (Table 4). Only language barriers and cultural considerations accounted for significant differences between the two groups, with 6.5% and 2.2% higher prevalence, respectively for immigrants compared to non-immigrants. Loneliness/boredom, language barriers, and cultural considerations significantly contributed to NPS for non-English-speaking immigrants and non-immigrants comparison, with the largest significant differences being for language barriers (10.0%) and cultural considerations (3.4%). There were no differences in the prevalence of

contributing factors between English-speaking immigrants and non-immigrants.

Discussion

This study compared the clinical and demographic characteristics, prevalence and severity of NPS, and associated caregiver distress for immigrants and non-immigrants living with dementia in RACHs referred to an Australian national dementia support service (DSA). It is important to note that our sample consisted of referrals who exhibited NPS at levels requiring specialist support. We analysed data collected from the first referral to DSA and as such, our findings reflect the level of symptoms requiring support and not the impact of support for NPS. With 24% of referrals from non-English-speaking countries, our sample was comparable to the national record that estimates 23% of Australians aged 65 and over were born in non-English-speaking countries [1]. Additionally, Alzheimer's disease, the most common form of dementia worldwide [47] was the most represented dementia subtype in both immigrant and non-immigrant groups. We found differences between immigrants and non-immigrants in age and dementia subtype, but not sex, partially supporting our first hypothesis. Our second hypothesis was also partially supported as we observed differences in the individual NPI domain severity scores between the two groups, but not in total severity scores. Our third hypothesis was not supported as there were no significant differences in caregiver distress between immigrants and non-immigrants living with dementia. However, there were differences in the prevalence of factors contributing to NPS between the two groups, thus confirming our fourth hypothesis.

Neuropsychiatric symptoms (Hypothesis 2)

Our findings indicated a significantly lower rate of hallucinations for immigrants and non-English-speaking immigrants compared to non-immigrants. Additionally, non-English-speaking immigrants were

Table 4 Differences in the prevalence of commonly identified contributing factors to NPS for non-immigrants and immigrant groups

Contributing factor	Non-immigrants %	Immigrants Difference [95% CI], <i>p</i>	NES immigrants Difference [95% CI], <i>p</i>	ES immigrants Difference [95% CI], <i>p</i>
Pain	58.6%	1.5% [0.1, 2.9], <i>p</i> =.031	2.6% [1.0, 4.2], <i>p</i> =.001	-0.6% [-2.7, 1.5], <i>p</i> =.600
Carer approach	36.3%	-0.1% [-1.5, 1.2], <i>p</i> =.861	0.1% [-1.5, 1.6], <i>p</i> =.927	-0.5% [-2.6, 1.5], <i>p</i> =.609
Mood disorders	26.5%	-0.6% [-1.9, 0.6], <i>p</i> =.334	0.3% [-1.2, 1.7], <i>p</i> =.734	-2.3% [-4.1, -0.5], <i>p</i> =.015
Over/under stimulation	15.5%	0.1% [-0.9, 1.2], <i>p</i> =.816	-0.2% [-1.4, 0.9], <i>p</i> =.711	0.8% [-0.7, 2.4], <i>p</i> =.295
Loneliness/boredom	15.2%	1.4% [0.4, 2.5], <i>p</i> =.006	2.0% [0.8, 3.2], <i>p</i> <.001	0.4% [-1.2, 1.9], <i>p</i> =.652
Language barrier	0.2%	6.5% [6.0, 7.1], <i>p</i> <.001	10.0% [9.1, 10.8], <i>p</i> <.001	-0.0% [-0.3, 0.2], <i>p</i> =.861
Cultural considerations	0.5%	2.2% [1.8, 2.6], <i>p</i> <.001	3.4% [2.8, 3.9], <i>p</i> <.001	-0.1% [-0.3, -0.0], <i>p</i> =.190

NPS Neuropsychiatric symptoms, NES Non-English-speaking, ES English-speaking, CI Confidence interval, *p*: probability

significantly lower on delusions and disinhibition than non-immigrants. The literature reports different types of NPS and higher frequencies of some NPS for ethnic minorities compared with their counterparts or native populations. Salazar and colleagues [11] compared NPI scores for ethnic groups living with Alzheimer's disease in the community ($n = 975$) reporting a higher frequency of hallucinations, night-time behaviour, and elation for people from Hispanic backgrounds. However, Wilson and colleagues [48] reported no differences in NPS presentation for Latino older adults and non-Latino older adults living with dementia in Brazil, reasoning that NPS are driven primarily by brain structure and functional changes than by sociodemographic characteristics. However, the findings of these studies are different to our study findings likely reflecting the differences in the study methodology (e.g., study design) and sample characteristics (e.g., sample size).

Our study found no differences in total NPI severity scores between immigrants and non-immigrants. This is partially due to differences in the direction of individual domains cancelling out the effect in the total severity scores, where immigrants were more likely to be higher on agitation/aggression but lower on hallucinations, compared to non-immigrants. This also applies to the comparison for non-English-speaking immigrants, which were more likely to be higher on agitation/aggression and depression/dysphoria, but lower on anxiety, disinhibition, and hallucinations. The higher severity of agitation/aggression is likely driven by communication difficulties as there was no difference for the English-speaking immigrant groups comparison. Cognitive decline can impair both the ability to express and comprehend spoken language and people living with dementia who have English as their additional language may lose their ability to communicate in English and subsequently use their first language as the primary language of communication [7, 8]. A study comparing verbal communication and psychiatric medication use by Greek and Italian residents with dementia in ethno-specific and mainstream RACHs in Australia found that communication difficulties were linked with higher rates of psychiatric prescriptions, ostensibly for managing NPS such as agitation and aggression [49]. There were no significant differences in NPI prevalence and total NPI severity between English-speaking immigrants and non-immigrants highlighting the impact of language and culture in NPS presentation in people living with dementia. Unlike non-English-speaking immigrants, English-speaking immigrants may not have experienced communication and/or cultural difficulties with their formal caregivers.

Caregiver distress (Hypothesis 3)

There was no difference in caregiver distress between caregivers of immigrants and non-immigrants living with dementia. This may be because our sample comprised of referrals for NPS support from RACHs and as such similar levels of distress have been reported by the caregivers across the groups. Further, formal caregivers working in RACHs may have had training on managing NPS and regularly supported such symptoms. Current literature suggests longer durations of care [50, 51] and more caregiving activities [50] are linked to increased informal caregiver distress and burden.

Contributing factors to NPS (Hypothesis 4)

Pain was the top contributing factor for NPS for both immigrant and non-immigrant groups, echoing previous findings that pain is commonly associated with NPS, particularly agitation/aggression [27, 52]. Given that pain is a frequently occurring contributing factor to NPS, future studies should focus on examining the association between pain and NPS. We found a higher rate of language barriers and cultural considerations for non-English-speaking immigrants compared to non-immigrants. Such findings support what is already known about the impact of culture on the patterns of NPS presentations for people living with dementia and where there is a lack of cultural considerations and communication options available in RACHs this may exacerbate agitation [53, 54]. For this study pain was assessed using PainChek®, an observational pain assessment instrument that alleviates the need for verbal communication and self-reporting of pain. While such tools are important for people who have lost language expression, lack of such options can be difficult for carers in understanding the expressions of pain by residents from immigrant backgrounds. Data from the Australian Royal Commission into Aged Care Quality and Safety provides examples of how care workers' inability to recognize pain expression of residents from immigrant backgrounds who have lost their English language led to poor care outcomes [55].

Strengths and limitations

This is one of the largest population-based studies and to the best of our knowledge, the first in Australia, to comparatively describe the clinical, demographic, and NPS characteristics of immigrants and non-immigrants living with dementia referred to an Australian dementia behaviour support service. The strengths of this study include the use of a large representative sample of people living with dementia aged 65 years and above, strengthening

the generalisability of our findings to older people living with dementia who received external support for NPS.

This study has some limitations that need to be noted. This was a retrospective observational cross-sectional study and bound by the limitations of observational studies such as the inability to attribute causation [56]. Our findings may be limited to participants aged 65 years and above and living in RACHs, although, there are lessons for community/home care that are relevant. Constrained by existing data variables, we could not investigate the potential confounding effects of racial and ethnic variations, medical conditions, medication profiles, and other covariates such as levels of cognitive function or activities of daily living. We determined immigrant status by documented country of birth, and we did not have access to the duration of stay in Australia (e.g., long-term immigrants or short-term immigrants) or English language proficiency; as such, we cannot be certain that people coming from non-English-speaking countries were speaking English as a first language or had developed and/or lost English as their second language. In general, contributing factors to NPS are complex and their identification is complicated by the (co-)presence of many behaviours often having numerous causes [57]. These factors are of various types including biological, psychological, social and/or environmental factors [25, 57]. Therefore, identifying any factor requires a robust and comprehensive assessment process and obtaining information through multiple sources including standardised and validated instruments, observation, interview, and review of clinical notes and social history. Due to the complexity of contributing factors and their assessment, their identification may be subject to bias and should be interpreted within this context. However, the risk of such bias is mitigated by the extensive clinical expertise and experience of DSA consultants. We have also not performed an adjustment for multiple testing due to the exploratory nature of our study. However, we have interpreted the significance of our findings in relation to a stricter alpha value of $<.001$ rather than $<.05$. Further, due to the exploratory nature of the hypotheses, our findings need to be replicated in future studies to confirm any conclusions.

Conclusion

Our findings reveal a mixed picture of NPS between immigrants and non-immigrants living with dementia that may give policymakers and service providers a better picture of the potential differences in demographic characteristics, prevalence and severity of NPS, and factors contributing to NPS. This may be useful for future service planning, in particular, but not limited to RACHs. Our findings call for increased awareness and education

on the impact of culture and language on NPS for people receiving residential care. Future studies where variables such as length of stay in Australia and English language proficiency can be determined are needed as are those with diverse study designs (e.g., longitudinal studies) to learn more about NPS presentations for different immigrant groups (e.g., first- and second-generation immigrants). Studies investigating the association between NPS and pain intensity in people living with dementia, with different migration histories, will also contribute to the appropriate management of NPS.

Abbreviations

DBMAS	Dementia Management Advisory Service
DSA	Dementia Support Australia
NPI	Neuropsychiatric Inventory
NPI-NH	Neuropsychiatry Inventory Nursing Home
NPI-Q	Neuropsychiatry Inventory Questionnaire
NPS	Neuropsychiatric symptoms
RACH(s)	Residential Aged Care Home(s)
SBRT	Severe Behaviour Response Team

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12877-023-04447-3>.

Additional file 1: Supplementary Table 1. Effect sizes, 95% CI and p-values for the descriptive characteristics of NES and ES immigrants.

Additional file 2: Supplementary Figure 1. Prevalence of each NPI domain for NES immigrants and non-immigrants. NES: non-English-speaking; NPI: neuropsychiatric inventory. Differences between the groups at the .05 level and .001 level are marked with * and **, respectively.

Additional file 3: Supplementary Figure 2. Prevalence of each NPI domain for ES immigrants and non-immigrants. NPI: neuropsychiatric inventory; ES: English-speaking. Differences between the groups at the .05 level are marked with *.

Additional file 4: Supplementary Table 2. Prevalence of each NPS domain for each group with the 95% confidence intervals.

Additional file 5: Supplementary Table 3. Linear regression model predicting total NPI severity scores from immigrant status controlling for age and sex.

Additional file 6: Supplementary Table 4. Linear regression model predicting total caregiver distress scores from immigrant status controlling for age and sex.

Acknowledgements

The authors thank The Dementia Centre, HammondCare, and Dementia Support Australia (DSA) for allowing us to use their data. We acknowledge the people living with dementia and their families who have been supported by DSA programs.

Authors' contributions

PC (first author) contributed to the study concept and design; contributed to data acquisition, analysis, and interpretation; conducted the literature review and drafted the manuscript. MA contributed to the study concept and design; contributed to data acquisition, analysis, and interpretation; and provided a critical review and comment on the manuscript. PC (third author) contributed to the study concept and design; contributed to data acquisition, analysis, and interpretation; and provided a critical review and comment on the manuscript. DW processed the data; performed statistical analysis; and provided a critical review and comment on the manuscript. TM contributed

to data acquisition, analysis, and interpretation; and provided a critical review and comment on the manuscript. DP contributed to the study concept and design; contributed to data acquisition, analysis, and interpretation; and provided a critical review and comment on the manuscript. All authors proofread and approved the final version of the manuscript.

Funding

The authors did not receive any specific funding for this study.

Availability of data and materials

Restrictions apply to the availability of data generated or analysed during this study to preserve data privacy. The corresponding author will, on request, provide details on the restrictions and any conditions under which access to some data may be provided.

Declarations

Ethics approval and consent to participate

The study was conducted in compliance with the approval granted by the University of New South Wales Human Research Committee (HC190049) which granted a waiver of informed consent for the study as the data were deidentified and analysed retrospectively and anonymously. The subsequent reciprocal approval (2022-03715) was granted by the Human Research Ethics Committee of Edith Cowan University via an exempt pathway. The study was also approved (R257) by the Research Governance Office of HammondCare. A data custodian at DSA handled the data according to ethical principles set out by the Declaration of Helsinki, and the National Health and Medical Research Council's National Statement on Ethical Conduct in Human Research.

Consent for publication

Not applicable.

Competing interests

The author(s) declared the following potential conflicts of interest with respect to the research, authorship and/or publication of this article: MA, DW and TM are staff members of The Dementia Centre, HammondCare; a research, education, and consultancy arm of HammondCare, an independent Christian charity which auspices the DSA programs – the Dementia Behavior Management Advisory Service (DBMAS) and the Severe Behavior Response Team (SBRT). MA is one of the originators of the PainChek instrument, which is marketed by PainChek Ltd. (ASX: PCK). He is also a shareholder of PainChek Ltd. He previously held the position of Senior Research Scientist (October 2018–May 2020) at PainChek Ltd. He had a granted patent titled 'A pain assessment method and system; PCT/AU2015/000501' in Australia, China, Japan, and the United States, which was assigned to PainChek Ltd. The other authors declare that they have no competing interests.

Author details

¹Centre for Research in Aged Care, School of Nursing and Midwifery, Edith Cowan University, 270 Joondalup Drive, Joondalup, Western Australia 6027, Australia. ²The Dementia Centre, HammondCare, Osborne Park, Western Australia, Australia. ³Curtin Medical School, Faculty of Health Sciences, Curtin University, Bentley, Western Australia, Australia. ⁴Sydney Pharmacy School, Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales, Australia. ⁵The Dementia Centre, HammondCare, St Leonards, New South Wales, Australia.

Received: 17 April 2023 Accepted: 1 November 2023

Published online: 10 November 2023

References

1. Australian Institute of Health and Welfare. Dementia in Australia. <https://www.aihw.gov.au/reports/dementia/dementia-in-aus/contents/dementia-in-vulnerable-groups/dementia-among-people-from-culturally-and-linguistically-diverse-backgrounds> (2023). Accessed 20 Feb 2023.
2. Australian Institute of Health and Welfare. People using aged care. <https://www.gen-agedcaredata.gov.au/Topics/People-using-aged-care#>
3. People%20from%20culturally%20and%20linguistically%20diverse%20backgrounds (2022). Accessed 20 Feb 2023.
4. Moon H, Badana ANS, Hwang SY, Sears JS, Haley WE. Dementia Prevalence in Older Adults: Variation by Race/Ethnicity and Immigrant Status. *Am J Geriatr Psychiatry*. 2019;27(3):241–50.
5. Parlevliet JL, Uysal-Bozkir Ö, Goudsmit M, van Campen JP, Kok RM, Ter Riet G, et al. Prevalence of mild cognitive impairment and dementia in older non-western immigrants in the Netherlands: a cross-sectional study. *Int J Geriatr Psychiatry*. 2016;31(9):1040–9.
6. Selten J-P, Termorshuizen F, van Sonsbeek M, Bogers J, Schmand B. Migration and dementia: a meta-analysis of epidemiological studies in Europe. *Psychol Med*. 2021;51(11):1838–45.
7. McNabney MK, Green AR, Burke M, Le ST, Butler D, Chun AK, et al. Complexities of care: Common components of models of care in geriatrics. *J Am Geriatr Soc*. 2022;70(7):1960–72.
8. Sagbakken M, Spilker RS, Ingebretsen R. Dementia and Migration: Family Care Patterns Merging With Public Care Services. *Qual Health Res*. 2018;28(1):16–29.
9. Tipping SA, Whiteside M. Language Reversion among People with Dementia from Culturally and Linguistically Diverse Backgrounds: The Family Experience. *Aust Soc Work*. 2015;68(2):184–97.
10. Söderman M, Rosendahl SP. Caring for Ethnic Older People Living with Dementia - Experiences of Nursing Staff. *J Cross Cult Gerontol*. 2016;31(3):311–26.
11. Makimoto K, Kang Y, Kobayashi S, Liao XY, Panuthai S, Sung HC, et al. Prevalence of behavioural and psychological symptoms of dementia in cognitively impaired elderly residents of long-term care facilities in East Asia: a cross-sectional study. *Psychogeriatr*. 2019;19(2):171–80.
12. Salazar R, Dwivedi AK, Royall DR. Cross-Ethnic Differences in the Severity of Neuropsychiatric Symptoms in Persons With Mild Cognitive Impairment and Alzheimer's Disease. *J Neuropsychiatry Clin Neurosci*. 2017;29(1):13–21.
13. Lennon JC, Aita SL, Bene VAD, Rhoads T, Resch ZJ, Eloi JM, et al. Black and White individuals differ in dementia prevalence, risk factors, and symptomatic presentation. *Alzheimers Dement*. 2022;18(8):1461–71.
14. Goudsmit M, van de Vorst I, van Campen J, Parlevliet J, Schmand B. Clinical characteristics and presenting symptoms of dementia - a case-control study of older ethnic minority patients in a Dutch urban memory clinic. *Aging Ment Health*. 2022;26(11):2277–84.
15. Loeffler DA. Modifiable, Non-Modifiable, and Clinical Factors Associated with Progression of Alzheimer's Disease. *J Alzheimers Dis*. 2021;80(1):1–27.
16. Dietlin S, Soto M, Kiyasova V, Pueyo M, de Mauleon A, Delrieu J, et al. Neuropsychiatric Symptoms and Risk of Progression to Alzheimer's Disease Among Mild Cognitive Impairment Subjects. *J Alzheimers Dis*. 2019;70(1):25–34.
17. Hongisto K, Hallikainen I, Selander T, Törmälehto S, Väättäinen S, Martikainen J, et al. Quality of life in relation to neuropsychiatric symptoms in Alzheimer's disease: 5-year prospective ALSOVA cohort study. *Int J Geriatr Psychiatry*. 2018;33(1):47–57.
18. Choi A, Martyr A, Clare L, Fossey J, Khan Z, McDermid J, et al. Impact of Psychotic Symptoms and Concurrent Neuropsychiatric Symptoms on the Quality of Life of People With Dementia Living in Nursing Homes. *J Am Med Dir Assoc*. 2022;23(9):1474–9.e1.
19. Truzzi A, Valente L, Engelhardt E, Laks J. The association between caregiver distress and individual neuropsychiatric symptoms of dementia. *Dement Neuropsychol*. 2013;7(3):286–91.
20. Delfino LL, Komatsu RS, Komatsu C, Neri AL, Cachioni M. Neuropsychiatric symptoms associated with family caregiver burden and depression. *Dement Neuropsychol*. 2021;15(1):128–35.
21. Terum TM, Andersen JR, Rongve A, Aarsland D, Svendsboe EJ, Testad I. The relationship of specific items on the Neuropsychiatric Inventory to caregiver burden in dementia: a systematic review. *Int J Geriatr Psychiatry*. 2017;32(7):703–17.
22. Koyama A, Matsushita M, Hashimoto M, Fujise N, Ishikawa T, Tanaka H, et al. Mental health among younger and older caregivers of dementia patients. *Psychogeriatr*. 2017;17(2):108–14.
23. Yang XQ, Vedel I, Khanassov V. The Cultural Diversity of Dementia Patients and Caregivers in Primary Care Case Management: a Pilot Mixed Methods Study. *Can Geriatr J*. 2021;24(3):184–94.

23. Chejor P, Laging B, Whitehead L, Porock D. Experiences of older immigrants living with dementia and their carers: a systematic review and meta-synthesis. *BMJ Open*. 2022;12(5): e059783.
24. Fauth EB, Gibbons A. Which behavioral and psychological symptoms of dementia are the most problematic? Variability by prevalence, intensity, distress ratings, and associations with caregiver depressive symptoms. *Int J Geriatr Psychiatry*. 2014;29(3):263–71.
25. Kales HC, Gitlin LN, Lyketsos CG. Assessment and management of behavioral and psychological symptoms of dementia. *BMJ*. 2015;350: h369.
26. Tran P, Schmidt K, Gallo J, Tuppo E, Scheinthal S, Chopra A, et al. Neuropsychiatric Symptoms and Medical Illness in Patients with Dementia: An Exploratory Study. *J Osteopath Med*. 2006;106(7):412–4.
27. Atee M, Morris T, Macfarlane S, Cunningham C. Pain in Dementia: Prevalence and Association With Neuropsychiatric Behaviors. *J Pain Symptom Manage*. 2021;61(6):1215–26.
28. Giménez-Llort L, Bernal ML, Docking R, Muntan-Soria A, Torres-Lista V, Bulbena A, et al. Pain in Older Adults With Dementia: A Survey in Spain. *Front Neurol*. 2020;11:592366.
29. Sink KM, Covinsky KE, Barnes DE, Newcomer RJ, Yaffe K. Caregiver characteristics are associated with neuropsychiatric symptoms of dementia. *J Am Geriatr Soc*. 2006;54(5):796–803.
30. Smith M, Hall GR, Gerdner L, Buckwalter KC. Application of the Progressively Lowered Stress Threshold Model across the continuum of care. *Nurs Clin North Am*. 2006;41(1):57–81, vi.
31. Loi SM, Atee M, Morris T, Whiting D, Macfarlane S, Cunningham C, et al. Clinico-demographics of people with younger-onset dementia and neuropsychiatric symptoms referred to an Australian dementia support service: A comparison study with older-onset dementia. *Aust N Z J Psychiatry*. 2022;56(12):1653–63.
32. Zhou G, Liu S, Yu X, Zhao X, Ma L, Shan P. High prevalence of sleep disorders and behavioral and psychological symptoms of dementia in late-onset Alzheimer disease: A study in Eastern China. *Medicine (Baltimore)*. 2019;98(50): e18405.
33. Chang KH, Huang CC, Chen CM, Wu HC, Kuo HC. Differences in Clinical Presentation of Behavioral and Psychological Symptoms of Dementia in Alzheimer's Disease According to Sex and Education Level. *J Alzheimers Dis*. 2020;78(2):711–9.
34. Temple J, Wilson T, Brijnath B, Radford K, LoGiudice D, Utomo A, et al. The role of demographic change in explaining the growth of Australia's older migrant population living with dementia, 2016–2051. *Aust N Z J Pub Health*. 2022;46(5):661–7.
35. Macfarlane S, Atee M, Morris T, Whiting D, Healy M, Alford M, et al. Evaluating the clinical impact of national dementia behaviour support programs on neuropsychiatric outcomes in Australia. *Front Psychiatry*. 2021;12: 652254.
36. Wood S, Cummings JL, Hsu MA, Barclay T, Wheatley MV, Yarema KT, et al. The use of the neuropsychiatric inventory in nursing home residents. Characterization and measurement. *Am J Geriatr Psychiatry*. 2000;8(1):75–83.
37. Kaufer DI, Cummings JL, Ketchel P, Smith V, MacMillan A, Shelley T, et al. Validation of the NPI-Q, a brief clinical form of the Neuropsychiatric Inventory. *J Neuropsychiatry Clin Neurosci*. 2000;12(2):233–9.
38. Atee M, Hoti K, Hughes JD. A Technical Note on the PainChek™ System: A Web Portal and Mobile Medical Device for Assessing Pain in People With Dementia. *Front Aging Neurosci*. 2018;10:117.
39. Atee M, Hoti K, Parsons R, Hughes JD. Pain Assessment in Dementia: Evaluation of a Point-of-Care Technological Solution. *J Alzheimer's Dis*. 2017;60:137–50.
40. Atee M, Hoti K, Hughes JD. Psychometric Evaluation of the Electronic Pain Assessment Tool: An Innovative Instrument for Individuals with Moderate-to-Severe Dementia. *Dement Geriatr Cogn Disord*. 2017;44(5–6):256–67.
41. World Population Review. English Speaking Countries. <https://worldpopulationreview.com/country-rankings/english-speaking-countries> (2022). Accessed 20 Nov 2022.
42. R Core Team. R: A Language and Environment for Statistical Computing. 4.2.2 ed. Vienna, Austria 2022.
43. Benjamin DJ, Berger JO, Johannesson M, Nosek BA, Wagenmakers EJ, Berk R, et al. Redefine statistical significance. *Nat Hum Behav*. 2018;2(1):6–10.
44. Sloan JA, Halyard M, El Naqa I, Mayo C. Lessons From Large-Scale Collection of Patient-Reported Outcomes: Implications for Big Data Aggregation and Analytics. *Int J Radiat Oncol Biol Phys*. 2016;95(3):922–9.
45. Kim HY. Statistical notes for clinical researchers: Chi-squared test and Fisher's exact test. *Restor Dent Endod*. 2017;42(2):152–5.
46. Cohen J. Statistical power analysis for the behavioral sciences. 2nd ed. Hillsdale: Erlbaum Associates; 1988.
47. World Health Organization. Dementia Key facts. <https://www.who.int/news-room/fact-sheets/detail/dementia> (2022). Accessed 27 Feb 2023.
48. Wilson RS, Capuano AW, Sampaio C, Leurgans SE, Barnes LL, Farfel JM, et al. Neuropsychiatric symptoms in Brazilians with mild cognitive impairment and dementia. *Alzheimers Dement (Amst)*. 2021; 13(1):[e12219 p]; <https://doi.org/10.1002/dad2.12219>.
49. Runci SJ, Eppingstall BJ, O'Connor DW. A comparison of verbal communication and psychiatric medication use by Greek and Italian residents with dementia in Australian ethno-specific and mainstream aged care facilities. *Int Psychogeriatr*. 2012;24(5):733–41.
50. Pinquart M, Sörensen S. Gender Differences in Caregiver Stressors, Social Resources, and Health: An Updated Meta-Analysis. *J Gerontol B Psychol Sci Soc Sci*. 2006;61:P33–45.
51. Chakraborty R, Jana A, Vibhute VM. Caregiving: a risk factor of poor health and depression among informal caregivers in India- A comparative analysis. *BMC Public Health*. 2023;23(1):42.
52. Ahn H, Horgas A. The relationship between pain and disruptive behaviors in nursing home residents with dementia. *BMC Geriatr*. 2013;13:14.
53. Hendrie HC, Baiyewu O, Eldemire D, Prince C. Cross-cultural perspectives: Caribbean, Native American, and Yoruba. *Int Psychogeriatr*. 1996;8(Suppl 3):483–6.
54. Mangone CA, Bueno AA, Allegri R, Arizaga R, Nitrini R, Ventura R, et al. Behavioral and psychological symptoms of dementia in Latin America. *Int Psychogeriatr*. 2000;12(S1):415–8.
55. Cain P, Alan J, Porock D. Emergency department transfers from residential aged care: what can we learn from secondary qualitative analysis of Australian Royal Commission data? *BMJ Open*. 2022;12(9): e063790.
56. Wang MTM, Bolland MJ, Grey A. Reporting of Limitations of Observational Research. *JAMA Intern Med*. 2015;175(9):1571–2.
57. Tible OP, Riese F, Savaskan E, von Gunten A. Best practice in the management of behavioural and psychological symptoms of dementia. *Ther Adv Neurol Disord*. 2017;10(8):297–309.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

