

2016

## Heat shock gene Sirtuin 1 regulates post-prandial lipid metabolism with relevance to nutrition and appetite regulation in diabetes

Ian J. Martins  
*Edith Cowan University*

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworkspost2013>



Part of the [Medicine and Health Sciences Commons](#)

---

[10.15344/2394-1499/2016/120](https://ro.ecu.edu.au/ecuworkspost2013/3554)

Martins, I. J. (2016). Heat shock gene Sirtuin 1 regulates post-prandial lipid metabolism with relevance to nutrition and appetite regulation in diabetes. *Int J Diab Clin Diagn*, 3, 20.

<http://dx.doi.org/10.15344/2394-1499/2016/120>

This Other is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/3554>

# Heat Shock Gene Sirtuin 1 Regulates Post-Prandial Lipid Metabolism with Relevance to Nutrition and Appetite Regulation in Diabetes

Ian James Martins<sup>1,2,3\*</sup>

<sup>1</sup>Centre of Excellence in Alzheimer's Disease Research and Care, School of Medical Sciences, , Edith Cowan University, 270 Joondalup Drive, Joondalup, 6027, Australia

<sup>2</sup>School of Psychiatry and Clinical Neurosciences, The University of Western Australia, Nedlands, 6009, Australia

<sup>3</sup>McCusker Alzheimer's Research Foundation, Holywood Medical Centre, 85 Monash Avenue, Suite 22, Nedlands, 6009, Australia

New discoveries in medicine are required to understand the importance of appetite regulation that is associated with the overconsumption of food in Type 2 and Type 3 diabetes. Food restriction in diabetes is essential to maintain the hepatic metabolism of dietary fat with relevance to defective post-prandial lipid metabolism and to the global non alcoholic fatty liver disease (NAFLD) epidemic [1,2]. Premature brain aging has become important with the development of Type 3 diabetes and Alzheimer's disease [3] that is associated with repression of the anti-aging gene Sirtuin 1 (Sirt 1) relevant to post-prandial lipid metabolism, amyloid beta metabolism (peptide involved in amyloid beta plaques) and circadian rhythm abnormalities in the brain biological clock associated with the development of NAFLD. Nutritional interventions such as very low carbohydrate diets have become important to diabetes (Figure 1) to reverse defective post-prandial lipid and amyloid beta metabolism without atherogenic lipoprotein formation [4,5] with the prevention of accelerated atherosclerosis in various communities. Western diets that are high in fat and glucose are linked to diabetes and NAFLD with anti-aging gene Sirt 1 transcriptional dysregulation [6] in cell and tissues associated with, hyperglycemia, mitochondrial apoptosis and delayed hepatic fat and amyloid beta metabolism (Figure 1).

## Publication History:

Received: September 15, 2016

Accepted: November 01, 2016

Published: November 03, 2016

## Keywords:

Amyloid beta, Anti-aging genes, diet, Heat shock gene, Heat shock proteins, NAFLD, Ontogeny, Post-prandial, Sirtuin 1, Transcription factors

proteins facilitate the rapid metabolism by the liver of amyloid beta by preventing its misfolding and aggregation in the brain [3,13,14]. Sirt 1 is also sensitive to  $\alpha$ -synuclein metabolism [15] and relevant to temperature alterations in  $\alpha$ -synuclein oligomer and amyloid beta formation [16, 17].

Down regulation of Sirt 1 expression and activity disturbs the nuclear and mitochondria interactions with effects on the metabolism of fatty acids, glucose and amyloid beta metabolism in diabetes [16, 17]. Heat shock protein (HSP) induces endoplasmic reticulum stress (ER) stress with delayed metabolism of amyloid beta/  $\alpha$ -synuclein oligomers associated with liver disease linked to NAFLD and neurodegenerative diseases. ER stress is associated with programmed cell death with relevance to mitochondrial apoptosis, defective post-prandial lipid metabolism and NAFLD [18-22] with Sirt 1 regulation of PGC1 associated with mitochondrial biogenesis after ingestion of various fat diets. Sirt 1 regulates HSP by deacetylation of heat shock factor (HSF) via peroxisome proliferator-activated receptor gamma co activator 1-alpha (PGC1 $\alpha$ ) as a critical repressor of HSF1-mediated transcriptional programs [23,24]. Sirt 1 is involved in body temperature regulation of the mammalian target of rapamycin (mTOR) signaling through the tumor suppressor tuberous sclerosis complex 1 with relevance to the expression of hepatic PGC-1 $\alpha$  and fibroblast growth factor 21 (FGF21) [25-29].

Sirt 1 regulation of cell senescence in diabetes involves cell ontogeny with transcriptional ontogeny defective via the transcriptional factor p53 involved in the regulation of various other anti-aging genes [30-35] such as klotho, p66shc, FOXO3a, micro RNA 34a [18] and transcription factors involve post-prandial lipid metabolism, metabolic activity, insulin resistance, cellular ontogeny, inflammation

**\*Corresponding Author:** Dr. Ian James Martins, School of Medical Sciences, Edith Cowan University, 270 Joondalup Drive, Joondalup, Western Australia 6027, Australia, Tel: +61863042574, E-mail: [i.martins@ecu.edu.au](mailto:i.martins@ecu.edu.au)

**Citation:** Martins IJ (2016) Heat Shock Gene Sirtuin 1 Regulates Post-Prandial Lipid Metabolism with Relevance to Nutrition and Appetite Regulation in Diabetes. Int J Diabetes Clin Diagn 3: 120. doi: <https://doi.org/10.15344/2394-1499/2016/120>

**Copyright:** © 2016 Martins. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: In diabetes and neurodegenerative diseases nutritional interventions are required to activate the anti-aging gene Sirt 1 and prevent defective liver lipid metabolism (NAFLD) and amyloid beta metabolism. Diets that are low in fat such as very low carbohydrate diets activate the calorie sensitive gene Sirt 1 with glucose metabolism connected to accelerated hepatic lipid/amyloid beta metabolism in diabetes and neurodegenerative diseases.

Accelerated aging that disturbs the brain to liver crosstalk [6,7] involves the calorie sensitive gene Sirtuin 1 (Sirt 1) that is a nicotinamide adenine dependent dinucleotide class III histone deacetylase involved in the prevention of defective post-prandial metabolism and NAFLD [8-10]. The intake of fat is sensitive to the regulation of the heat shock gene Sirt 1 that is responsible for the metabolism of heat shock proteins (HSP) [3] that are produced by living cells in response to temperature regulation above physiological levels [11, 12]. Heat shock

and xenobiotic metabolism [18,36] (Figure 2). In the developing and developed world diabetic treatment has become important with defective Sirt 1 gene expression determined by miR-34a [18, 36] related to defective cell proliferation in the brain and the liver (Figure 2). Defective thermoregulation may be relevant to ingestion of food (Sirt 1 defective) with inappropriate post-prandial lipid and amyloid beta metabolism that inactivate magnesium therapy that may now be relevant to HSP and amyloid beta metabolism with relevance to myocardial infarction [3,37].




Figure 2: Appetite regulation in diabetes involve the maintenance of the heat shock gene Sirtuin 1 that is essential for thermo regulation function and hepatic lipid metabolism. The repression of the anti-aging gene Sirt 1 involves p53/mTOR dysregulation of the other anti-aging genes (klotho, p66shc, FOXO3a) associated with hepatic lipid, heat shock protein and amyloid beta metabolism. FGF 21 treatment in diabetes has become important to NAFLD [38] and myocardial infarction treatment [39, 40] but defective cell transcriptional ontogeny in the liver and brain related to mi-34a inhibition of Sirtuin 1/transcription factors interactions inactivates hepatic lipid metabolism with the induction of NAFLD, defective amyloid beta metabolism and neurodegenerative disease.

Food restriction and fasting regulate Sirt 1 and FGF21 involved in the prevention of the metabolic syndrome and maintenance of high density lipoprotein levels [39-41]. FGF21 therapy [42, 43] in Type 2 and Type 3 diabetes may be ineffective with the development of cardiac ageing with relevance to core body temperature regulation that is determined by mi-34a/Sirt 1 gene expression with low adiponectin levels [36,44] (Figure 2). Healthy diets that maintain Sirt 1 activity in the brain and liver have become important to many diabetics in the developing and developed world with relevance to food technology that involve the hepatic metabolism of bacterial lipopolysaccharides (LPS) that has become important to the reversal of Type 2 diabetes and Type 3 diabetes [45, 46]. Plasma LPS and xenobiotic levels [47] have risen in various developed countries and consumption of activators such as leucine, pyruvic acid and magnesium may

supersede LPS inhibition of Sirt 1 [18, 37] with prevention of defective cell transcriptional ontogeny and membrane transformations in the brain and liver [45, 46]. Appetite dysregulation in diabetes may require these activators of Sirt 1 with relevance to the maintenance of neurons and the treatment of Type 3 diabetes with defective apelinergic system that involves thermo dysregulation.

## Conclusion

Dyslipidemia is one of the key risk factors for cardiovascular disease in diabetes. The management of dyslipidemia in diabetes continues to remain controversial and improvements in the characteristic diabetic dyslipidemia of high triglyceride and low HDL may not indicate that defective cell ontogeny is underway early in life. Diabetes and defective hepatic cell transcriptional programs induce delayed post-prandial lipid metabolism associated with Western diets rich in fat and glucose. The clinical management of diabetes in the young and elderly now not only involves appetite regulation with calorie restricted diets that maintain the heat shock gene Sirt 1 expression but also careful core body temperature (37°C) to activate hepatic and brain Sirt 1. The biological active release of FGF21 is connected to Sirt 1 activation and glucose homeostasis with relevance to treatment of dyslipidemia, NAFLD, cardiovascular disease and neurodegenerative diseases. Consumption of Sirt 1 inhibitors such as alcohol, suramin and palmitic acid should be avoided to prevent defective liver and brain cell ontogeny in the young and the elderly to prevent hyperglycemia induced oxidative stress and myocardial infarction.

## Funding

This work was supported by grants from Edith Cowan University, the McCusker Alzheimer's Research Foundation and the National Health and Medical Research Council.

## References

1. Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, et al. (2016) Global epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology* 64: 73-84.
2. Martins IJ (2014) Induction of NAFLD with Increased Risk of Obesity and Chronic Diseases in Developed Countries. *Open J Endocr Metab Dis* 4: 90-110.
3. Martins IJ (2016) Type 3 diabetes with links to NAFLD and Other Chronic Diseases in the Western World. *International J Diab* 1: 1-5.
4. Chehade JM, Gladysz M, Mooradian AD (2013) Dyslipidemia in type 2 diabetes: prevalence, pathophysiology, and management. *Drugs* 73: 327-339.
5. Boekholdt SM, Hovingh GK, Mora S, Arsenault BJ, Amarenco P4, et al. (2014) Very low levels of atherogenic lipoproteins and the risk for cardiovascular events: a meta-analysis of statin trials. *J Am Coll Cardiol* 64: 485-494.
6. Martins IJ (2016) Diet and Nutrition reverse Type 3 Diabetes and Accelerated Aging linked to Global chronic diseases. *J Diab Res Ther* 2.
7. Liang Q, Zhong L, Zhang J, Wang Y, Bornstein SR, et al. (2014) FGF21 maintains glucose homeostasis by mediating the cross talk between liver and brain during prolonged fasting. *Diabetes* 63: 4064-4075.
8. Bordone L, Cohen D, Robinson A, Motta MC, van Veen E, et al. (2007) SIRT1 transgenic mice show phenotypes resembling calorie restriction. *Aging Cell* 6: 759-767.
9. Xu F, Gao Z, Zhang J, Rivera CA, Yin J, et al. (2010) Lack of SIRT1 (Mammalian Sirtuin 1) activity leads to liver steatosis in the SIRT1<sup>+/-</sup> mice: a role of lipid mobilization and inflammation. *Endocrinology* 151: 2504-2514.
10. Purushotham A, Xu Q, Li X (2012) Systemic SIRT1 insufficiency results in disruption of energy homeostasis and steroid hormone metabolism upon high-fat-diet feeding. *FASEB J* 26: 656-667.

11. Tomita T, Hamazaki J, Hirayama S, McBurney MW, Yashiroda H, et al. (2015) Sirt1-deficiency causes defective protein quality control. *Sci Rep* 5: 12613.
12. Rensing L, Monnerjahn C (1996) Heat shock proteins and circadian rhythms. *Chronobiol Int* 13: 239-250.
13. Kakimura J, Kitamura Y, Takata K, Umeki M, Suzuki S, et al. (2002) Microglial activation and amyloid-beta clearance induced by exogenous heat-shock proteins. *FASEB J* 16: 601-603.
14. Wilhelmus MM, de Waal RM, Verbeek MM (2007) Heat shock proteins and amateur chaperones in amyloid-Beta accumulation and clearance in Alzheimer's disease. *Mol Neurobiol* 35: 203-216.
15. Martins IJ (2015) Diabetes and Cholesterol Dyshomeostasis Involve Abnormal a-Synuclein and Amyloid Beta Transport in Neurodegenerative Diseases. *Austin Alzheimers J Parkinsons Dis* 2: 1020-28.
16. Ariesandi W, Chang CF, Chen TE, Chen YR (2013) Temperature-dependent structural changes of Parkinson's alpha-synuclein reveal the role of pre-existing oligomers in alpha-synuclein fibrillization. *PLoS One* 8: e53487.
17. Ghavami M, Rezaei M, Ejtehadi R, Lotfi M, Sokargozar MA, et al. (2013) Physiological Temperature Has a Crucial Role in Amyloid Beta in the Absence and Presence of Hydrophobic and Hydrophilic Nanoparticles. *ACS Chem Neurosci* 4: 375-378.
18. Martins IJ (2015) Unhealthy Nutrigenomic Diets Accelerate NAFLD and Adiposity in Global communities. *J Mol Genet Med* 9: 1-11.
19. Tang BL (2016) Sirt1 and the Mitochondria. *Mol Cells* 39: 87-95.
20. Malhi H, Kaufman RJ (2011) Endoplasmic reticulum stress in liver disease. *J Hepatol* 54: 795-809.
21. Ashraf NU, Sheikh TA (2015) Endoplasmic reticulum stress and Oxidative stress in the pathogenesis of Non-alcoholic fatty liver disease. *Free Radic Res* 49: 1405-1418.
22. Bravo R, Gutierrez T, Paredes F, Gatica D, Rodriguez AE, et al. (2012) Endoplasmic reticulum: ER stress regulates mitochondrial bioenergetics. *Int J Biochem Cell Biol* 44: 16-20.
23. Amat R, Planavila A, Chen SL, Iglesias R, Giralt M, et al. (2009) SIRT1 controls the transcription of the peroxisome proliferator-activated receptor-gamma Co-activator-1alpha (PGC-1alpha) gene in skeletal muscle through the PGC-1alpha autoregulatory loop and interaction with MyoD. *J Biol Chem* 284: 21872-21880.
24. Rodgers JT, Lerin C, Gerhart-Hines Z, Puigserver P (2008) Metabolic adaptations through the PGC-1 alpha and SIRT1 pathways. *FEBS Lett* 582: 46-53.
25. Cantó C, Auwerx J (2009) PGC-1alpha, SIRT1 and AMPK, an energy sensing network that controls energy expenditure. *Curr Opin Lipidol* 20: 98-105.
26. Higashida K, Kim SH, Jung SR, Asaka M, Holloszy JO, et al. (2013) Effects of resveratrol and SIRT1 on PGC-1 $\alpha$  activity and mitochondrial biogenesis: a reevaluation. *PLoS Biol* 11: e1001603.
27. Cornu M, Oppliger W, Albert V, Robitaille AM, Trapani F, et al. (2014) Hepatic mTORC1 controls locomotor activity, body temperature, and lipid metabolism through FGF21. *Proc Natl Acad Sci U S A* 111: 11592-11599.
28. Ghosh HS, McBurney M, Robbins PD (2010) SIRT1 negatively regulates the mammalian target of rapamycin. *PLoS One* 5: e9199.
29. Huang J, Dibble CC, Matsuzaki M, Manning BD (2008) The TSC1-TSC2 complex is required for proper activation of mTOR complex 2. *Mol Cell Biol* 28: 4104-4115.
30. Martins IJ (2016) Anti-Aging Genes Improve Appetite Regulation and Reverse Cell Senescence and Apoptosis in Global Populations. *Advances in Aging Research* 5: 9-26.
31. Kuro-o M (2009) Klotho and aging. *Biochim Biophys Acta* 1790: 1049-1058.
32. Yan H, Jihong Y, Feng Z, Xiaomei X, Xiaohan Z, et al. (2014) Sirtuin 1-mediated inhibition of p66shc expression alleviates liver ischemia/reperfusion injury. *Crit Care Med* 42: e373-381.
33. Haga S, Morita N, Irani K, Fujiyoshi, M, et al (2010) p66Shc has a pivotal function in impaired liver regeneration in aged mice by a redox-dependent mechanism. *Special Issue – Hepatic and Pancreatic Systems. Lab Invest* 90: 1718-1726.
34. Carbajo-Pescador S, Mauriz JL, García-Palomo A, González-Gallego JI (2014) FoxO proteins: regulation and molecular targets in liver cancer. *Curr Med Chem* 21: 1231-1246.
35. Tikhonovich I, Cox J, Weinman SA (2013) Forkhead box class O transcription factors in liver function and disease. *J Gastroenterol Hepatol* 28 Suppl 1: 125-131.
36. Martins IJ (2016) The role of Clinical Proteomics, Lipidomics and Genomics in the diagnosis of Alzheimer's disease. *Proteome* 4: 1-19.
37. Martins IJ (2016) Magnesium Therapy Prevents Senescence with the Reversal of Diabetes and Alzheimer's Disease. *Health* 8: 694-710.
38. Zhu S, Wu Y, Ye X, Ma L, et al. (2016) FGF21 ameliorates nonalcoholic fatty liver disease by inducing autophagy. *Mol Cell Biochem* 420: 107-119.
39. Domouzoglou EM, Naka KK, Vlahos AP, Papafaklis MI, Michalis LK, et al. (2015) Fibroblast growth factors in cardiovascular disease: The emerging role of FGF21. *Am J Physiol Heart Circ Physiol* 309: H1029-1038.
40. Boon RA, Iekushi K, Lechner S, Seeger T, Fischer A, et al. (2013) MicroRNA-34a regulates cardiac ageing and function. *Nature* 495: 107-110.
41. Shang W, Yu X, Wang H, Chen T, Fang Y, et al. (2015) Fibroblast growth factor 21 enhances cholesterol efflux in THP-1 macrophage-derived foam cells. *Mol Med Rep* 11: 503-508.
42. Zhang X, Yeung DC, Karpisek M, Stejskal D, Zhou ZG, et al. (2008) Serum FGF21 levels are increased in obesity and are independently associated with the metabolic syndrome in humans. *Diabetes* 57: 1246-1253.
43. So WY, Leung PS (2016) Fibroblast Growth Factor 21 As an Emerging Therapeutic Target for Type 2 Diabetes Mellitus. *Med Res Rev* 36: 672-704.
44. Qiao L, Yoo Hs, Bosco C, Lee B, Feng GS, et al. (2014) Adiponectin reduces thermogenesis by inhibiting brown adipose tissue activation in mice. *Diabetologia* 57: 1027-1036.
45. Martins IJ (2016) Bacterial Lipopolysaccharides Change Membrane Fluidity with Relevance to Phospholipid and Amyloid Beta Dynamics in Alzheimer's Disease. *J Microb Biochem Technol.* 8: 322-324.
46. Martins IJ (2015) Diabetes and Organ Dysfunction in the Developing and Developed. *Global J of Med Res F Diseases* 15: 1-6.
47. Martins IJ (2013) Increased Risk for Obesity and Diabetes with Neurodegeneration in Developing Countries. *J Mol Genet Med* S1: 1-8.
48. Martins IJ (2015) Nutritional diets accelerate amyloid beta metabolism and prevent the induction of chronic diseases and Alzheimer's disease. *Photon ebooks.* 1-48.