

2018

## Organic carbon sequestration and storage in vegetated coastal habitats along the western coast of the Arabian Gulf

M Cusack

V Saderne

A Arias-Ortiz

Pere Masque

*Edith Cowan University, Australia*

P K. Krishnakumar

*See next page for additional authors*

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworkspost2013>



Part of the [Terrestrial and Aquatic Ecology Commons](#)

---

[10.1088/1748-9326/aac899](https://ro.ecu.edu.au/ecuworkspost2013/4499)

Cusack, M., Saderne, V., Arias-Ortiz, A., Masqué, P., Krishnakumar, P. K., Rabaoui, L., ... & Elyas, A. A. (2018). Organic carbon sequestration and storage in vegetated coastal habitats along the western coast of the Arabian Gulf.

*Environmental Research Letters*, 13(7). Available [here](#)

This Letter to the Editor is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/4499>

---

## Authors

M Cusack, V Saderne, A Arias-Ortiz, Pere Masque, P K. Krishnakumar, L Rabaoui, M A. Qusem, P Prihartato, R A. Loughland, A A. Elyas, and C M. Duarte

LETTER • OPEN ACCESS

# Organic carbon sequestration and storage in vegetated coastal habitats along the western coast of the Arabian Gulf

To cite this article: M Cusack *et al* 2018 *Environ. Res. Lett.* **13** 074007

View the [article online](#) for updates and enhancements.

## Related content

- [Geoengineering with seagrasses: is credit due where credit is given?](#)  
Sophia C Johannessen and Robie W Macdonald
- [A global map of mangrove forest soil carbon at 30 m spatial resolution](#)  
Jonathan Sanderman, Tomislav Hengl, Greg Fiske et al.
- [Soil greenhouse gas emissions reduce the contribution of mangrove plants to the atmospheric cooling effect](#)  
Guangcheng Chen, Bin Chen, Dan Yu et al.

## Environmental Research Letters



## LETTER

## OPEN ACCESS

RECEIVED  
8 April 2018

ACCEPTED FOR PUBLICATION  
30 May 2018

PUBLISHED  
27 June 2018

Original content from  
this work may be used  
under the terms of the  
[Creative Commons  
Attribution 3.0 licence](#).

Any further distribution  
of this work must  
maintain attribution to  
the author(s) and the  
title of the work, journal  
citation and DOI.



# Organic carbon sequestration and storage in vegetated coastal habitats along the western coast of the Arabian Gulf

M Cusack<sup>1,8</sup>, V Saderne<sup>1</sup>, A Arias-Ortiz<sup>2</sup>, P Masqué<sup>2,3,4</sup>, P K Krishnakumar<sup>5,6</sup>, L Rabaoui<sup>5,6</sup>,  
M A Qurban<sup>5,6</sup>, A M Qasem<sup>7</sup>, P Prihartato<sup>7</sup>, R A Loughland<sup>7</sup>, A A Elyas<sup>7</sup> and C M Duarte<sup>1</sup>

<sup>1</sup> King Abdullah University of Science and Technology, Red Sea Research Centre, Thuwal 23955-6900, Saudi Arabia

<sup>2</sup> Departament de Física & Institut de Ciència i Tecnologia Ambientals, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

<sup>3</sup> School of Science, Centre for Marine Ecosystems Research, Edith Cowan University, 270 Joondalup Drive, Joondalup WA 6027, Australia

<sup>4</sup> Oceans Institute & School of Physics, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

<sup>5</sup> Marine Studies section, Center for Environment and Water, Research Institute, King Fahd University of Petroleum and Minerals (KFUPM), P B No. 391, Dhahran 31261, Saudi Arabia

<sup>6</sup> Geosciences Department, The College of Petroleum Engineering & Geosciences, King Fahd University of Petroleum and Minerals (KFUPM), Dhahran 31261, Saudi Arabia

<sup>7</sup> Environmental Protection Department, Saudi Aramco, Dhahran, Saudi Arabia

<sup>8</sup> Author to whom any correspondence should be addressed.

E-mail: [michael.cusack@kaust.edu.sa](mailto:michael.cusack@kaust.edu.sa)

**Keywords:** blue carbon, carbon sink, marine coastal ecosystems, organic carbon burial

Supplementary material for this article is available [online](#)

## Abstract

Certain coastal ecosystems such as mangrove, saltmarsh and seagrass habitats have been identified as significant natural carbon sinks, through the sequestration and storage of carbon in their biomass and sediments, collectively known as 'blue carbon' ecosystems. These ecosystems can often thrive in extreme environments where terrestrial systems otherwise survive at the limit of their existence, such as in arid and desert regions of the globe. To further our understanding of the capability of blue carbon ecosystems to sequester and store carbon in such extreme climates, we measured carbon sediment stocks in 25 sites along the Western Arabian Gulf coast. While seagrass meadows and saltmarsh habitats were widely distributed along the coast, mangrove stands were much reduced as a result of anthropogenic pressures, with 90% of stands having been lost over the last century. Carbon stocks in 1 m deep surface sediments were similar across all three blue carbon habitats, with comparable stocks for saltmarsh ( $81 \pm 22 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$ ), seagrass ( $76 \pm 20 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$ ) and mangroves ( $76 \pm 23 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$ ). We recorded a 38% decrease in carbon stocks between mature established mangrove stands ( $91 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$ ) and recently planted mangroves ( $56 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$ ). Mangroves also had the lowest carbon stock per total area owing to their very limited spatial coverage along the coast. The largest stock per total area belonged to seagrass beds as a result of their large spatial coverage within the Gulf. We employed  $^{210}\text{Pb}$  dating to determine the sediment accretion rates in each ecosystem and found mangrove habitats to be the most efficient carbon sequesters over the past century, with the highest carbon burial rate of the three ecosystems ( $19 \text{ g C}_{\text{org}} \text{ m}^{-2} \text{ yr}^{-1}$ ), followed by seagrass ( $9 \text{ g C}_{\text{org}} \text{ m}^{-2} \text{ yr}^{-1}$ ) and saltmarshes ( $8 \text{ g C}_{\text{org}} \text{ m}^{-2} \text{ yr}^{-1}$ ). In this work, we describe a comprehensive comparison of sediment stocks in different blue carbon ecosystems within a single marine environment and across a large geographical area, and discuss our results in a global context for other blue carbon ecosystems in the dry tropics.

## 1. Introduction

Vegetated coastal habitats, including mangroves, seagrass meadows and saltmarshes, can sequester and store significant amounts of organic carbon (Duarte *et al* 2005, Fourqurean *et al* 2012). This has prompted increased interest in the potential role of these ecosystems in supporting climate change and adaptation policies, through so-called blue carbon strategies (Nellemann *et al* 2009, Mcleod *et al* 2011, Duarte *et al* 2013). However, studies on carbon sequestration in blue carbon habitats to-date have been disproportionately focused on sites in temperate or tropical climates (Chmura *et al* 2003, Donato *et al* 2011, Adame *et al* 2013, Fourqurean *et al* 2012, Brown *et al* 2016, Serrano *et al* 2016, Macreadie *et al* 2017). Similar studies in arid regions of the world are relatively scarce, where temperature extremes and low rainfall may affect net ecosystem productivity and primary productivity. Recent studies in arid regions such as those adjoining the Red Sea report some of the lowest short-term carbon sequestration rates ( $15 \text{ g C}_{\text{org}} \text{ m}^{-2} \text{ yr}^{-1}$ ) and sediment stocks ( $43 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$  in 1 m-thick soils) in mangrove forests (Almahasheer *et al* 2017), while (Schile *et al* 2017) also reported modest carbon stocks in blue carbon ecosystems ( $49\text{--}156 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$ ) within the Arabian Gulf coast of the United Arab Emirates. The distribution of organic carbon in mangrove forests of the Egyptian Red Sea coastline has been described by (Madkour *et al* 2014), who found moderate levels of organic carbon ( $190 \text{ Mg C}_{\text{org}} \text{ ha}^{-1}$ ) in mangrove sediments. However, studies and data on carbon stocks and sequestration rates for blue carbon ecosystems in the arid Middle East region remain critically low.

The Saudi Arabian coasts of the Arabian Gulf and the Red Sea comprise highly productive vegetated coastal ecosystems. Saltmarshes form vast ecosystems of halophytes known locally as *sabkha* (Qurban *et al* 2011). All of these ecosystems are at risk or in decline due to local pressures such as industrial activities and land reclamation for coastal development (Loughland *et al* 2012), which has resulted in the loss of around 90% of the mangrove cover (Loughland *et al* 2012) and unknown, but large, spans of *sabkha*, and possibly seagrass meadows. Almahasheer (2018) recently determined the spatial coverage of mangrove stands in the Arabian Gulf and found that mangrove coverage in Saudi Arabia, Qatar and Bahrain is stable, while stands in Iran have been in decline over recent years. The Middle East and the Gulf of Arabia basin is expected to be especially vulnerable to the negative impacts of climate change, and it is predicted that by the end of this century certain population centres in the region are likely to experience extreme temperatures, owing to the consequences of increasing concentrations of anthropogenic greenhouse gases (Pal and Eltahir 2016).

Here we present a comprehensive analysis on carbon stocks and  $\text{C}_{\text{org}}$  accumulation rates for mangrove, saltmarsh and seagrass habitats along the Western Gulf coast. This study will contribute important information from a climatically extreme region of the globe, which is underrepresented in the literature, thereby contributing to a more balanced account of global patterns in carbon sequestration and stocks in blue carbon habitats. In literature, the Arabian Gulf and the Persian Gulf are interchangeable, however for this study we refer to the study area as the Arabian Gulf.

## 2. Methods

### 2.1. Study region

#### 2.1.1. Blue carbon habitats along the western Arabian Gulf coastline

The Kingdom of Saudi Arabia has approximately 560 km of coastline in the Arabian Gulf, which covers an area of  $226\,000 \text{ km}^2$  and stretches almost 1000 km in length along its central axis (Sheppard *et al* 2010). The Gulf is a shallow, semi-enclosed sea with an average depth of 35–40 m, with the only opening to the Indian Ocean (Gulf of Oman) via the relatively narrow Strait of Hormuz (50 km). A number of factors such as limited circulation, extreme summer air temperature ( $>50^\circ\text{C}$ ) and shallow depth result in high evaporation, hypersalinity, and low water exchange with the open ocean (Kazemi *et al* 2012, Fard *et al* 2015). Due to the geographic location of the Gulf, surrounded by a largely arid region, there is little freshwater input from land to the Saudi coast and virtually no water borne sediment reaches the Gulf from the surrounding land mass. Thus, sediments of the Gulf originate mainly from carbonate deposits from sea water and deposited airborne sediment of terrestrial origin, which comprise one third of Gulf sediments Sugden (1963), and sediments delivered by the Tigris and Euphrates rivers in the north and transported to the Saudi coast. The Gulf suffers particularly high anthropogenic pressure owing to the large reserves of crude oil, which represent 57%–66% of the world's oil reserves (Vaezi *et al* 2015), and thus the presence of a large number of related industries and associated discharges. The industrial development of the region, along with high population growth driven by the large employment created by these industries, has led to significant coastal development, rapidly transforming previously pristine coastal environments through urbanisation, land reclamation and dredging (Lokier and Fiorini 2016). The consequent industrial and sewage effluents, wastewater discharges from desalination plants, shipping and oil tanker traffic and emissions from oil production have all contributed to negatively impact the Gulf environment (Sheppard *et al* 2010, Naser 2014, Loughland *et al* 2012).




Figure 1. Map of sampling sites: Zone 1) three seagrass sites at Uqair, Zone 2) three seagrass, three mangrove and three saltmarsh sites at Abu-Ali Island, Zone 3) three seagrass, four mangrove and three saltmarsh sites at Ras Tanura/Safwa and Zone 4) three seagrass sites at Safaniya.

Almahasheer (2018) estimated that around 165 km<sup>2</sup> fragmented mangrove stands are distributed unevenly around Arabian Gulf coastlines. Mangrove ecosystems along the Gulf coast of Saudi Arabia are limited in their distribution, existing mostly in sheltered areas along the coast (Saenger 2011). Mangrove development in the Gulf is restricted largely due to natural environmental pressure, such as aridity, high salinity and seasonal temperature fluctuations from cold to extreme heat. Mangroves on the west coast are represented by a single species, *Avicennia marina*, which is adapted to withstand the local extreme environmental conditions. On the Iranian coast, a mixture of *Avicennia marina* and *Rhizophora mucronata* can be found (Zahed *et al* 2010). Anthropogenic pressures, such as land reclamation and coastal development, have led to the loss of an estimated 90% of the original mangrove ecosystems along the Saudi coast, with much of the remaining mangrove area likely to be lost within a decade at current lost rates (Loughland *et al* 2012), except for areas under protection.

Supratidal flats occupy 30%–40% of the Saudi coastline (Basson *et al* 1977), and mainly consist of mud or very fine sand in sheltered areas where wave energy is limited. Saltmarshes, or sabkhas, of the Saudi Gulf coastline have been surveyed by Al-Abdulkader and Loughland (2012), and were found to be dominated by salt tolerant or halophytic perennial grasses such as *Aeluropus lagopoides* and *Bienertia cycloptera*. Seagrass beds, including three species: *Halodule uninervis* (most widely distributed), *Halophila stipulacea* and *Halophila ovalis* (less common) (Price and Coles 1992), occur at 46% of the coastal and 30% of off-

shore sites of the Gulf coast (Erftemeijer and Shuaib 2012). Seagrass meadows in the region are under local environmental pressure due to dredging, land reclamation and associated changes in hydrology, together with pressures derived from climate change and marine pollution.

This study focuses on vegetated coastal habitats along the Gulf coast of Saudi Arabia, spanning 400 km of the coastline, extending from 25° 43' N (Uqair) on the southwest coast to 27° 58' N (Safaniya) on the northwest coast (table 1, figure 1). Sediment core samples were taken in replicate at 25 locations: 12 seagrass habitats, seven mangrove and six saltmarsh habitats (table 1). The mangroves sampled included both recently planted mangroves in Abu-Ali, from a successful replanting project undertaken in the early 1990s, and natural mangrove forests in the protected areas of Ecopark and Safwa (table 1).

#### 2.1.2. Sediment sampling and analysis

PVC cores of 140 cm length and 7 cm internal diameter were employed to collect undisturbed sediment core samples. Cores were manually hammered into the sediment until resistance prevented further percussion (table 1). Compression of sediments during coring was corrected for by distributing the spatial discordances proportionally between the expected and the observed soil column layers and was accounted for in the calculations of bulk density,  $C_{org}$  stocks and accretion rates. The sediment was cut in 1 cm thick segments. Each slice was then dried at 60 °C until constant weight in order to determine bulk density (DBD: g cm<sup>-3</sup>). The dried sediment slices were then ground by pestle and

**Table 1.** Summary of sediment parameters per core, mass accumulation rate (MAR) and sediment accretion rate (SAR)  $\pm$  uncertainty, estimated using  $^{210}\text{Pb}$  dating over the past decades to century.  $C_{\text{org}}$  (%), dry bulk density (DBD) are means  $\pm$  SE, stocks extrapolated to surface 1 m depth ( $\pm$  SE).

Region	Habitat	Core ID	Latitude ( $^{\circ}\text{N}$ )	Longitude ( $^{\circ}\text{E}$ )	Sediment depth (cm)	Mean $C_{\text{org}}$ content (%)	DBD ( $\text{g cm}^{-3}$ )	Stock ( $\text{Mg } C_{\text{org}} \text{ ha}^{-1}$ )	MAR ( $\text{g cm}^{-2} \text{ yr}^{-1}$ )	SAR ( $\text{mm yr}^{-1}$ )	$C_{\text{org}}$ burial rate ( $\text{g } C_{\text{org}} \text{ m}^{-2} \text{ yr}^{-1}$ )
Safaniya	Seagrass	SSGC1	27.97364	48.77404	109	$0.54 \pm 0.03$	$1.3 \pm 0.03$	$74 \pm 4$	$0.34 \pm 0.04$	$2.4 \pm 0.3$	$19 \pm 2$
Safaniya	Seagrass	SSGC2	27.97679	48.77877	90	$0.56 \pm 0.01$	$1.3 \pm 0.02$	$73 \pm 2$	n/a	n/a	n/a
Safaniya	Seagrass	SSGC3	27.98142	48.78462	110	$0.50 \pm 0.04$	$1.1 \pm 0.02$	$62 \pm 4$	$0.23 \pm 0.03$	$1.7 \pm 0.2$	$9.0 \pm 1.4$
Abu-Ali	Mangrove	AAMGC1	27.28603	49.56685	102	$0.36 \pm 0.02$	$1.4 \pm 0.02$	$55 \pm 3$	$0.25 \pm 0.02$	$1.59 \pm 0.10$	$8.7 \pm 0.8$
Abu-Ali	Mangrove	AAMGC2	27.28384	49.56403	103	$0.39 \pm 0.02$	$1.3 \pm 0.03$	$51 \pm 3$	$0.176 \pm 0.013$	$1.24 \pm 0.09$	$6.2 \pm 0.6$
Abu-Ali	Mangrove	AAMGC3	27.28283	49.56523	103	$0.46 \pm 0.05$	$1.4 \pm 0.02$	$63 \pm 2$	$0.37 \pm 0.02$	$2.6 \pm 0.2$	$23 \pm 3$
Abu-Ali	Saltmarsh	AASMC1	27.28777	49.56953	103	$0.53 \pm 0.07$	$1.4 \pm 0.04$	$77 \pm 4$	$0.178 \pm 0.008$	$1.28 \pm 0.06$	$9.4 \pm 0.8$
Abu-Ali	Saltmarsh	AASMC2	27.31332	49.56401	72	$0.73 \pm 0.07$	$1.0 \pm 0.05$	$75 \pm 5$	$0.102 \pm 0.007$	$0.96 \pm 0.07$	$6.0 \pm 0.6$
Abu-Ali	Saltmarsh	AASMC3	27.29719	49.48471	67	$0.54 \pm 0.08$	$1.5 \pm 0.06$	$68 \pm 4$	$0.123 \pm 0.012$	$0.87 \pm 0.09$	$5.1 \pm 0.6$
Abu-Ali	Seagrass	AASGC1	27.26978	49.54546	35	$0.76 \pm 0.01$	$1.3 \pm 0.01$	$95 \pm 1.4$	$0.21 \pm 0.06$	$1.6 \pm 0.5$	$16 \pm 4$
Abu-Ali	Seagrass	AASGC2	27.25976	49.57006	118	$0.98 \pm 0.12$	$0.9 \pm 0.02$	$83 \pm 4$	$0.09 \pm 0.02$	$0.9 \pm 0.2$	$4.9 \pm 1.2$
Abu-Ali	Seagrass	AASGC3	27.27886	49.54524	116	$1.12 \pm 0.04$	$1.2 \pm 0.02$	$127 \pm 1.3$	n/a	n/a	n/a
Ecopark	Mangrove	RTMGC1	26.71674	50.02111	54	$0.82 \pm 0.07$	$1.2 \pm 0.02$	$99 \pm 4$	$0.50 \pm 0.05$	$3.60 \pm 0.40$	$38 \pm 4$
Ecopark	Mangrove	SAMGC1	26.65781	50.00534	28	$0.87 \pm 0.07$	$1.3 \pm 0.03$	$113 \pm 12$	$0.26 \pm 0.10$	$2.00 \pm 0.70$	$22 \pm 9$
Ecopark	Saltmarsh	RTSMC1	26.65761	50.01003	40	$0.94 \pm 0.03$	$1.4 \pm 0.08$	$53 \pm 3$	$0.082 \pm 0.010$	$0.60 \pm 0.07$	$3.1 \pm 0.4$
Ecopark	Saltmarsh	RTSMC2	26.71998	50.02104	98	$0.40 \pm 0.11$	$1.2 \pm 0.02$	$104 \pm 7$	$0.13 \pm 0.03$	$1.2 \pm 0.3$	$12 \pm 3$
Ecopark	Saltmarsh	SASMC1	26.73849	49.99526	117	$0.81 \pm 0.08$	$1.4 \pm 0.02$	$113 \pm 7$	$0.119 \pm 0.014$	$1.04 \pm 0.11$	$11.2 \pm 1.6$
Ecopark	Seagrass	RTSGC3	26.72047	49.99885	106	$0.74 \pm 0.04$	$1.2 \pm 0.02$	$86 \pm 2$	$0.10 \pm 0.05$	$0.8 \pm 0.4$	$7 \pm 3$
Ecopark	Seagrass	RTSGC2	26.70883	50.01225	61	$0.60 \pm 0.05$	$1.2 \pm 0.02$	$71 \pm 6$	$0.08 \pm 0.02$	$0.60 \pm 0.2$	$3.8 \pm 0.9$
Ecopark	Seagrass	RTSGC1	26.68064	50.01184	104	$0.68 \pm 0.04$	$1.1 \pm 0.02$	$75 \pm 3$	n/a	n/a	n/a
Safwa	Mangrove	SAMGC3	26.64246	50.01462	43	$0.54 \pm 0.02$	$1.5 \pm 0.02$	$82 \pm 1.1$	$0.28 \pm 0.09$	$1.5 \pm 0.5$	$12 \pm 4$
Safwa	Mangrove	SAMGC2	26.63699	50.01067	83	$0.48 \pm 0.01$	$1.5 \pm 0.02$	$71 \pm 2$	$0.44 \pm 0.06$	$2.7 \pm 0.4$	$20 \pm 3$
Uqair	Seagrass	UQSGC1	25.72452	50.23141	93	$0.49 \pm 0.01$	$1.1 \pm 0.01$	$55 \pm 1.3$	n/a	n/a	n/a
Uqair	Seagrass	UQSGC2	25.72891	50.22990	87	$0.56 \pm 0.03$	$1.0 \pm 0.01$	$58 \pm 4$	$0.16 \pm 0.06$	$1.3 \pm 0.5$	$6 \pm 2$
Uqair	Seagrass	UQSGC3	25.73403	50.22677	76	$0.59 \pm 0.03$	$0.9 \pm 0.02$	$55 \pm 3$	$0.16 \pm 0.06$	$1.3 \pm 0.5$	$6 \pm 2$



mortar. Every fifth slice down to 20 cm depth was analysed for organic carbon (% OC) on a CHN analyser (Thermo Scientific CHNS element analyser). A 1 g sub-sample of ground sample was acidified with 4% HCL to remove inorganic carbon, centrifuged (3400 revolutions per minute, for 5 min), and the supernatant with acid residues was carefully removed by pipette, avoiding resuspension. The sample was then washed with Milli-Q water, centrifuged and the supernatant removed. The residual samples were then re-dried and encapsulated for organic carbon analyses.

Sediment accumulation rates for the last decades-century were determined using the  $^{210}\text{Pb}$  method. The concentrations of  $^{210}\text{Pb}$  in the upper 19 cm were determined in the fraction  $<125\ \mu\text{m}$  at the Edith Cowen University (Australia) by alpha spectrometry through the measurement of its granddaughter  $^{210}\text{Po}$ , assuming radioactive equilibrium between both radionuclides (Sanchez-Cabeza *et al* 1998). The concentrations of excess  $^{210}\text{Pb}$  used to obtain the age models were determined as the difference between total  $^{210}\text{Pb}$  and  $^{226}\text{Ra}$  (supported  $^{210}\text{Pb}$ ). Concentrations of  $^{226}\text{Ra}$  were determined for selected samples along each core by low-background liquid scintillation counting method (Wallac 1220 Quantulus) adapted from (Masqué *et al* 2002). These concentrations were found to be comparable with the concentrations of total  $^{210}\text{Pb}$  at depth below the excess  $^{210}\text{Pb}$  horizons. Analyses of reagent blanks, replicates and a reference material (IAEA-315, marine sediments) were carried out for both  $^{210}\text{Pb}$  and  $^{226}\text{Ra}$  to assess for any contamination and to ensure reproducibility of the results. Average sediment accumulation rates (expressed in  $\text{g DW cm}^{-2}\text{ yr}^{-1}$ ) for the last decades/century were estimated using the Constant Flux: Constant Sedimentation model, (Krishnaswami *et al* 1971) and accretion rates ( $\text{mm yr}^{-1}$ ) were calculated using the dry bulk density (DBD) data of each core (table 1).

$\text{C}_{\text{org}}$  stocks per unit area were calculated by summing  $\text{C}_{\text{org}}$  stocks in each segment interval ( $\text{g C}_{\text{org}}\text{ cm}^{-2}$ ) down to 20 cm; segments for which we did not measure  $\text{C}_{\text{org}}$  data were interpolated.  $\text{C}_{\text{org}}$  data deeper than 20 cm were extrapolated to the equivalent of 1 m depth to bring values in line with globally reported stocks. Average short-term  $\text{C}_{\text{org}}$  burial rates were estimated by multiplying sediment accumulation rates ( $\text{g cm}^{-2}\text{ yr}^{-1}$ ) by the fraction of C accumulated to 100 year depth determined by  $^{210}\text{Pb}$  dating.

### 3. Results and discussion

The length of sediment penetrated and collected ranged from 28–118 cm, and were, in general, deeper in seagrass sediments (table 1). Mean bulk density of sediments was similar for the three habitats, around  $1.4\text{ g cm}^{-3}$ , with no evident geographical difference or significant changes with depth. The highest bulk densities ( $>1.5\text{ g cm}^{-3}$ ) were found in saltmarsh cores,


which were typically comprised of coarser sandy sediments. Organic carbon content ( $\text{C}_{\text{org}}\%$  DW) ranged from 0.3% to a maximum of 1.4%, with a mean  $\text{C}_{\text{org}}$  of  $0.6 \pm 0.2\%$  and median of 0.6%.

Mean  $\text{C}_{\text{org}}$  stocks ( $\pm\text{SE}$ ), calculated in the upper meter of sediment were similar across habitats, at  $76 \pm 23$ ,  $81 \pm 23$  and  $76 \pm 20\text{ Mg C}_{\text{org}}\text{ ha}^{-1}$  for mangrove, saltmarsh and seagrass respectively (table 2). Carbon stocks in the upper meter ranged two fold within habitats, from 51–113  $\text{Mg C}_{\text{org}}\text{ ha}^{-1}$  in the mangrove sediments, 53–113  $\text{Mg C}_{\text{org}}\text{ ha}^{-1}$  in saltmarshes and 55–127  $\text{Mg C}_{\text{org}}\text{ ha}^{-1}$  in seagrass sediments (figure 2; table 1). The studied mangrove sites included both recently planted (Abu-Ali; early 1990s) and natural mangrove stands (Ecopark and Safwa).  $\text{C}_{\text{org}}$  stocks were considerably lower for the planted ( $56\text{ Mg C}_{\text{org}}\text{ ha}^{-1}$ ) relative to mature mangroves ( $91\text{ Mg C}_{\text{org}}\text{ ha}^{-1}$ ). Taking the area coverage of remaining mangrove stands along the Saudi coast (256 ha), we estimate a total of 20 Gg  $\text{C}_{\text{org}}$  stored in the top 1 m of sediment of mangrove ecosystems in the Arabian Gulf coast of Saudi Arabia, and 1254 Gg  $\text{C}_{\text{org}}$  for the entire Gulf, based on mangrove coverage ( $165.5\text{ km}^2$ ) estimated by Almahasheer (2018). Unfortunately we do not know when the original mangrove stands along the coast were destroyed, as human activities have impacted mangrove plantations on the Gulf coast for centuries. However, if we consider that the remaining mangrove stands represent only 10% of the original coverage, then we can estimate that, assuming all sediment  $\text{C}_{\text{org}}$  was mineralised, a maximum of around 660  $\text{CO}_2\text{eq}$  may have been potentially emitted as a result of their destruction. Although the exact extent of the coverage of seagrass and saltmarshes is unknown, around  $7000\text{ km}^2$  of seagrass meadows in the Arabian Gulf have been mapped (Erftemeijer and Shuail 2012). Based on this area coverage, we estimate that seagrass meadows of the Arabian Gulf store around 150 Tg of  $\text{C}_{\text{org}}$  in the top meter of sediment.

The  $^{210}\text{Pb}$  concentration profiles of 21 out of 25 cores showed mixing was a significant issue in the seagrass corers, all having a relatively thick mixed layer in the upper cms. Mangrove corers in Abu-Ali showed evidence of mixing in the top 1–4 cm for AAMGC1 and AAMGC3, and 1–7 cm for AAMGC2 (figure S1 available at [stacks.iop.org/ERL/13/074007/mmedia](https://stacks.iop.org/ERL/13/074007/mmedia)) and this is likely a result of the recent replanting in the 1990s which would have considerably disturbed the sediments. Mixing was not an issue in saltmarsh sediments, probably owing to the reduced influence of bioturbation.

Mass and sediment accumulation rate (MAR, SAR) varied substantially between ecosystems (figure 2; table 2), with mangroves having the highest average MAR and SAR of  $0.34 \pm 0.13\text{ g cm}^{-2}\text{ yr}^{-1}$  and  $0.22 \pm 0.17\text{ mm yr}^{-1}$ , followed by seagrass at  $0.17 \pm 0.08\text{ g cm}^{-2}\text{ yr}^{-1}$  and  $0.13 \pm 0.12\text{ mm yr}^{-1}$  and finally saltmarsh with  $0.12 \pm 0.03\text{ g cm}^{-2}\text{ yr}^{-1}$  and





**Table 2.** Mean  $C_{org}$  stocks in the top 1 m sediment, MAR and SAR, and modern (last century) burial rates for mangrove, saltmarsh and seagrass habitats in the Arabian Gulf.

Ecosystem	No. of samples	$C_{org}$ stock ( $Mg C_{org} ha^{-1}$ )	MAR ( $g cm^{-2} yr^{-1}$ )	SAR ( $cm yr^{-1}$ )	Modern burial rate ( $g C_{org} m^{-2} yr^{-1}$ )
Mangrove	7	$76 \pm 23$	$0.34 \pm 0.13$	$0.21 \pm 0.09$	$19 \pm 11$
Saltmarsh	6	$81 \pm 20$	$0.12 \pm 0.03$	$0.10 \pm 0.01$	$8 \pm 4$
Seagrass	12	$79 \pm 2$	$0.17 \pm 0.08$	$0.13 \pm 0.05$	$9 \pm 6$

$0.10 \pm 0.06$  mm  $yr^{-1}$ . Variability in local  $C_{org}$  burial rates during the last decades largely reflected the sediment accretion rates across all sampling sites, with similar values in saltmarshes and seagrasses ( $8 \pm 4$  and  $9 \pm 6$   $g C_{org} m^{-2} yr^{-1}$ ), and the highest rate of burial found in mangrove habitats ( $19 \pm 11$   $g C_{org} m^{-2} yr^{-1}$ ; table 2). However, it should be noted that it was not possible to determine the MAR, and thus the burial rate of  $C_{org}$ , for four of the seagrass sediments due to surface mixing of the sediment, which precluded reliable dating with  $^{210}Pb$ . Moreover, values for MAR and SAR among individual seagrass sites exhibited larger variability relative to the other sites, ranging from  $0.08 \pm 0.02$  to  $0.34 \pm 0.04$   $g cm^{-2} yr^{-1}$  and  $0.06 \pm 0.2$  to  $0.24 \pm 0.3$   $cm yr^{-1}$ , respectively. We estimate that on an annual basis mangrove stands of the Arabian Gulf coastline currently sequester  $3.1 \pm 1.8$   $Gg C_{org} yr^{-1}$ . If the mangroves that have been lost over the centuries

were restored, mangroves could annually sequester as much as 28  $Gg C_{org} yr^{-1}$ , increasing the stored  $C_{org}$  stock to 180  $Gg$ . Seagrasses of the Gulf, based on the coverage of 7000  $km^2$  estimated by Erftemeijer and Shuail (2012), sequester  $66 \pm 40$   $Gg C_{org} yr^{-1}$ . Given the annual sequestration rates of mangrove and seagrass, we estimate that the top metre of sediments in mangroves and seagrasses have been accumulating for 400 and 800 years respectively.

The widespread sampling undertaken in this study covering a large geographical area demonstrates that blue carbon stocks for this region are considerably less variable, both spatially and between ecosystems, compared to climatically similar parts of the globe. For example, (Schile *et al* 2017) reported a large variation of  $C_{org}$  sediment stocks in United Arab Emirates ranging from 2–515  $Mg C_{org} ha^{-1}$ . The average  $C_{org}$  stocks in the upper 1 m of sediments in the studied

blue carbon ecosystems were substantially lower than global mean local  $C_{org}$  stocks reported by (Duarte *et al* 2013) for saltmarshes ( $81 \pm 23$  in the Arabian Gulf vs.  $162 \text{ Mg } C_{org} \text{ ha}^{-1}$  as a global mean), mangroves ( $76 \pm 23$  in the Arabian Gulf vs.  $255 \text{ Mg } C_{org} \text{ ha}^{-1}$  as a global mean) and seagrasses ( $76 \pm 20$  in the Arabian Gulf vs.  $140 \text{ Mg } C_{org} \text{ ha}^{-1}$  as a global mean). However, our sediment stocks of  $C_{org}$  were comparable to those reported by (Schile *et al* 2017) for the climatically and geographically similar United Arab Emirates, who reported stocks of  $80 \text{ Mg } C_{org} \text{ ha}^{-1}$  in the upper meter of saltmarsh sediments. Furthermore, (Schile *et al* 2017) described a significant difference in carbon stocks between planted mangrove and mature mangrove stands, whereby planted mangrove sediments in UAE ( $102 \text{ Mg } C_{org} \text{ ha}^{-1}$ ) contained around 35% less  $C_{org}$  compared to mature mangrove stands ( $156 \text{ Mg } C_{org} \text{ ha}^{-1}$ ). In our study,  $C_{org}$  stocks were 38% lower for the planted mangroves of Abu-Ali ( $56 \text{ Mg } C_{org} \text{ ha}^{-1}$ ) relative to mature mangrove  $C_{org}$  stocks ( $91 \text{ Mg } C_{org} \text{ ha}^{-1}$ ). Lovelock *et al* (2017) recently stated that restored mangrove forests take around 20–25 years to match the  $C_{org}$  sediment stocks of undisturbed mangroves. 25 years since plantation, mangrove sediments at Abu-Ali contain around 62% less  $C_{org}$  compared to mature mangrove stands 100 km to the south.

The low  $C_{org}$  stocks for mangrove sediments in this region may be a result of the harsh environment, where *Avicennia marina* exists at its latitudinal limits, and the absence of rivers supplying sediments to these mangrove stands. Mangrove heights here rarely exceed 5 m, typically reaching between 1–1.5 m in height, reflecting the limiting environment characterised by extreme temperatures (both low and high), low rainfall, high salinity and limited nutrient availability. Furthermore, the Gulf War oil spill in 1991 adversely affected the recently planted mangroves on the shorelines of Abu Ali (Al-Abdulkader and Loughland 2012). Atwood *et al* (2017) recently described global patterns of soil carbon stocks by compiling data and comparing  $C_{org}$  stocks in mangrove sediments across the globe. However, measurements from arid regions included in that study were limited to the Red Sea region and North West Mexico, with no studies from the Arabian Gulf included. As Almahasheer (2018) recently noted, the Arabian Gulf has significant spatial coverage of mangrove stands ( $165.5 \text{ km}^2$ ), and this would contribute considerably to the global estimates of sequestered  $C_{org}$ . Combining our data for mangrove  $C_{org}$  stocks of the Saudi Arabian Gulf coast ( $91 \text{ Mg } C_{org} \text{ ha}^{-1}$  (mature mangrove stands only)) with that of Baja California reported by (Ezcurra *et al* 2016,  $232 \text{ Mg } C_{org} \text{ ha}^{-1}$ ), the Red Sea coast of Egypt (Atwood *et al* 2017,  $185 \text{ Mg } C_{org} \text{ ha}^{-1}$ ), the Red Sea coast of Saudi Arabia (Almahasheer 2018,  $43 \text{ Mg } C_{org} \text{ ha}^{-1}$ ) and the Gulf coast of the UAE (Schile *et al* 2017,  $156 \text{ Mg } C_{org} \text{ ha}^{-1}$

(mature mangrove stands only)), we calculate that mangrove sediments in tropical arid regions store on average  $139 \text{ Mg } C_{org} \text{ ha}^{-1}$ . If we consider that mangrove coverage for the Red Sea is  $135 \text{ km}^2$  (Almahasheer *et al* 2017),  $165.5 \text{ km}^2$  in the Arabian Gulf (Almahasheer 2018) and  $1873 \text{ km}^2$  in Baja California (Ezcurra *et al* 2016), we can estimate that mangrove sediments to 1 m depth in tropical arid climates store  $30 \text{ Tg } C_{org}$ . Previous estimates on global  $C_{org}$  stocks such as those of (Donato *et al* 2011 and Duarte *et al* 2013) were calculated based on measurements taken in tropical Indo-Pacific climates, which did not include data from the tropical arid zones. This has led to a probable over-estimation of global mangrove  $C_{org}$  stocks which have been biased by an emphasis on high carbon density mangrove systems, overlooking regions of less carbon dense mangrove sediments. Atwood *et al* (2017) calculated an average global carbon stock in mangrove sediments across a range of latitudes and climates of  $283 \pm 193 \text{ Mg } C \text{ ha}^{-1}$ , which we can revise slightly downwards with the inclusion of our study to  $281 \pm 194 \text{ Mg } C \text{ ha}^{-1}$ .

Published data on  $C_{org}$  content of seagrass sediments for comparison are currently lacking relative to mangrove sites. Fourqurean *et al* (2012) compiled  $C_{org}$  data from seagrass sediments across the globe and estimated a global mean stock in the upper meter of  $140 \text{ Mg } C_{org} \text{ ha}^{-1}$ , almost double the stocks we measured in seagrass meadows in the Arabian Gulf. This global average only included one study from an area with similar geographical characteristics to our study sites (Abu Dhabi), where (Campbell *et al* 2015) estimated a carbon stock of  $49 \text{ Mg } C_{org} \text{ ha}^{-1}$ , which is more than one third less than our estimates for the Saudi Arabian coast ( $76 \text{ Mg } C_{org} \text{ ha}^{-1}$ ). (Arias-Ortiz *et al* 2018) measured  $128 \pm 7 \text{ Mg } C_{org} \text{ ha}^{-1}$  in Western Australian seagrass sediments, one of the largest assemblages of seagrasses worldwide which exists in a similarly arid environment. Contrary to most published studies that show that seagrass sediments typically contain significantly less  $C_{org}$  compared to saltmarsh and mangrove sediments (Duarte *et al* 2013, Murray *et al* 2011, Siikamäki *et al* 2013, Ewers Lewis *et al* 2018), we found that sediments in Arabian Gulf seagrass meadows contain comparable stocks of  $C_{org}$  to mangrove and saltmarsh sediments.

The capacity of seagrass sediments to store carbon is believed to be influenced by local environmental conditions and landscape configuration, where exchanges of carbon may occur between habitats (Hyndes *et al* 2014). For example, (Walton *et al* 2014) demonstrated the tight coupling between arid mangroves and seagrass beds; they noted that the contribution of mangroves to seagrass sediment  $C_{org}$  stocks is often negligible in arid regions, yet there is  $C_{org}$  transfer from seagrass to mangrove, meaning seagrass beds are supporting mangroves stands. A recent study by Gullström *et al* (2018) observed that seagrass meadows covering a

larger area with clearly defined limits tended to contain more  $C_{org}$  per area relative to a patchy seagrass landscape adjacent to mangrove stands.  $C_{org}$  stocks in each study area in this work exhibited low spatial variability i.e. within seagrass meadows of Safaniya, Uqair and Ecopark, although they did vary within the Abu-Ali region (83–127 Mg  $C_{org}$  ha<sup>-1</sup>). The low areal coverage of mangrove stands along the Gulf coast may also limit the capacity of Gulf seagrass sediments to store carbon, as pointed out in a recent study by (Chen *et al* 2017) who suggested that proximity to mangroves can enhance seagrass  $C_{org}$  stocks. These findings highlight the large variability in stocks which can occur within relatively small areas, which has important implications when we attempt to assess the relative risk of CO<sub>2</sub> emissions if these habitats are destroyed (Lovelock *et al* 2017).

The variability of  $C_{org}$  stocks and %  $C_{org}$ , both spatially and within ecosystems, may be partially explained by their capability to successfully vertically accrete and bury carbon in sediments. Local  $C_{org}$  burial rates for blue carbon ecosystems in this work were around ten times lower than the global averages reported by (Duarte *et al* 2013) for mangroves, saltmarsh and seagrasses (163, 218 and 138 g  $C_{org}$  m<sup>-2</sup> yr<sup>-1</sup> respectively), and further differ in that mangrove ecosystems in our study sequester almost twice as much carbon as saltmarshes, in contrast to the global norm (McLeod *et al* 2011). The SAR derived in this work reveal that accretion rates in the study area fall right at the lower end of the global ranges for mangrove (0.3–21 mm yr<sup>-1</sup>), saltmarsh (0.4–61 mm yr<sup>-1</sup>) and seagrass meadows (0.6–6 mm yr<sup>-1</sup>) (Duarte *et al* 2013). Breithaupt *et al* (2012) reported a median SAR in mangrove stands of 2.8 mm yr<sup>-1</sup> based on 19 studies, which is comparable with our data (2.2 ± 0.3 mm yr<sup>-1</sup>), even though the study areas reported in (Breithaupt *et al* 2012) tended to receive riverine input or were at least in regions of high precipitation, unlike our study region. Other measurements of burial rates of  $C_{org}$  in arid regions are few. Almahasheer *et al* (2017) reported similar short-term (centennial) burial rates in mangroves to this study of 15 g  $C_{org}$  m<sup>-2</sup> yr<sup>-1</sup>. Ezcurra *et al* (2016) reported a large range of  $C_{org}$  burial rates in Baja California of 10–260 g  $C_{org}$  m<sup>-2</sup> yr<sup>-1</sup>. The low SAR and  $C_{org}$  burial rates observed across all ecosystems we sampled are most likely related to the restricted terrigenous inputs limited to aeolian dust deposition, as there are no permanent fluvial inputs anywhere along this stretch of coast. The higher mean SAR value for mangroves may be related to the efficiency of this habitat to trap both marine sediments and wind-borne dust amongst its aboveground biomass, in contrast to saltmarshes which are more exposed to wind removal processes, or removal by underwater currents such as in seagrass meadows. Furthermore, the mangrove stands in this study, except for Abu Ali, were surrounded by reclaimed land and urban devel-

opment/infrastructure, which may increase natural sedimentation rates and deposition.

Short-term  $C_{org}$  accumulation rates, as determined by <sup>210</sup>Pb dating, have been shown to be significantly higher than long-term accumulation rates, possibly due to the slow but continuous remineralisation of carbon in sediments over time (Choi and Wang 2004). Maher *et al* (2017) determined that carbon below an average depth of 40 cm (equivalent to 100 years of sediment accumulation) in mangrove sediments was susceptible to mineralisation and subsequent export. Baskaran *et al* (2017) further hypothesised that carbon mineralisation takes place over the sedimentary column resulting in significant losses of  $C_{org}$  in the long-term at deeper depths. Thus, our results on short-term  $C_{org}$  burial may overestimate the long-term burial capability of these ecosystems to sequester carbon. However, we found no consistent trend of decreasing  $C_{org}$  concentrations with depth, suggesting that mineralisation, if occurring, was masked by the variability in  $C_{org}$  over the past century.

The estimates of carbon stocks presented here highlight the importance of the protection of these threatened marine habitats. Whereas the carbon stocks currently held in mangrove sediments are modest, this only represents 10% of the historical coverage, as 90% of the mangrove cover has been lost. Hence, there is ample scope for mangrove restoration, as exemplified by replanting efforts along the Saudi coast of the Red Sea. Moreover, the comparison of  $C_{org}$  stocks in restored vs. undisturbed mangrove sediments in our survey shows that  $C_{org}$  stocks are rebuilt rapidly following restoration, as mangroves planted around 25 years ago have already developed carbon stocks that are only one third lower than nearby undisturbed mangroves forests (figure 2(a)).

## 4. Conclusions

This study offers a major advance in our understanding of Arabian Gulf coastal ecosystems and their capacity to sequester carbon. Furthermore, this work contributes to balance the global database on blue carbon habitat  $C_{org}$  storage by including vegetated coastal habitats in arid regions, underrepresented in current global assessments, such as that recently published by (Atwood *et al* 2017). Our results show that mangroves, seagrass and saltmarshes along the western coast of the Arabian Gulf are acting as active CO<sub>2</sub> sinks, burying substantial amounts of carbon in their sediments on an annual basis, and contain moderate stocks of sequestered carbon, comparable per unit area among habitats, but largest at the regional scale in seagrass meadows. The destruction or disturbance of these habitats could result in the remineralisation of a fraction of the stored carbon, giving rise to significant CO<sub>2</sub> emissions, while also making the ecosystems incapable

of acting as carbon sinks in the future. Restoration of these habitats, particularly mangroves that have suffered the greatest losses, offer opportunities to contribute to mitigate climate change through CO<sub>2</sub> sequestration, while also helping to protect the shorelines from sea level rise. The Middle East is especially vulnerable to future projected climate change, being an already extremely arid environment with very limited rainfall, few permanent fresh water sources and an increasing population. However, the results presented in this work indicate that despite the extreme environment, coastal blue carbon ecosystems store low to moderate levels of carbon and are successfully sequestering carbon on an annual basis. This study demonstrably shows for the first time the value of mangrove, saltmarsh and seagrass habitats as carbon sinks on the western coastline of the Arabian Gulf.

## Acknowledgments

This work was supported by a grant from Saudi Aramco. Further funding was provided to PM by the Generalitat de Catalunya (MERS, 2017SGR 1588) and an Australian Research Council LIEF Project (LE170100219). This work is contributing to the ICTA 'Unit of Excellence' (MinECo, MDM2015-0552). The authors thank the staff at the Marine Studies department of the Centre for Environment and Water of King Fahd University of Petroleum and Minerals, for their support and expertise, especially R Lindo, R Magalles, P Bacquiran and M Lopez. We also thank I Schulz, N Gerdali, D Prabowo and R Diaz at KAUST and G Salgado at ECU for their technical support.

## ORCID iDs

M Cusack  <https://orcid.org/0000-0002-9804-2641>

## References

- Al-Abdulkader K A and Loughland R A 2012 *Marine Atlas: Western Arabian Gulf* (Saudi Aramco Environmental Protection)
- Adame M F, Kauffman J B, Medina I, Gamboa J N, Torres O, Caamal J P, Reza M and Herrera-Silveira J A 2013 Carbon stocks of tropical coastal wetlands within the karstic landscape of the Mexican Caribbean *PLoS ONE* **8** e56569
- Almahasheer H 2018 Spatial coverage of mangrove communities in the Arabian Gulf *Environ. Monit. Assess.* **190** 85
- Almahasheer H, Serrano O, Duarte C M, Arias-Ortiz A, Masque P and Irigoien X 2017 Low carbon sink capacity of Red Sea mangroves *Sci. Rep.* **7** 9700
- Arias-Ortiz A *et al* 2018 A marine heatwave drives massive losses from the world's largest seagrass carbon stocks *Nat. Clim. Change* **8** 338–44
- Atwood T B *et al* 2017 Global patterns in mangrove soil carbon stocks and losses *Nat. Clim. Change* **7** 523–8
- Baskaran M, Bianchi T S and Filley T R 2017 Inconsistencies between <sup>14</sup>C and short-lived radionuclides-based sediment accumulation rates: effects of long-term remineralization *J. Environ. Radioact.* **174** 10–6
- Basson P W, Burchard Jr J, Hardy J and Price A 1977 *Biomes of the Western Arabian Gulf; Marine Life and Environments of Saudi Arabia* (Dhahran: Aramco Department of Loss Prevention and Environmental Affairs)
- Breithaupt J L, Smoak J M, Smith T J, Sanders C J and Hoare A 2012 Organic carbon burial rates in mangrove sediments: strengthening the global budget *Glob. Biogeochem. Cycles* **26** GB3011
- Brown D R *et al* 2016 Seagrass, mangrove and saltmarsh sedimentary carbon stocks in an urban estuary; Coffs Harbour, Australia *Reg. Stud. Mar. Sci.* **8** 1–6
- Campbell J E, Lacey E A, Decker R A, Crooks S and Fourqurean J W 2015 Carbon storage in seagrass beds of Abu Dhabi, United Arab Emirates *Estuaries Coasts* **38** 242–51
- Chen G, Azkab M H, Chmura G L, Chen S, Sastrosuwondo P, Ma Z, Dharmawan I W E, Yin X and Chen B 2017 Mangroves as a major source of soil carbon storage in adjacent seagrass meadows *Sci. Rep.* **7** 42406
- Chmura G L, Anisfeld S C, Cahoon D R and Lynch J C 2003 Global carbon sequestration in tidal, saline wetland soils *Glob. Biogeochem. Cycles* **17** 22–1
- Choi Y and Wang Y 2004 Dynamics of carbon sequestration in a coastal wetland using radiocarbon measurements *Glob. Biogeochem. Cycles* **18** GB4016
- Donato D C, Kauffman J B, Murdiyarso D, Kurnianto S, Stidham M and Kanninen M 2011 Mangroves among the most carbon-rich forests in the tropics *Nat. Geosci.* **4** 293–7
- Duarte C M, Losada I J, Hendriks I E, Mazarrasa I and Marba N 2013 The role of coastal plant communities for climate change mitigation and adaptation *Nat. Clim. Change* **3** 961–8
- Duarte C M, Middelburg J J and Caraco N 2005 Major role of marine vegetation on the oceanic carbon cycle *Biogeosciences* **2** 1–8
- Erftemeijer P L and Shuaib D A 2012 Seagrass habitats in the Arabian Gulf: distribution, tolerance thresholds and threats *Aquat. Ecosyst. Health Manage.* **15** 73–83
- Ewers Lewis C J, Carnell P E, Sanderman J, Baldock J A and Macreadie P I 2018 Variability and vulnerability of coastal 'blue carbon' stocks: a case study from southeast Australia *Ecosystems* **21** 263
- Ezcurra P, Ezcurra E, Garcillán P P, Costa M T and Aburto-Oropeza O 2016 Coastal landforms and accumulation of mangrove peat increase carbon sequestration and storage *Proc. Natl Acad. Sci.* **113** 4404–9
- Fard N J H, Ravanbakhsh M, Ramezani Z, Ahmadi M, Angali K A and Javid A Z 2015 Determination of mercury and vanadium concentration in *Johnius belangerii* (C) fish in Musa estuary in Persian Gulf *Mar. Pollut. Bull.* **97** 499–505
- Fourqurean J W *et al* 2012 Seagrass ecosystems as a globally significant carbon stock *Nat. Geosci.* **5** 505–9
- Gullström M *et al* 2018 Blue carbon storage in tropical seagrass meadows relates to carbonate stock dynamics, plant–sediment processes, and landscape context: insights from the western Indian Ocean *Ecosystems* **21** 551
- Hyndes G A, Nagelkerken I, Mcleod R J, Connolly R M, Lavery P S and Vanderklift M A 2014 Mechanisms and ecological role of carbon transfer within coastal seascapes *Biol. Rev.* **89** 232–54
- Kazemi A, Bakhtiari A R, Kheirabadi N, Barani H and Haidari B 2012 Distribution patterns of metals contamination in sediments based on type regional development on the intertidal coastal zones of the Persian Gulf, Iran *Bull. Environ. Contam. Toxicol.* **88** 100–3
- Krishnaswami S, Lal D, Prabhu N and Tamhane A 1971 Olivines: revelation of tracks of charged particles *Science* **174** 287–91
- Lokier S W and Fiorini F 2016 Temporal evolution of a carbonate coastal system, Abu Dhabi, United Arab Emirates *Mar. Geol.* **381** 102–13
- Loughland R A, Al-Abdulkader K A, Wyllie A and Burwell B O 2012 *Anthropogenic Induced Geomorphological Change along the Western Arabian Gulf coast* (Rijeka: INTECH Open Access Publisher)



- Lovelock C E *et al* 2017 Assessing the risk of carbon dioxide emissions from blue carbon ecosystems *Front. Ecol. Environ.* **15** 257–65
- Macreadie P I *et al* 2017 Carbon sequestration by Australian tidal marshes *Sci. Rep.* **7** 44071
- Madkour H A, Mansour A M, Ahmed A E-H N and El-Taher A 2014 Environmental texture and geochemistry of the sediments of a subtropical mangrove ecosystem and surrounding areas, Red Sea Coast, Egypt *Arabian J. Geosci.* **7** 3427–40
- Maher D T, Santos I R, Schulz K G, Call M, Jacobsen G E and Sanders C J 2017 Blue carbon oxidation revealed by radiogenic and stable isotopes in a mangrove system *Geophys. Res. Lett.* **44** 4889–96
- Masqué P, Sanchez-Cabeza J, Bruach J, Palacios E and Canals M 2002 Balance and residence times of 210 Pb and 210 Po in surface waters of the northwestern Mediterranean Sea *Cont. Shelf Res.* **22** 2127–46
- McLeod E, Chmura G L, Bouillon S, Salm R, Björk M, Duarte C M, Lovelock C E, Schlesinger W H and Silliman B R 2011 A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO<sub>2</sub> *Front. Ecol. Environ.* **9** 552–60
- Murray B C, Pendleton L, Jenkins W A and Sifleet S 2011 Green payments for blue carbon: economic incentives for protecting threatened coastal habitats *Nicholas Institute for Environmental Policy Solutions, Report NI* (North Carolina: Duke University) p 11
- Naser H A 2014 Marine ecosystem diversity in the Arabian Gulf: threats and conservation *Biodiversity—The Dynamic Balance of the Planet* ed O Grillo (Rijeka: InTech)
- Nellemann C, Corcoran E, Duarte C, Valdés L, de Young C, Fonseca L and Grimsditch G 2009 Blue Carbon: a rapid response assessment *United Nations Environment Programme, GRID-Arendal* (Norway: GRID-Arendal) p 80
- Pal J S and Eltahir E A B 2016 Future temperature in southwest Asia projected to exceed a threshold for human adaptability *Nat. Clim. Change* **6** 197–200
- Price A R and Coles S L 1992 Aspects of seagrass ecology along the western Arabian Gulf coast *Hydrobiologia* **234** 129–41
- Qurban M A, Krishnakumar P K, Joydas T V, Ashraf M, Manikandan T T, Abdulkader K P and Loughland K A 2011 Overview of Gulf Marine habitats *Marine Atlas of the Saudi Arabian Waters of the Arabian Gulf* ed A Khaled and R A L Al-Abdulkader (Dhahran: Saudi Aramco)
- Saenger P 2011 Mangroves and salt marshes *Marine Atlas of the Saudi Arabian Waters of the Arabian Gulf* ed Khaled A and Al-Abdulkader R A L (Dhahran: Saudi Aramco)
- Sanchez-Cabeza J A, Masqué P and Ani-Ragolta I 1998 210Pb and 210Po analysis in sediments and soils by microwave acid digestion *J. Radioanal. Nucl. Chem.* **227** 19–22
- Schile L M, Kauffman J B, Crooks S, Fourqurean J W, Glavan J and Megonigal J P 2017 Limits on carbon sequestration in arid blue carbon ecosystems *Ecol. Appl.* **27** 859–74
- Serrano O, Ricart A M, Lavery P S, Mateo M A, Arias-Ortiz A, Masqué P, Rozaimi M, Steven A and DUARTE C M 2016 Key biogeochemical factors affecting soil carbon storage in Posidonia meadows *Biogeosciences* **13** 4581–94
- Sheppard C, Al-Husiani M, Al-Jamali F, Al-Yamani F, Baldwin R, Bishop J, Benzoni F, Dutrieux E, Dulvy N K and Durvasula S R V 2010 The Gulf: a young sea in decline *Mar. Pollut. Bull.* **60** 13–38
- Siikamäki J, Sanchirico J N, Jardine S, McLaughlin D and Morris D 2013 Blue carbon: coastal ecosystems, their carbon storage, and potential for reducing emissions *Environ. Sci. Policy Sustain. Dev.* **55** 14–29
- Sugden W 1963 Some aspects of sedimentation in the Persian Gulf *J. Sediment. Res.* **33** 355–64
- Vaezi A, Karbassi A and Fakhraee M 2015 Assessing the trace metal pollution in the sediments of Mahshahr Bay, Persian Gulf, via a novel pollution index *Environ. Monit. Assess.* **187** 613
- Walton M E M, Al-Maslamani I, Skov M W, Al-Shaikh I, Al-Ansari I S, Kennedy H A and Le Vay L 2014 Outwelling from arid mangrove systems is sustained by inwelling of seagrass productivity *Mar. Ecol. Prog. Ser.* **507** 125–37
- Zahed M A, Rouhani F, Mohajeri S, Bateni F and Mohajeri L 2010 An overview of Iranian mangrove ecosystems, northern part of the Persian Gulf and Oman Sea *Acta Ecol. Sin.* **30** 240–4