

2018

The association between subclass-specific IgG Fc N-glycosylation profiles and hypertension in the Uygur, Kazak, Kirgiz, and Tajik populations

J. N. Liu

Mamatyusupu Dolikun

Jerko Štambuk

Irena Trbojević-Akmačić

J. Zhang

See next page for additional authors

10.1038/s41371-018-0071-0

This is an Author's Accepted Manuscript of: Liu, J. N., Dolikun, M., Štambuk, J., Trbojević-Akmačić, I., Zhang, J., Wang, H., ... & Liu, D. (2018). The association between subclass-specific IgG Fc N-glycosylation profiles and hypertension in the Uygur, Kazak, Kirgiz, and Tajik populations. *Journal of human hypertension*, 32, p 555–563. Original article is available [here](#)

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/4712>

Authors

J. N. Liu, Mamatyusupu Dolikun, Jerko Štambuk, Irena Trbojević-Akmačić, J. Zhang, Hao Wang, D. Q. Zheng, X. Y. Zhang, H. L. Peng, Z. Y. Zhao, D. Liu, Yang Sun, Q. Sun, Q. H. Li, J. X. Zhang, Ming Sun, W. J. Cao, Ana Momčilović, Genadij Razdorov, L. J. Wu, Alyce Russell, Y. X. Wang, Song Manshu, Gordan Lauc, and Wei Wang

1 **The association between subclass-specific IgG Fc N-glycosylation**
2 **profiles and hypertension in the Uygur, Kazak, Kirgiz and Tajik**
3 **populations**

4 JN Liu^{1,#}, M Dolikun^{2,#}, J Štambuk^{3,#}, I Trbojević-Akmačić³, J Zhang¹, H Wang¹, DQ
5 Zheng¹, XY Zhang¹, HL Peng¹, ZY Zhao¹, D Liu¹, Y Sun¹, Q Sun¹, QH Li¹, JX
6 Zhang¹, M Sun¹, WJ Cao¹, A Momčilović³, G. Razdorov³, LJ Wu¹, A Russell⁴, YX
7 Wang¹, MS Song^{1,4}, G Lauc^{3,5}, W Wang^{1,4}

8 ¹Beijing Key Laboratory of Clinical Epidemiology, School of Public Health, Capital
9 Medical University, Beijing, China.

10 ²College of the Life Sciences and Technology, Xinjiang University, Urumqi, China.

11 ³Genos Glycoscience Research Laboratory, Zagreb, Croatia.

12 ⁴School of Medical and Health Sciences, Edith Cowan University, Perth, Australia.

13 ⁵Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia.

14 Correspondence:

15 Associate Professor MS Song, School of Public Health, Capital Medical University,

16 No. 10 Xitoutiao, You An Men Wai, Fengtai District, Beijing 100069, China

17 E-mail: songms@ccmu.edu.cn

18 Tel.: 0086 10 83911497

19 &

1 Associate Professor YX Wang, School of Public Health, Capital Medical University,

2 No. 10 Xitoutiao, You An Men Wai, Fengtai District, Beijing 100069, China

3 E-mail: wangy@ccmu.edu.cn

4 Tel: 0086 10 83911779, Fax: 0086 10 83911508

5 #These authors contributed equally to this work.

1 **ABSTRACT**

2 Hypertension results from the interaction of genetic and acquired factors. IgG occurs
3 in the form of different subclasses, of which the effector functions show significant
4 variation. The detailed differences between the glycosylation profiles of the individual
5 IgG subclasses may be lost in a profiling method for total IgG *N*-glycosylation. In this
6 study, subclass-specific IgG Fc glycosylation profile was investigated in the four
7 northwestern Chinese minority populations, namely, Uygur (UIG), Kazak (KZK),
8 Kirgiz (KGZ) and Tajik (TJK), composed of 274 hypertensive patients and 356
9 healthy controls. The results showed that 10 directly measured IgG *N*-glycan traits (i.e.
10 IgG1G0F, IgG2G0F, IgG2G1FN, IgG2G1FS, IgG2G2S, IgG4G0F, IgG4G1FS,
11 IgG4G1S, IgG4G2FS, and IgG4G2N) representing galactosylation and sialylation are
12 significantly associated with hypertension, with IgG4 consistently showed weaker
13 associations of its sialylation, across the four ethnic groups. We observed a modest
14 improvement on the AUC of ROC-curve when the IgG Fc *N*-glycan traits are added
15 into the glycan-based model (difference between AUCs, 0.044, 95% CI: 0.016-0.072,
16 $P= 0.002$). The AUC of the diagnostic model indicated that the subclass-specific IgG
17 Fc *N*-glycan profiles provide more information reinforcing current models utilizing
18 age, gender, BMI and ethnicity, and demonstrate the potential of subclass-specific
19 IgG Fc *N*-glycosylation profiles to serve as a biomarker for hypertension. Further

1 research is however required to determine the additive value of subclass-specific IgG

2 Fc *N*-glycosylation on top of biomarkers which are currently used.

3

1

Summary Table

2 *What is known about topic?*

3 ● The prevalence of hypertension in the Chinese minority populations residing in
4 the northwestern China ranked at the forefront of China.

5 ● Previous studies showed an association between hypertension and IgG *N*-
6 glycome composition in European (i.e. Croatian Korčula, Croatian Vis, and
7 Scottish Orkney), Asian (i.e. Chinese Han and Chinese Kazak) populations.

8 ● The relationship between IgG glycosylation and hypertension in other
9 populations remains unknown.

10 ● Previous studies on total IgG *N*-glycans do not allow to distinguish between Fc
11 and Fab associated *N*-glycans, as such the contribution of changes in Fab *N*-
12 glycosylation cannot be excluded.

13 *What this study adds?*

14 ● This study provides evidence that the decreased IgG galactosylation and
15 sialylation were correlated with hypertension in the four ethnic groups (i.e. Uyгур,
16 Kazak, Kirgiz and Tajik) residing in the northwestern China.

17 ● We describe the differences between the plasma Fc glycosylation profiles
18 particularly of IgG1, IgG2 and IgG4 in detail, which may be lost in a profiling
19 method for total IgG *N*-glycosylation.

20 ● The AUC of the glycan-based model indicated that the IgG glycan profiles
21 provide more information reinforcing current models utilizing gender, age, BMI
22 and ethnicity, and demonstrate the potential of subclass-specific IgG Fc *N*-
23 glycosylation profiles to serve as a biomarker for hypertension.

24

1 INTRODUCTION

2 Hypertension is a chronic, asymptomatic condition that increases individuals' risk
3 of developing several diseases, including coronary artery disease, stroke, and renal
4 failure.¹ Hypertension was the leading risk factor for global disease in 2010,² and over
5 20% of adults (≥ 18 years) worldwide were hypertensive, accounting for nearly 10
6 million deaths in 2014.³ The prevalence of hypertension continues to increase in
7 China from 18% in 2002 to 23.2% in 2015,⁴ in which participants were recruited by
8 the use of a stratified multistage cluster sampling design.

9 Glycosylation is one of the most widespread post-translational modifications
10 capable of increasing significant structural diversity to proteins, which is known to
11 reflect the physiological state of an organism and changes thereof.⁵ For
12 immunoglobulin G (IgG), which occupies a central role in the immune system, it is
13 known that the conserved *N*-glycan located at asparagine 297 on the fragment
14 crystallisable (Fc) part of IgG can modulate inflammatory responses: a lack of core
15 fucose, galactose and *N*-acetylneuraminic (sialic) acid increases the ability of IgG to
16 induce antibody-dependent cell-mediated cytotoxicity (ADCC).⁶ In spite of the
17 pivotal role of IgG glycosylation in many physiological and pathological processes,
18 the monitoring of IgG glycosylation is often neglected in clinical and immunological
19 research.⁷

1 IgGs consist of four subclasses (i.e. IgG1–4), which bind their antigen targets via
2 the fragment antigen binding (Fab) domain and exert their effector functions via the
3 fragment crystallizable (Fc) domain.⁸ Fc glycans are essential structural components
4 of the IgG molecule and minor changes in glycan composition can significantly alter
5 the conformation of the Fc region changing the interaction with receptor proteins and
6 thus modulating the effector functions of IgG. The assignment of glycans to the
7 specific Fc glycosylation sites of IgG subclasses is pivotal for deducing the functional
8 implications of the observed glycosylation features.⁹ Additionally, each of the IgG
9 subclasses has a unique binding profile to each Fc gamma receptors (FcγRs), and their
10 expression profiles are highly variable between different immune cells of myeloid and
11 NK cell origin.¹⁰ Most of the modulating effects of IgG Fc glycans have been reported
12 for IgG1 and may not apply to IgG2, stressing the relevance of subclass and site
13 specific IgG glycosylation profiling.⁹ The differences between the Fc *N*-glycosylation
14 profiles of the individual IgG subclasses may be lost in total IgG *N*-glycosylation (Fc
15 and Fab associated *N*-glycans).

16 The prevalence of hypertension in the Chinese minority populations residing in
17 the Xinjiang Uygur Autonomous Region located in the northwestern China is
18 relatively higher than that in the Chinese Han in China.¹¹ The northwestern Chinese
19 minority populations, including Uygur (UIG), Kazak (KZK), Kirgiz (KGZ) and Tajik

1 (TJK), etc., are overwhelmingly Muslim, and have their own language, religious
2 beliefs and lifestyles differed largely from either Han Chinese or European
3 populations.¹² They are classically well-defined isolated populations, and these ethnic
4 groups differ substantially in terms of morphological and genetic characteristics.¹³
5 These differences in genetics and environmental factors together with abnormal
6 higher prevalence of hypertension might imply different pathogenesis of hypertension.

7 Our previous study has reported that individual variation of *N*-glycosylation of
8 IgG might contribute to hypertension in European (i.e. Croatian Korčula, Croatian
9 Vis, and Scottish Orkney) and Asian (i.e. Chinese Han and Chinese Kazak)
10 populations.^{14,15} Investigating other populations would be an expedient way to
11 evaluate the generalizability of the current findings. In this study, we further validated
12 these findings by investigating Fc *N*-glycosylation profiles of human IgG subclasses,
13 investigated to what extent the Fc *N*-glycome of IgG subclasses correlates with the
14 occurrence of hypertension, and demonstrated the potential of subclass-specific IgG
15 Fc *N*-glycosylation profiles to serve as a biomarker for hypertension in the four
16 northwestern Chinese minority populations.

17 **MATERIALS AND METHODS**

18 Study population

19 This population-based, case-control survey was conducted in the Xinjiang Uyghur

1 Autonomous Region, China, from August 2014 to August 2016. A total of 630
2 participants were recruited from 170 UIG, 150 KZK, 168 KGZ and 142 TJK. The
3 demographic characteristics of the participants, including ethnicity, gender, age, and
4 history of medications, were collected by a questionnaire.

5 Participants were included if they aged more than 18 years, self-reported the
6 respective UIG, KZK, KGZ and TJK ethnicity without intermarriage history with
7 other ethnic groups within at least the past three generations. Individuals with a
8 diagnosis of specific severe diseases concerning the cardiovascular system,
9 respiratory system, genitourinary system, digestive system, and hematic system, were
10 excluded. All participants gave written informed consent. This study was approved by
11 the local community leaders and the ethics committee of the Capital Medical
12 University, Beijing, China.

13 Anthropometric measurements, including weight, height, etc., were obtained
14 using standardized techniques.¹⁴ The body mass index (BMI) was calculated by the
15 formula weight (in kilograms)/height (in square meters). Peripheral bloods of the
16 participants were collected in EDTA-anti-coagulated tubes after an overnight fast.
17 Fasting blood glucose (FBG), high-density lipoprotein cholesterol (HDL), low-density
18 lipoprotein cholesterol (LDL), triglycerides (TG) and serum total cholesterol (TC)
19 were assayed under the standard procedures in the local hospital.

1 Dyslipidemia defined based on the 2016 Guidelines for Prevention and
2 Treatment of Dyslipidemia in Adults in China: TC \geq 6.2 mmol/L, TG \geq 2.3
3 mmol/L, LDL \geq 4.1 mmol/L or HDL $<$ 1.0 mmol/L.¹⁶ Systolic blood pressure (SBP)
4 and diastolic blood pressure (DBP) were measured three times in a day with a
5 mercury sphygmomanometer. Hypertension was defined as an average SBP \geq 140
6 mmHg or an average DBP \geq 90 mmHg, or self-reported current hypertension.¹⁷

7 IgG Fc *N*-glycopeptide analysis

8 IgG was isolated from the plasma samples of all the participants after overnight
9 fasting, and each IgG subclass was analyzed by nano-reverse phase UPLC-MS using
10 a sheath-flow ESI sprayer interface as described previously.^{18,19} Glycopeptide data
11 were extracted by in-house python script. In short, m/z was recalibrated internally,
12 and intensities of top four isotopologues per charge per glycopeptide were extracted
13 from LC elution bins. Glycopeptide area was produced by summing intensities across
14 all relevant isotopologues and charge states. The *N*-glycopeptide signals were
15 normalized by dividing each by the total signal intensity of all *N*-glycopeptides
16 belonging to that subclass, yielding percentage data amounting to 100% per subclass.
17 Along with the 20 glycopeptide structures per subclass of IgG, an additional 7 derived
18 traits (agalactosylation, galactosylation, monogalactosylation, digalactosylation,
19 sialylation, bisecting *N*-acetylglucosamine [GlcNAc] and fucosylation) representing

1 composite traits were defined by taking into account these glycopeptide structures

2 amounts per each subclass:^{8,14}

3
$$\text{Agalactosylation (A)} = G0 + G0F + G0FN + G0N$$

4
$$\text{Galactosylation (G)} = G1 + G1F + G1FN + G1N + G2 + G2F + G2FN + G2N$$

5
$$\text{Monogalactosylation (M)} = G1 + G1F + G1FN + G1N$$

6
$$\text{Digalactosylation (D)} = G2 + G2F + G2FN + G2N$$

7
$$\text{Sialylation (S)} = G1FS + G1FNS + G1S + G1NS + G2FS + G2FNS + G2NS +$$

8
$$G2S$$

9
$$\text{Bisecting GlcNAc (B)} = G0FN + G0N + G1FN + G1FNS + G1N + G1NS +$$

10
$$G2FN + G2FNS + G2NS + G2N$$

11
$$\text{Fucosylation (F)} = G0F + G0FN + G1F + G1FN + G1FNS + G1FS + G2F +$$

12
$$G2FN + G2FNS + G2FS$$

13 Statistical analysis

14 To remove experimental variation from measurements, normalization and batch

15 correction were performed on the LC-MS glycopeptide data.⁸ Normality distributions

16 were checked by the Kolmogorov-Smirnov tests. Continuous variables underlying

17 normality distribution were described by mean together with standard deviation (SD),

18 otherwise median together with interquartile range were used. Considering that nearly

19 all the glycan traits cannot be assumed normal distributed, Mann-Whitney U test was

1 used to compare the differences of IgG subclass-specific Fc glycopeptides between
2 the hypertension patients and the healthy controls. Associations between glycans and
3 disease status were performed using a logistic regression with age, gender, ethnicity,
4 BMI, FBG and dyslipidemia included as covariates. Pooling analysis was carried out
5 by introducing dummy variables representing the different ethnic groups.

6 Dimension reduction based on LASSO (least absolute shrinkage and selection
7 operator) method was performed to select *N*-glycopeptide traits used in the
8 discriminant analysis.¹⁴ To classify the hypertension patients with the healthy controls,
9 a logistic regression model was applied. Two different predicted models were
10 considered: a baseline model included gender, age, BMI and ethnicity as the
11 covariates, and a glycan-based model included the significantly differed IgG Fc
12 glycopeptides together with the aforementioned covariates. Receiver operating
13 characteristic (ROC) curve analyses were used to evaluate the diagnostic ability in
14 classifying the hypertension patients from the healthy controls. The area under the
15 receiver operating characteristic curve (AUC) of the 2 classification models were
16 compared using MedCalc (Version 17.9.7).

17 All the analyses were performed using the SPSS (Version 21.0, IBM) software
18 and R packages (Version 2.7.2). For the multiple corrections, the false discovery rate
19 (FDR) was used based on the Benjamini-Hochberg procedure.²⁰ All the tests were 2-

1 sided, and $P < 0.05$ was considered statistically significant.

2 **RESULTS**

3 By performing LC-ESI-MS analysis on IgG glycopeptides, Fc glycosylation
4 profiles of IgG1, IgG2 and IgG4 were determined for 630 participants (i.e. 170 UIG,
5 150 KZK, 168 KGZ, and 142 TJK) in the study. Fc glycopeptides of IgG2 and IgG3
6 share the same peptide sequence,²¹ and so the commonly used protein G enrichment
7 of IgG gives a joint profile for both IgG2 and IgG3. It has been shown to capture IgG1,
8 2 and 4, but has a much lower binding affinity for IgG3 under the enrichment
9 conditions we use,⁷ thus allowing for near-separate glycoprofiling of IgG2.

10 All 630 participants provided complete biometrics and glycan level data, and
11 were therefore used in the further analyses. Descriptive information on the
12 hypertension patients and the healthy controls for each of the four ethnic groups are
13 shown separately (Table S1) and pooled together in Table 1. The age, gender, BMI,
14 SBP, DBP, TC, HDL, LDL and FBG significantly differed in hypertension patients
15 compared to healthy controls.

16 The comparisons of directly measured subclass-specific IgG Fc glycomic
17 composition between hypertension patients and healthy controls of each ethnicity and
18 of the pooled participants from all four ethnic groups are shown in Table S2, while
19 derived traits are shown in Table S3 and Figure 1. Subclass-specific IgG Fc *N*-

1 glycopeptides were significantly associated with hypertension in the pooled
2 participants from all four ethnic groups after controlling for gender, age, BMI,
3 dyslipidemia, FBG and ethnicity as summarized in Table S2 and Table S3. Ten
4 directly measured IgG subclass-specific Fc glycopeptide structures and 14 derived
5 glycan traits are significantly associated with hypertension in the pooling of the four
6 ethnic groups, primarily reflecting the decrease of galactosylation and sialylation in
7 hypertension patients. Subclass-specific IgG Fc glycopeptides that statistically
8 differed between the hypertensive patients and healthy controls in the pooling of four
9 ethnic groups were not consistently significant among the comparisons between each
10 ethnicity, although these differences were in similar trends (Table S2 and Table S3).

11 Two different models were built using Logistic regression (Table 4, Figure 2).
12 The AUC of 0.688 (95% CI: 0.650-0.724) in the baseline model utilized gender, age,
13 BMI and ethnicity as the covariates, and 0.732 (95% CI: 0.696-0.766) in the glycan-
14 based model utilized five significantly differed subclass-specific IgG Fc glycopeptide
15 *N*-glycans (IgG1G0F, IgG2G0F, IgG4G0F, IgG4G2FS and IgG4G2N) together with
16 the aforementioned covariates, respectively. The AUCs showed modest improvement
17 in the glycan-based diagnostic model compared to the baseline model, with an
18 increment of 0.044 (95% CI: 0.016-0.072, $P = 0.002$).

19 **DISCUSSION**

1 Using LC-ESI-MS, this study represents the first comprehensive analysis of Fc
2 *N*-glycosylation profiling of human IgG subclasses showing similar associations with
3 hypertension in the four ethnic minorities in northwestern China. In this study, ten
4 directly measured subclass-specific IgG Fc *N*-glycan traits indicative of
5 galactosylation and sialylation presented a significant association with hypertension.
6 The IgG subclasses displayed subtle differences in directly measured glycopeptide
7 structures, with significant differences in sialylation resulting primarily between the
8 IgG1 and IgG2 subclasses, the potential reason for any underlying difference in
9 biological role. When the five selected glycan traits (IgG1G0F, IgG2G0F, IgG4G0F,
10 IgG4G2FS and IgG4G2N) were added into the glycan-based model, the AUC showed
11 modest improvement (difference between AUCs, 0.044, 95% CI: 0.016-0.072, $P =$
12 0.002).

13 We previously demonstrated that plasma *N*-glycome is associated with blood
14 pressure, with increased triantennary glycans and decreased core-fucoseglycans in the
15 subjects with elevated blood pressure, in both Chinese (Han and Kazak) and Croatian
16 (Korčula and Vis).²² Previously we showed that galactosylation and sialylation of
17 total IgG decreased, and fucosylated structures and bisecting GlcNAc of total IgG
18 increased in patients of hypertension.¹⁵ In addition, 14 IgG subclass-specific Fc
19 glycopeptide structures, along with one derived glycosylation trait in subclasses IgG2

1 and IgG4, are correlated with SBP and/or DBP in the Kazakh population.¹⁴ Here, we
2 advanced the understanding of the association between the subclass-specific IgG Fc
3 glycopeptides, which may be lost in total IgG *N*-glycosylation, and hypertension in
4 the four northwestern Chinese minority populations. Our findings of decreasing
5 galactosylation and sialylation in hypertensive patients are consistent with those in
6 Chinese Han, Kazak and European populations. However, the non-significant changes
7 of core-fucosylation and bisecting GlcNAc in hypertension patients and healthy
8 controls are inconsistent with our previous findings in Chinese Han, Kazak and
9 European populations.

10 Glycans do not have a direct genetic template, thus glycan structures attached to
11 proteins are determined by complex dynamic interactions between genetic and
12 environmental factors.²³ The *HNF1A* gene, known for its common variants associated
13 with type 2 diabetes (T2DM),²⁴ is a genetic locus significant associated with total
14 plasma *N*-glycome as a regulator of fucosylation identified in the individuals of
15 European ancestry.²⁶ Moreover, *HNF1A* regulates expression of *FUT8*,²⁶ and is
16 associated with the susceptibility of T2DM in the individuals of European ancestry.²⁷
17 However, some variants in *HNF1A* and *FUT8* were not statistically associated with
18 the susceptibility of T2DM patients of Uyghur.²⁵ The Chinese ethnic groups (e.g.,
19 Tajik, Uyghur, Kazakh, Kirgiz and Hui) are admixed populations with both eastern

1 and western Eurasian ancestries, thus resulting in complex genetic diversity.¹³ The
2 widespread differences in genetic structure and environmental conditions among these
3 ethnic groups may partly explain the inconsistent association between IgG Fc *N*-
4 glycans and hypertension among the different ethnic groups.¹⁴

5 The marked decrease in galactosylation and increase in agalactosylation of the
6 three subclasses (IgG 1, 2 and 4) observed in this study have been demonstrated in
7 many other diseases and traits, such as the Parkinson's disease and the effects of age
8 on allergic disease.^{8,26} Additionally, in many autoimmune diseases, such as
9 rheumatoid arthritis (RA), systemic lupus erythematosus, juvenile onset chronic
10 arthritis, and Crohn's disease, marked increase in agalactosylation has been
11 demonstrated.²⁷⁻²⁹ It can be assumed that the increase in agalactosylation and the
12 resulting pro-inflammatory effect may be associated with many other diseases. Lack
13 of galactosylation is associated with a pro-inflammatory state of IgG through
14 activation of the complement cascade.³⁰ The previous study also demonstrated that
15 high galactosylation promotes cooperative signaling of the FcγRIIB with dectin-1
16 resulting in an inhibitory signaling pathway that blocks pro-inflammatory effector
17 functions.³¹ A possible explanation for this decreased galactosylation is post-
18 translational modifications of the enzyme β4-galactosyltransferase-1.³² Thus the
19 decrease in galactosylation may be the common pathogeny of multiple diseases

1 associated with decreased immunosuppressive potential of circulating IgG.

2 We find that sialylation of IgG1 and IgG2 decreased in individuals with the state
3 of hypertension. With regards to IgG4, the observed differences could have been
4 attributed to a decrease in analytical precision resulting from the low concentrations
5 of IgG4 in comparison to other IgG subclasses. More work is required, in particular
6 on antigen specific IgG subclasses, to improve our understanding of the role of IgG
7 glycosylation.

8 Sialylation plays an important role in the inflammatory potential of IgG.
9 Addition of sialic acid to IgG converts its function from pro- to anti-inflammatory by
10 decreased binding to Fcγ receptors.³³ IgG acquires anti-inflammatory properties upon
11 Fc sialylation, which is reduced upon the induction of an antigen-specific immune
12 response. This differential sialylation may provide a switch from innate anti-
13 inflammatory activity in the steady state to generating adaptive pro-inflammatory
14 effects upon antigenic challenge.³² Therefore decreased sialylation might play
15 pathogenic roles in hypertension, but the pathogenic mechanism of sialylation was
16 unclear.

17 Alterations of the immune response have been implicated in the pathogenesis of
18 hypertension for more than 5 decades.¹⁵ Chae et al observed significantly graded
19 relationships between blood pressure and levels of sICAM-1 as well as IL-6, which

1 suggested the increased blood pressure should be a stimulus for inflammation.³⁴
2 Marvar et al further found that hypertension has been involved in both central and
3 peripheral mechanisms of Tlymphocyte activation and vascular inflammation
4 produced by angiotensin II-induced hypertension.³⁵ Additionally, several
5 inflammatory markers including C-reactive protein (CRP) are strongly associated
6 with the risk of hypertension.³⁶ Therefore, the association between inflammation and
7 hypertension suggest that IgG glycosylation might contribute to the pathogenesis of
8 hypertension via the alternation of the inflammation, and as such could be a valuable
9 prognostic model for hypertension.

10 As glycan profiles are associated with genetic, metabolic and environmental
11 influences, this adds to their predictive potential. We observed a modest improvement
12 when adding the IgG Fc *N*-glycopeptide profiles to clinical risk factors. The AUC of
13 the glycan-based model indicated that the IgG glycan profiles provide more
14 information reinforcing current models utilizing gender, age, BMI and ethnicity.

15 The case-control nature of this study makes it impossible to infer the causal-
16 effect relationship between IgG Fc *N*-glycans and hypertension. One of the
17 shortcomings in this study is the lack of prospective follow-up for the hypertension
18 outcome. While the study design adopted in this project shows a significant
19 association between hypertension and IgG subclass-specific Fc *N*-glycopeptide

1 glycosylation, it is not conclusive as to whether the expressed IgG subclass Fc *N*-
2 glycopeptide glycosylation proceeds the symptoms of hypertension in the study
3 cohorts. As such, no causal claims can be made as the AUCs can only be interpreted
4 in a case-control setting where patients have already been diagnosed for and are being
5 treated for this health event. Notwithstanding this, the current study not only presents
6 the potential for the IgG subclass-specific glycan information to be indicative of
7 hypertension, but additionally reinforces current diagnostic models of hypertension.
8 Another potential limitation of this study was that some cases were on their
9 antihypertensive medication when their blood samples were collected. Thus, it was
10 not possible to distinguish between antihypertensive medication effects on the IgG
11 glycome and effects associated with the pathophysiology of the disease itself. It
12 should be noted that other risk factors related to hypertension, such as family history
13 of hypertension and smoking, were not obtainable during this study and were not
14 included. The absence of controlling for potential risk factors may confound the
15 association, but at certain extend the consistent results among populations
16 investigated with different ethnic background might help this study to profile
17 objectively the association observed between IgG glycosylation and hypertension.
18 Future studies in prospective cohorts should be conducted to investigate how this
19 biomarker potential could be used for personalized approaches in prevention and

1 treatment of the disease as well as its role in hypertension complications.

2 To conclude, we found that the robust associations of subclass-specific IgG Fc
3 *N*- glycosylation profiles may serve as an informative biomarker indicative of the pro-
4 inflammatory and biological state shared by hypertension. Future studies should be
5 directed at in-depth pathophysiological insights that can be derived from IgG Fc *N*-
6 glycan associations with hypertension. It is expected that prospective follow-up
7 studies with additional information about inflammatory biomarkers and genetic
8 studies might further shed light on the causality, the potential as a biomarker for
9 complex inflammatory processes, and the true predictive capacity of IgG subclass
10 specific Fc *N*-glycans in hypertension.

11 **CONFLICT OF INTEREST**

12 G. Lauc is the founder and owner of Genos Ltd, a private research organization that
13 specializes in high-throughput glycomic analysis and has several patents in this field.

14 J. Štambuk, I. Trbojević-Akmačić, A. Momčilović and G. Razdorov are employees of
15 Genos Ltd.

16 **ACKNOWLEDGEMENTS**

17 This study was supported by the National Natural Science Foundation of China
18 (81573215, 31460285 and 81370083), Australian National Health and Medical
19 Research Council (NH&MRC-APP1046711). Genos has received funding from

1 European Commission FP7 grants MIMOmics (contract #305280), HTP-GlycoMet
2 (contract #324400) and PainOmics (contract #602736), and H2020 grants GlySign
3 (contract #722095), GlyCoCan (contract #676412), SYSCID (contract #733100) and
4 IMforFuture (contract #721815) as well as funding from the European Structural and
5 Investments funds for projects "New generation of high throughput glycoanalytical
6 services (contract #KK.01.2.1.01.0003) and "Croatian National Centre of Research
7 Excellence in Personalized Healthcare" (contract #KK.01.1.1.01.0010). Hao Wang
8 was supported by China Scholarship Council (CSC No. 201708110200). The authors
9 thank the directors and relevant staff of relevant Health Bureaus for their support for
10 recruiting the Uygur, Kazak, Kirgiz and Tajik participants, and these volunteers and
11 community leaders for their participation and support.

12 REFERENCES

- 13 1. World Health Organization. A global brief on hypertension: Silent killer, global public health crisis
14 (2013). http://www.who.int/cardiovascular_diseases/publications/global_brief_hypertension/en/
15 (Date of access: 11/28/2017)
- 16 2. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, *et al.* A comparative risk
17 assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21
18 regions, 1990-2010: a systematic analysis for the global burden of disease study 2010. *Lancet.* 2012;
19 **380**(9859): 2224-60.
- 20 3. World Health Organization. Global status report on noncommunicable diseases 2014: World
21 Health Organization. 2014.
- 22 4. Wang Z, Chen Z, Zhang L, Wang X, Hao G, Zhang Z, *et al.* Status of hypertension in China: results
23 from the China hypertension survey, 2012-2015. *Circulation.* 2018. doi:
24 10.1161/CIRCULATIONAHA.117.032380.
- 25 5. Varki A. Biological roles of oligosaccharides: all of the theories are correct. *Glycobiology.* 1993; **3**(2):
26 97-130.

- 1 6.Plomp R, Ruhaak LR, Uh HW, Reiding KR, Selman M, Houwing-Duistermaat JJ, *et al.* Subclass-
2 specific IgG glycosylation is associated with markers of inflammation and metabolic health. *Sci Rep.*
3 2017; **7**(1): 12325.
- 4 7.Wuhrer M, Stam JC, Fe VDG, Koeleman CA, Verrips CT, Dolhain RJ, *et al.* Glycosylation profiling
5 of immunoglobulin G (IgG) subclasses from human serum. *Proteomics.* 2007; **7**(22): 4070-81.
- 6 8.Pezer M, Stambuk J, Perica M, Razdorov G, Banic I, Vuckovic F, *et al.* Effects of allergic diseases
7 and age on the composition of serum IgG glycome in children. *Sci Rep.* 2016; **6**: 33198.
- 8 9.Bakoviä MP, Selman MH, Hoffmann M, Rudan I, Campbell H, Deelder AM, *et al.* High-throughput
9 IgG Fc *N*-glycosylation profiling by mass spectrometry of glycopeptides. *J Proteome Res.* 2013;
10 **12**(2): 821-31.
- 11 10.Vidarsson G, Dekkers G, Rispens T. IgG subclasses and allotypes: from structure to effector
12 functions. *Front Immunol.* 2014; **5**: 520.
- 13 11.Keser T, Vuckovic F, Barrios C, Zierer J, Wahl A, Akinkuolie AO, *et al.* Effects of statins on the
14 immunoglobulin G glycome. *Biochim Biophys Acta.* 2017; **1861**(5 Pt A): 1152-58.
- 15 12.Zhai F, He Y, Wang Z, Hu Y. Status and characteristic of dietary intake of 12 minority nationalities
16 in China. *Wei Sheng Yan Jiu.* 2007; **36**(5): 539-41.
- 17 13.Li J, Lou H, Yang X, Lu D, Li S, Jin L, *et al.* Genetic architectures of ADME genes in five Eurasian
18 admixed populations and implications for drug safety and efficacy. *J Med Genet.* 2014; **51**(9): 614-22.
- 19 14.Gao Q, Dolikun M, J Š, Wang H, Zhao F, Yiliham N, *et al.* Immunoglobulin G *N*-Glycans as
20 potential postgenomic biomarkers for hypertension in the Kazakh population. *Omic A Journal of*
21 *Integrative Biology.* 2017; **21**(7): 380.
- 22 15.Wang Y, Klarić L, Yu X, Thaqi K, Dong J, Novokmet M, *et al.* The association between
23 glycosylation of immunoglobulin G and hypertension. *Medicine.* 2016; **95**(17): e3379.
- 24 16.Junren C, Runlin G, Shuiping Z, Guoping L, Dong Z, Jianjun L. Guidelines for the prevention and
25 treatment of dyslipidemia in Chinese adults (Revised Edition 2016). *Chinese Circulation Journal.*
26 2016; **31**(10): 7-28.
- 27 17.Gu D, Reynolds K, Wu X, Chen J, Duan X, Muntner P, *et al.* Prevalence, awareness, treatment, and
28 control of hypertension in china. *Hypertension.* 2002; **40**(6): 920-7.
- 29 18.Selman MH, Derks RJ, Bondt A, Palmblad M, Schoenmaker B, Koeleman CA, *et al.* Fc specific
30 IgG glycosylation profiling by robust nano-reverse phase HPLC-MS using a sheath-flow ESI sprayer
31 interface. *J Proteomics.* 2012; **75**(4): 1318-29.
- 32 19.Gornik O, Pavic T, Lauc G. Alternative glycosylation modulates function of IgG and other proteins
33 - implications on evolution and disease. *Biochim Biophys Acta.* 2012; **1820**(9): 1318-26.
- 34 20.Benjamini Y, Hochberg Y. Controlling The false discovery rate - A practical and powerful approach
35 to multiple testing. *Journal of the Royal Statistical Society.* 1995; **57**(1): 289-300.
- 36 21.Plomp R, Bondt A, de Haan N, Rombouts Y, Wuhrer M. Recent advances in clinical
37 glycoproteomics of immunoglobulins (Igs). *Mol Cell Proteomics.* 2016; **15**(7): 2217-28.
- 38 22.Lu JP, Knežević A, Wang YX, Rudan I, Campbell H, Zou ZK, *et al.* Screening novel biomarkers

- 1 for metabolic syndrome by profiling human plasma N-glycans in Chinese Han and Croatian
2 populations. *J Proteome Res.* 2011; **10**(11): 4959-69.
- 3 23.Krištić J, Zoldoš V, Lauc G. Complex genetics of protein N -Glycosylation. In: Endo T, Seeberger
4 PH, Hart GW, Wong C-H, Taniguchi N, (eds). *Glycoscience: biology and medicine*. Blackwell
5 Publishing Ltd: Japan, 2014, pp 1-7
- 6 24.Voight BF, Scott LJ, Steinthorsdottir V, Morris AP, Dina C, Welch RP, *et al.* Twelve type 2
7 diabetes susceptibility loci identified through large-scale association analysis. *Nat Genet.* 2010; **42**(7):
8 579-89.
- 9 25.Zhao F, Mamatjusupu D, Wang Y, Fang H, Wang H, Gao Q, *et al.* The Uyghur population and
10 genetic susceptibility to type 2 diabetes: potential role for variants in CAPN10,APM1 and FUT6
11 genes. *Journal of Cellular & Molecular Medicine.* 2016; **20**(11): 2138-47.
- 12 26.Russell AC, Simurina M, Garcia MT, Novokmet M, Wang Y, Rudan I, *et al.* The N-glycosylation
13 of immunoglobulin G as a novel biomarker of Parkinson's disease. *Glycobiology.* 2017; **27**(5): 501-
14 10.
- 15 27.Parekh R, Isenberg D, Rook G, Roitt I, Dwek R, Rademacher T. A comparative analysis of disease-
16 associated changes in the galactosylation of serum IgG. *J Autoimmun.* 1989; **2**(2): 101-14.
- 17 28.Parekh RB, Dwek RA, Sutton BJ, Fernandes DL, Leung A, Stanworth D, *et al.* Association of
18 rheumatoid arthritis and primary osteoarthritis with changes in the glycosylation pattern of total
19 serum IgG. *Nature.* 1985; **316**(6027): 452-7.
- 20 29.Bond A, Alavi A, Axford JS, Bourke BE, Bruckner FE, Kerr MA, *et al.* A detailed lectin analysis of
21 IgG glycosylation, demonstrating disease specific changes in terminal galactose and N-
22 acetylglucosamine. *J Autoimmun.* 1997; **10**(1): 77-85.
- 23 30.Mihai S, Nimmerjahn F. The role of Fc receptors and complement in autoimmunity. *Autoimmun*
24 *Rev.* 2013; **12**(6): 657-60.
- 25 31.Karsten CM, Pandey MK, Figge J, Kilchenstein R, Taylor PR, Rosas M, *et al.* Anti-inflammatory
26 activity of IgG1 mediated by Fc galactosylation and association of FcγRIIB and dectin-1. *Nat Med.*
27 2012; **18**(9): 1401-06.
- 28 32.Dall'Olio F, Vanhooren V, Chen CC, Slagboom PE, Wuhrer M, Franceschi C. N-glycomic
29 biomarkers of biological aging and longevity: a link with inflammaging. *Ageing Res Rev.* 2013; **12**(2):
30 685-98.
- 31 33.Kaneko Y, Nimmerjahn F, Ravetch JV. Anti-inflammatory activity of immunoglobulin G resulting
32 from Fc sialylation. *Science.* 2006; **313**(5787): 670-3.
- 33 34.Chae CU, Lee RT, Rifai N, Ridker PM. Blood pressure and inflammation in apparently healthy men.
34 *Hypertension.* 2001; **38**(3): 399-403.
- 35 35.Marvar PJ, Thabet SR, Guzik TJ, Lob HE, McCann LA, Weyand C, *et al.* Central and peripheral
36 mechanisms of T-lymphocyte activation and vascular inflammation produced by angiotensin II-
37 induced hypertension. *Circ Res.* 2010; **107**(2): 263-70.
- 38 36.Solak Y, Afsar B, Vaziri ND, Aslan G, Yalcin CE, Covic A, *et al.* Hypertension as an autoimmune

1 and inflammatory disease. *Hypertens Res.* 2016; **39**(8): 567-73.

2 **Figure Legends:**

3

4 **Figure 1** Comparisons of derived subclass-specific IgG Fc *N*-glycomic composition
5 between hypertension patients and healthy controls.

6 Each box represents the 25th to 75th percentiles. Lines inside the boxes represent the
7 median. Lines outside the boxes represent the 2.5th and 97.5th percentiles.

8 **Figure 2** ROC curve illustrated the performance of the models in diagnosing
9 hypertension of the four ethnic groups.

10 When glycan traits are added into the glycan-based model, the AUCs indicated that
11 the IgG glycan profiles provide more information reinforcing the baseline models
12 utilizing gender, age, BMI and ethnicity. (difference between AUCs, 0.044, 95% CI:
13 0.016 - 0.072, $P= 0.002$).

14 ROC = receiver operating characteristic; AUC = area under the curve

15




Figure 1




Figure 2

1 **Table 1.** Characteristics of the study subjects in pooling of the four ethnic groups

Variables	Hypertension (n=274)	Control group (n=356)	<i>P</i> values
Age (years)	57.07±12.42	49.83±14.88	<0.001*
Gender (male)	126 (50.80%)	122 (40.90 %)	<0.001*
BMI (kg/m ²)	26.76±4.59	24.92±4.71	<0.001*
SBP (mmHg)	151.20±18.44	112.67±11.83	<0.001*
DBP (mmHg)	94.58±12.49	71.72±9.18	<0.001*
TC (mmol/L)	4.72±1.11	4.41±1.20	0.004*
TG (mmol/L)	2.82±0.99	2.84±1.37	0.989
HDL (mmol/L)	1.83±0.75	1.61±0.70	<0.001*
LDL (mmol/L)	2.72±1.62	2.29±1.10	<0.001*
FBG (mmol/L)	5.90±1.40	5.40±1.31	<0.001*
Dyslipidemia	206 (75.18%)	289 (81.12%)	0.069

2 Data are shown as mean ± standard deviation.

3 BMI: body mass index; SBP: systolic blood pressure; DBP: diastolic blood pressure; TC: Total
4 cholesterol; TG: Triglycerides; HDL: high-density lipoprotein cholesterol; LDL: low-density
5 lipoprotein cholesterol; FBG: fasting blood glucose.

6 Mann-Whitney U test: Age, BMI, SBP, DBP, TC, TG, HDL, LDL, FBG, Dyslipidemia; χ^2 test:
7 Gender.

8 * $P < 0.05$ was considered statistically significant.

1 **Table 2.** Directly measured subclass-specific IgG Fc *N*-glycomic composition in the hypertension
 2 patients and the healthy controls for the pooling of the four ethnic groups.

Glycans	Median (IQR)		AOR (95% CI)	<i>P</i> -adjusted	FDR
	Hypertension	Control group			
IgG1G0F	25.79 (21.39,30.29)	23.20 (19.69,27.92)	1.05 (1.02,1.08)	5.00E-04	7.50E-03*
IgG2G0F	32.96 (27.35,39.22)	30.62 (25.11,35.99)	1.05 (1.02,1.07)	2.00E-04	1.20E-02*
IgG2G1FN	4.67 (3.71,5.47)	4.83 (3.92,5.66)	0.80 (0.69,0.92)	2.50E-03	2.14E-02*
IgG2G1FS	3.64 (3.07,4.28)	3.99 (3.22,4.52)	0.74 (0.61,0.90)	2.40E-03	2.40E-02*
IgG2G2S	0.42 (0.25,0.62)	0.45 (0.23,0.78)	0.44 (0.25-0.79)	6.00E-03	4.00E-02*
IgG4G0F	17.53 (12.12,24.14)	14.66 (10.81,20.97)	1.05 (1.02,1.07)	3.00E-04	6.00E-03*
IgG4G1FS	2.36 (1.77,2.92)	2.42 (1.67,3.07)	0.73 (0.59,0.91)	5.20E-03	3.90E-02*
IgG4G1S	0.33 (0.25,0.40)	0.35 (0.24,0.44)	0.10 (0.02,0.42)	1.70E-03	2.04E-02*
IgG4G2FS	5.40 (3.89,7.64)	6.35 (4.45,8.55)	0.89 (0.83,0.95)	2.00E-04	1.20E-02*
IgG4G2N	1.74 (1.09,2.29)	1.77 (1.23,2.46)	0.79 (0.66,0.94)	7.50E-03	4.50E-02*

3 False discovery rate (FDR) was controlled using Benjamini–Hochberg procedure.

4 Only the directly measured subclass-specific IgG Fc *N*-glycan traits which presented significant
 5 differences describing glycomic composition are shown. The other directly measured *N*-glycan
 6 traits are available in Supplementary Table S2.

7 Logistic regression controlled for gender, age, BMI, dyslipidemia, FBG and ethnicity

8 * *P*<0.05 was considered statistically significant.

1 **Table 3.** Derived subclass-specific IgG Fc *N*-glycomic composition in the hypertension patients and
 2 healthy controls for the pooling of the four ethnic groups.

Glycans	Median (IQR)		AOR (95% CI)	<i>P</i> -adjusted	FDR
	Hypertension	Control group			
A1	37.04 (31.81,43.13)	33.76 (27.55,39.74)	1.04 (1.02,1.06)	1.30E-03	6.07E-03*
M1	37.50 (35.63,38.86)	37.93 (36.17,39.37)	0.93 (0.87,0.99)	1.50E-02	3.50E-02*
D1	12.38 (10.37,15.20)	14.23 (11.35,17.79)	0.95 (0.90,0.99)	2.14E-02	4.28E-02*
G1	50.28 (46.67,53.90)	52.74 (48.48,56.46)	0.96 (0.93,0.99)	6.00E-03	1.87E-02*
S1	11.58 (9.23,13.86)	12.20 (9.69,14.92)	0.93 (0.89,0.98)	1.04E-02	2.91E-02*
A2	42.59 (36.51,49.42)	39.25 (32.65,45.60)	1.04 (1.02,1.06)	5.00E-04	4.67E-03*
M2	33.01 (31.01,35.18)	33.87 (31.95,35.35)	0.91 (0.86,0.96)	1.00E-03	5.60E-03*
D2	11.81 (9.68,15.11)	13.82 (10.72,17.12)	0.93 (0.89,0.97)	2.30E-03	9.20E-03*
G2	45.39 (41.10,50.02)	47.90 (43.56,52.31)	0.95 (0.92,0.98)	4.00E-04	5.60E-03*
S2	10.29 (8.05,12.83)	11.54 (8.74,14.43)	0.93 (0.89,0.98)	5.10E-03	1.79E-02*
A4	24.30 (17.77,31.96)	20.26 (15.26,27.63)	1.03 (1.01,1.05)	8.00E-04	5.60E-03*
M4	26.89 (23.81,29.82)	27.76 (24.02,30.96)	0.96 (0.93,0.99)	1.88E-02	4.05E-02*
D4	22.10 (18.15,25.35)	23.82 (20.28,27.80)	0.96 (0.93,0.99)	1.25E-02	3.18E-02*
G4	48.17 (45.17,52.74)	51.39 (47.15,56.26)	0.95 (0.92,0.97)	1.00E-04	2.80E-03*

3 False discovery rate (FDR) was controlled using Benjamini–Hochberg procedure.

4 A, agalactosylation; M, monogalactosylation; D, digalactosylation; G, galactosylation; S, sialylation;

5 B, bisecting GlcNAc; F, fucosylation. Number represents specific IgG subclass.

6 Only the derived subclass-specific IgG Fc *N*-glycan traits which presented significant differences

7 describing glycomic composition are shown. The other derived *N*-glycan traits are available in

8 Supplementary Table S3

9 Logistic regression controlled for gender, age, BMI, dyslipidemia, FBG, and ethnicity

10 * *P*<0.05 was considered statistically significant.

1 **Table 4.** The glycan-based model included the significantly differed subclass-specific IgG Fc
 2 glycopeptide *N*-glycans together with gender, age, BMI and ethnicity

Variables	β	S.E.	Wald χ^2	AOR	<i>P</i> values	95% CI
Gender	-0.37	0.18	4.12	0.69	4.24E-02	(0.48,0.99)
Age	0.02	0.01	11.85	1.03	5.77E-04	(1.01,1.04)
BMI	0.09	0.02	22.59	1.10	2.00E-06	(1.06,1.14)
Ethnicity	-0.30	0.09	10.67	0.74	1.09E-03	(0.62,0.89)
IgG1G0F	0.01	0.02	0.44	1.01	5.07E-01	(0.98,1.05)
IgG2G0F	0.00	0.02	0.02	1.00	8.87E-01	(0.97,1.04)
IgG4G0F	0.02	0.02	2.29	1.02	1.30E-01	(0.99,1.06)
IgG4G2FS	-0.12	0.04	9.56	0.88	1.99E-03	(0.82,0.96)
IgG4G2N	-0.20	0.11	3.54	0.82	6.00E-02	(0.67,1.01)

3 AOR = adjusted odds ratio; 95%CI = 95% confidence interval.

Table S1. Characteristics of the study subjects in the four ethnic groups: Uyгур, Kazak, Kirgiz and Tajik

Variables	UIG			KZK			Hypertension (n=90)
	Hypertension (n=90)	Control group (n=80)	<i>P</i> values	Hypertension (n=52)	Control group (n=98)	<i>P</i> values	
Age (years)	62.88±11.62	59.19±13.83	0.061	49.33±10.48	41.68±10.97	<0.001*	54.85±12.92
Gender (male)	25(27.8%)	40(50.0%)	0.338	25(48.1%)	20(20.4%)	<0.001*	38(42.2%)
BMI (kg/m ²)	25.41±4.95	24.71±4.76	0.349	25.83±4.08	22.68±4.38	<0.001*	28.97±3.90
SBP (mmHg)	151.09±18.44	115.75±8.83	<0.001*	149.46±15.08	105.92±12.67	<0.001*	156.37±22.93
DBP (mmHg)	96.39±12.75	73.44±6.14	<0.001*	95.58±9.18	68.27±9.97	<0.001*	92.37±15.99
TC (mmol/L)	4.88±1.29	4.77±0.95	0.54	4.42±0.76	3.35±0.96	0.004*	4.55±1.08
TG (mmol/L)	1.84±0.46	1.89±0.40	0.518	1.88±0.71	1.15±0.53	0.989	1.70±0.44
HDL (mmol/L)	3.61±2.21	3.09±0.77	0.049	2.75±0.78	2.19±0.93	<0.001*	2.72±0.81
LDL (mmol/L)	2.90±0.73	3.32±1.48	0.020*	1.85±0.50	1.59±0.79	<0.001*	2.72±0.83
FBG (mmol/L)	6.23±1.41	6.11±1.02	0.542	5.37±1.10	4.56±1.00	<0.001*	5.83±1.48

Data are shown as mean ± standard deviation.

BMI: body mass index; SBP: systolic blood pressure; DBP: diastolic blood pressure; TC: Total cholesterol; TG: Total triglyceride; Mann-Whitney U test: Age, BMI, SBP, DBP, TC, TG, HDL, LDL, FBG; χ^2 test: Gender.

* $P < 0.05$ was considered statistically significant.

KGZ		TJK			
Control group (n=78)	<i>P</i> values	Hypertension (n=65)	Control group (n=77)	<i>P</i> values	
40.96±12.12	<0.001*	54.85±12.92	40.96±12.12	<0.001*	
27 (34.6%)	0.091	38(56.7%)	34(33.7%)	0.091	
27.53±3.70	0.026	28.97±3.90	27.53±3.70	0.026	
117.79±9.43	<0.001*	156.37±22.93	117.79±9.43	<0.001*	
72.84±10.08	<0.001*	92.37±15.99	72.84±10.08	<0.001*	
4.79±1.23	0.223	4.56±1.12	4.74±1.28	0.279	
1.68±0.41	0.805	2.74±0.84	2.78±1.25	0.834	
2.93±1.04	0.180	1.71±0.52	1.72±0.45	0.979	
2.75±1.20	0.864	2.72±0.81	2.93±1.04	0.180	
5.44±1.50	0.128	5.80±1.50	5.40±1.54	0.124	

‡: Triglycerides; HDL: high-density lipoprotein cholesterol; LDL: low-density lipoprotein cholesterol;

FBG: fasting blood glucose.

40.96 12.12 40.96±12.12
27.53 3.70 27.53±3.7
117.79 9.43 117.79±9.43
72.84 10.08 72.84±10.08
4.74 1.28 4.74±1.28
1.72 0.45 1.72±0.45
2.93 1.04 2.93±1.04
2.78 1.25 2.78±1.25
5.40 1.54 5.4±1.54

117.79	72.84	4.7391	1.7168	2.9322	2.7762	5.4038
9.433	10.081	1.27868	.44732	1.03752	1.24591	1.53688

Variables	UIG		KZK		KGZ		T
	Hypertensio n (n=90)	Controls (n=80)	Hypertensi on (n=52)	Controls (n=98)	Hypertensio n (n=90)	Controls (n=78)	
Age (years)	62.88±11.62	59.19±13.83	49.33±10.48	41.68±10.97	54.85±12.92	40.96±12.12	54.85±12.92
Gender (male)	25(27.8%)	40(50.0%)	25(48.1%)	20(20.4%)	38(42.2%)	27 (34.6%)	38(56.7%)

JK	
Controls (n=77)	
40.96±12.12	
34(33.7%)	

Table S2. Directly measured subclass-specific IgG Fc *N*-glycomic composition in hypertension p

Glycans	UIG			KZK		
	Mann-Whitney test	Logistic regression		Mann-Whitney test	Logistic regression	
	Z	<i>P</i> values	<i>P</i> -adjusted	Z	<i>P</i> values	<i>P</i> -adjusted
IgG1G0	-0.71	4.80E-01	1.91E-01	-0.19	8.50E-01	1.45E-01
IgG1G0F	-2.28	2.24E-02	9.77E-02	-4.77	1.84E-06	2.90E-03
IgG1G0FN	-1.53	1.25E-01	1.53E-01	-0.99	3.22E-01	6.54E-01
IgG1G0N	-0.64	5.23E-01	2.28E-01	-3.02	2.49E-03	1.92E-02
IgG1G1	-1.88	6.01E-02	6.69E-02	-0.35	7.25E-01	3.01E-01
IgG1G1F	-0.42	6.73E-01	6.53E-01	-2.91	3.56E-03	1.94E-02
IgG1G1FN	-0.21	8.31E-01	9.07E-01	-3.94	8.11E-05	1.09E-02
IgG1G1FNS	-0.18	8.57E-01	4.08E-01	-5.34	9.14E-08	2.00E-04
IgG1G1FS	-1.40	1.60E-01	1.42E-01	-1.90	5.70E-02	7.16E-01
IgG1G1N	-0.22	8.29E-01	1.58E-01	-1.66	9.64E-02	7.16E-02
IgG1G1NS	-1.23	2.20E-01	4.74E-01	-2.00	4.53E-02	3.16E-02
IgG1G1S	-1.55	1.21E-01	5.81E-01	-1.71	8.73E-02	7.99E-02
IgG1G2	-1.44	1.51E-01	1.33E-01	-2.54	1.10E-02	5.79E-02
IgG1G2F	-0.87	3.83E-01	6.96E-01	-4.42	9.74E-06	3.00E-03
IgG1G2FN	-1.56	1.19E-01	5.16E-01	-2.19	2.87E-02	1.96E-01
IgG1G2FNS	-0.36	7.21E-01	5.38E-01	-2.74	6.06E-03	1.60E-03
IgG1G2FS	-0.96	3.39E-01	5.54E-01	-2.78	5.43E-03	1.21E-01
IgG1G2N	0.00	1.00E+00	2.16E-01	-1.26	2.09E-01	4.23E-01
IgG1G2NS	0.00	1.00E+00	7.02E-01	-1.54	1.23E-01	1.46E-02
IgG1G2S	-1.99	4.61E-02	1.25E-01	-1.50	1.33E-01	3.94E-01
IgG2G0	-1.23	2.20E-01	1.11E-01	-2.01	4.40E-02	6.91E-02
IgG2G0F	-0.09	9.28E-01	8.80E-01	-5.89	3.91E-09	<.0001
IgG2G0FN	-0.33	7.45E-01	6.34E-01	-1.82	6.93E-02	1.85E-01
IgG2G0N	-0.70	4.86E-01	1.51E-01	-3.96	7.59E-05	3.70E-03
IgG2G1	-1.54	1.25E-01	8.18E-01	-0.91	3.62E-01	4.51E-01
IgG2G1F	-0.54	5.88E-01	4.15E-01	-3.43	6.00E-04	2.30E-03
IgG2G1FN	-0.35	7.25E-01	6.65E-01	-6.52	7.23E-11	<.0001
IgG2G1FNS	-1.11	2.66E-01	1.52E-01	-4.95	7.49E-07	<.0001
IgG2G1FS	-0.27	7.84E-01	7.85E-01	-3.72	2.02E-04	1.00E-03
IgG2G1N	-0.29	7.70E-01	9.62E-01	-0.19	8.50E-01	2.59E-01
IgG2G1NS	0.00	1.00E+00	7.67E-01	-0.25	7.99E-01	6.72E-01
IgG2G1S	-1.35	1.76E-01	1.59E-01	-2.51	1.20E-02	3.48E-02
IgG2G2	-0.49	6.28E-01	3.74E-01	-4.48	7.53E-06	1.23E-02
IgG2G2F	-0.01	9.92E-01	7.76E-01	-4.70	2.56E-06	2.00E-04
IgG2G2FN	-0.10	9.21E-01	7.62E-01	-4.11	3.94E-05	7.00E-04
IgG2G2FNS	-0.14	8.88E-01	9.34E-01	-0.59	5.54E-01	2.74E-01
IgG2G2FS	-0.20	8.44E-01	8.18E-01	-3.05	2.27E-03	1.49E-02
IgG2G2N	-1.34	1.79E-01	7.50E-01	-4.94	7.96E-07	5.00E-04
IgG2G2NS	0.00	1.00E+00	8.84E-01	-1.33	1.82E-01	1.35E-02
IgG2G2S	-0.03	9.73E-01	1.22E-01	-1.63	1.03E-01	7.65E-01
IgG4G0	-0.67	5.05E-01	1.72E-01	-2.98	2.87E-03	5.09E-01
IgG4G0F	-0.17	8.65E-01	8.49E-01	-7.34	2.18E-13	<0.0001
IgG4G0FN	-0.29	7.72E-01	6.19E-01	-5.75	8.73E-09	<.0001
IgG4G0N	-0.81	4.15E-01	3.01E-01	-2.97	3.02E-03	1.09E-01

IgG4G1	-1.43	1.51E-01	3.02E-01	-4.60	4.13E-06	<.0001
IgG4G1F	-1.11	2.65E-01	3.03E-01	-0.83	4.05E-01	3.42E-01
IgG4G1FN	-0.52	6.05E-01	5.13E-01	-1.55	1.22E-01	2.54E-01
IgG4G1FNS	-0.17	8.63E-01	7.64E-01	-0.57	5.67E-01	3.59E-01
IgG4G1FS	-0.38	7.02E-01	1.42E-01	-0.40	6.87E-01	9.60E-01
IgG4G1N	-1.06	2.89E-01	3.19E-01	-4.94	7.80E-07	<.0001
IgG4G1NS	-0.95	3.40E-01	4.43E-01	-1.57	1.16E-01	2.90E-01
IgG4G1S	-0.93	3.54E-01	5.35E-02	-0.78	4.34E-01	8.20E-03
IgG4G2	-0.24	8.09E-01	6.68E-01	-4.00	6.33E-05	1.80E-03
IgG4G2F	-1.85	6.48E-02	2.00E-01	-2.72	6.43E-03	2.88E-02
IgG4G2FN	-1.94	5.28E-02	2.67E-01	-1.61	1.07E-01	4.29E-01
IgG4G2FNS	-2.50	1.23E-02	3.20E-01	-3.04	2.33E-03	1.39E-01
IgG4G2FS	-2.01	4.46E-02	5.72E-02	-4.02	5.82E-05	1.00E-03
IgG4G2N	-0.87	3.83E-01	4.52E-01	-7.17	7.42E-13	<0.0001
IgG4G2NS	-1.16	2.48E-01	6.47E-01	-2.32	2.02E-02	1.13E-01
IgG4G2S	-0.20	8.44E-01	2.58E-01	-1.15	2.49E-01	5.04E-01

False discovery rate (FDR) was controlled using Benjamini–Hochberg procedure.

Bold in *P*- adjusted represent $P < 0.05$.

Logistic regression controlled for gender, age, BMI, dyslipidemia, FBG and ethnicity

patients and healthy controls for each of the four ethnic groups and the pooling of the four ethn

KGZ			TJK		
Mann-Whitney test		Logistic regression	Mann-Whitney test		Logistic regression
Z	P values	P-adjusted	Z	P values	P-adjusted
-1.08	2.79E-01	3.81E-01	-1.91	5.61E-02	4.73E-01
-0.88	3.80E-01	8.61E-01	-3.19	1.40E-03	7.78E-01
-0.74	4.57E-01	6.09E-01	-3.47	5.28E-04	8.90E-01
-1.16	2.45E-01	3.42E-01	-2.15	3.19E-02	3.02E-01
-0.07	9.43E-01	5.32E-01	-0.53	5.99E-01	3.95E-01
-0.13	8.96E-01	7.12E-01	-0.71	4.75E-01	5.89E-01
-0.31	7.60E-01	6.98E-01	-0.13	8.98E-01	2.99E-01
-0.37	7.08E-01	6.76E-01	-0.63	5.30E-01	7.11E-01
-0.45	6.51E-01	5.21E-01	-0.85	3.97E-01	7.67E-01
-0.39	6.96E-01	4.37E-01	-0.11	9.13E-01	2.51E-01
-1.36	1.72E-01	2.92E-01	0.00	1.00E+00	1.37E-01
-0.43	6.67E-01	5.37E-01	-2.40	1.64E-02	1.96E-01
-0.46	6.44E-01	7.53E-01	-1.97	4.89E-02	9.02E-01
-0.86	3.92E-01	8.06E-01	-3.38	7.34E-04	6.31E-01
-0.88	3.77E-01	9.71E-01	-2.12	3.43E-02	5.61E-01
-0.72	4.73E-01	7.59E-01	-1.94	5.29E-02	9.52E-01
-0.63	5.26E-01	3.27E-01	-3.65	2.58E-04	9.34E-01
-0.34	7.31E-01	8.12E-01	-0.92	3.58E-01	6.63E-01
-0.22	8.29E-01	8.33E-01	0.00	1.00E+00	5.03E-01
-0.48	6.28E-01	4.19E-01	-2.12	3.40E-02	9.20E-01
-0.08	9.33E-01	2.97E-01	-0.04	9.66E-01	4.58E-01
-1.04	2.99E-01	5.14E-01	-3.21	1.35E-03	8.59E-01
-1.40	1.63E-01	1.77E-01	-2.89	3.80E-03	8.09E-01
-0.10	9.20E-01	5.94E-01	-0.07	9.42E-01	5.43E-01
-0.26	7.98E-01	7.70E-01	-0.46	6.44E-01	2.25E-01
-0.51	6.11E-01	3.48E-01	-1.46	1.45E-01	4.24E-01
-0.67	5.02E-01	5.10E-01	-1.00	3.17E-01	9.57E-01
-0.66	5.08E-01	7.10E-01	-0.19	8.46E-01	9.17E-01
-1.60	1.11E-01	1.70E-02	-1.88	5.97E-02	9.82E-01
-0.67	5.02E-01	7.39E-01	-0.70	4.86E-01	2.71E-01
-0.99	3.24E-01	5.81E-01	0.00	1.00E+00	5.34E-01
-1.68	9.24E-02	7.86E-02	-0.68	4.97E-01	6.35E-01
-0.09	9.25E-01	7.49E-01	-1.48	1.38E-01	2.95E-01
-1.37	1.71E-01	2.78E-01	-2.94	3.24E-03	3.87E-01
-1.26	2.08E-01	2.28E-01	-1.80	7.21E-02	8.81E-01
-1.43	1.52E-01	2.71E-01	-0.81	4.16E-01	8.57E-01
-1.52	1.28E-01	9.91E-02	-3.26	1.12E-03	7.99E-01
-0.22	8.25E-01	7.08E-01	-0.66	5.08E-01	3.16E-01
-0.82	4.10E-01	5.18E-01	0.00	1.00E+00	1.77E-01
-1.39	1.64E-01	2.20E-01	-3.00	2.74E-03	6.84E-01
-1.21	2.26E-01	3.60E-01	-1.57	1.17E-01	6.56E-01
-0.57	5.72E-01	9.69E-01	-2.45	1.44E-02	7.34E-01
-0.04	9.65E-01	7.18E-01	-3.09	2.01E-03	8.91E-01
-0.99	3.24E-01	3.01E-01	-1.09	2.76E-01	2.93E-01

-0.32	7.53E-01	9.37E-01	-2.10	3.57E-02	6.72E-01
-0.45	6.51E-01	9.73E-01	-0.02	9.82E-01	4.80E-01
-0.24	8.10E-01	9.98E-01	-1.17	2.43E-01	8.69E-01
-1.30	1.94E-01	3.99E-01	-0.48	6.34E-01	8.04E-01
-0.83	4.06E-01	2.02E-01	-1.14	2.55E-01	2.33E-01
-0.43	6.64E-01	9.44E-01	-2.53	1.14E-02	5.84E-01
-1.13	2.58E-01	5.43E-01	-1.54	1.22E-01	6.62E-02
-0.78	4.35E-01	6.13E-01	-0.11	9.10E-01	9.78E-01
-0.19	8.49E-01	4.58E-01	-0.93	3.50E-01	4.83E-01
-0.07	9.40E-01	8.82E-01	-2.65	8.08E-03	6.42E-01
-0.84	3.99E-01	7.07E-01	-1.32	1.87E-01	9.21E-01
-1.16	2.47E-01	2.81E-01	-2.48	1.30E-02	9.83E-01
-0.45	6.51E-01	3.93E-01	-2.67	7.51E-03	5.42E-01
-0.73	4.63E-01	8.06E-01	-1.26	2.08E-01	8.39E-01
-1.90	5.69E-02	3.88E-01	0.00	1.00E+00	7.36E-01
-0.97	3.32E-01	5.75E-01	-1.79	7.42E-02	6.22E-01

nic groups.

Mann-Whitney test		Pooling of 4 ethnic groups	
Z	P values	Logistic regression <i>P-adjusted</i>	FDR
-0.19	8.50E-01	4.02E-01	4.72E-01
-4.77	1.84E-06	5.00E-04	7.50E-03
-0.99	3.22E-01	2.72E-01	3.63E-01
-3.02	2.49E-03	8.70E-01	8.70E-01
-0.35	7.25E-01	7.66E-01	8.21E-01
-2.91	3.56E-03	3.05E-01	3.90E-01
-3.94	8.11E-05	4.65E-02	1.21E-01
-5.34	9.14E-08	8.50E-01	8.65E-01
-1.90	5.70E-02	6.46E-02	1.55E-01
-1.66	9.64E-02	4.53E-01	5.13E-01
-2.00	4.53E-02	7.09E-02	1.58E-01
-1.71	8.73E-02	4.70E-02	1.18E-01
-2.54	1.10E-02	4.01E-02	1.20E-01
-4.42	9.74E-06	3.73E-02	1.18E-01
-2.19	2.87E-02	6.17E-02	1.54E-01
-2.74	6.06E-03	8.39E-01	8.83E-01
-2.78	5.43E-03	6.94E-02	1.60E-01
-1.26	2.09E-01	8.92E-02	1.85E-01
-1.54	1.23E-01	4.26E-01	4.91E-01
-1.50	1.33E-01	3.41E-02	1.14E-01
-2.01	4.40E-02	3.33E-01	4.00E-01
-5.89	3.91E-09	2.00E-04	1.20E-02
-1.82	6.93E-02	2.73E-01	3.55E-01
-3.96	7.59E-05	8.44E-01	8.73E-01
-0.91	3.62E-01	1.42E-01	2.44E-01
-3.43	6.00E-04	3.18E-01	3.97E-01
-6.52	7.23E-11	2.50E-03	2.14E-02
-4.95	7.49E-07	9.36E-02	1.81E-01
-3.72	2.02E-04	2.40E-03	2.40E-02
-0.19	8.50E-01	2.31E-01	3.15E-01
-0.25	7.99E-01	4.61E-01	5.13E-01
-2.51	1.20E-02	1.24E-02	6.76E-02
-4.48	7.53E-06	2.26E-02	1.04E-01
-4.70	2.56E-06	2.98E-02	1.12E-01
-4.11	3.94E-05	2.75E-02	1.18E-01
-0.59	5.54E-01	1.70E-01	2.69E-01
-3.05	2.27E-03	2.15E-02	9.92E-02
-4.94	7.96E-07	1.41E-01	2.49E-01
-1.33	1.82E-01	4.30E-02	1.23E-01
-1.63	1.03E-01	6.00E-03	4.00E-02
-2.98	2.87E-03	1.86E-01	2.87E-01
-7.34	2.18E-13	3.00E-04	6.00E-03
-5.75	8.73E-09	1.59E-01	2.57E-01
-2.97	3.02E-03	2.03E-01	2.98E-01

-4.60	4.13E-06	9.41E-02	1.76E-01
-0.83	4.05E-01	6.98E-01	7.61E-01
-1.55	1.22E-01	8.77E-02	1.88E-01
-0.57	5.67E-01	2.26E-01	3.16E-01
-0.40	6.87E-01	5.20E-03	3.47E-02
-4.94	7.80E-07	2.15E-01	3.07E-01
-1.57	1.16E-01	1.43E-01	2.39E-01
-0.78	4.34E-01	1.70E-03	2.04E-02
-4.00	6.33E-05	3.22E-01	3.94E-01
-2.72	6.43E-03	2.80E-02	1.05E-01
-1.61	1.07E-01	1.10E-01	1.99E-01
-3.04	2.33E-03	3.01E-02	1.00E-01
-4.02	5.82E-05	2.00E-04	1.20E-02
-7.17	7.42E-13	7.50E-03	4.50E-02
-2.32	2.02E-02	1.98E-01	2.96E-01
-1.15	2.49E-01	9.29E-02	1.86E-01

Table S3. Derived subclass-specific IgG Fc N-glycomic composition in hypertension patients and h

Glycans	UIG			KZK		
	Mann-Whitney test		Logistic regression	Mann-Whitney test		Logistic regression
	Z	P values	P-adjusted	Z	P values	P-adjusted
A1	-1.83	6.74E-02	1.53E-01	-4.41	1.03E-05	4.20E-03
M1	-0.73	4.64E-01	5.90E-01	-3.76	1.73E-04	6.41E-02
D1	-1.25	2.10E-01	4.97E-01	-4.32	1.53E-05	3.50E-03
G1	-1.25	2.11E-01	4.62E-01	-4.33	1.48E-05	2.50E-03
S1	-1.82	6.83E-02	1.54E-01	-2.64	8.34E-03	1.81E-01
B1	-0.46	6.49E-01	6.89E-01	-0.56	5.75E-01	7.74E-01
F1	-2.28	2.25E-02	2.88E-02	-1.16	2.44E-01	9.98E-01
A2	-0.01	9.94E-01	8.93E-01	-5.45	4.94E-08	<.0001
M2	-0.50	6.17E-01	4.74E-01	-5.32	1.02E-07	<.0001
D2	-0.46	6.49E-01	8.51E-01	-5.17	2.30E-07	<.0001
G2	-0.18	8.55E-01	8.09E-01	-5.47	4.42E-08	<.0001
S2	-0.34	7.37E-01	7.74E-01	-3.23	1.22E-03	5.80E-03
B2	-0.14	8.90E-01	8.57E-01	-2.03	4.20E-02	1.91E-02
F2	-1.28	2.01E-01	5.14E-01	-0.46	6.47E-01	4.22E-01
A4	-0.28	7.82E-01	7.50E-01	-7.23	4.95E-13	<0.0001
M4	-0.26	7.91E-01	8.61E-01	-4.82	1.45E-06	2.00E-04
D4	-0.05	9.62E-01	9.67E-01	-6.31	2.71E-10	<.0001
G4	-0.17	8.65E-01	8.59E-01	-8.00	1.24E-15	<0.0001
S4	-0.66	5.12E-01	8.18E-01	-1.70	8.95E-02	2.04E-01
B4	-0.05	9.64E-01	8.58E-01	-0.09	9.31E-01	5.68E-01
F4	-0.98	3.29E-01	2.43E-01	-3.72	1.96E-04	2.90E-03

False discovery rate (FDR) was controlled using Benjamini–Hochberg procedure.

Bold in P- adjusted represent $P < 0.05$.

Logistic regression controlled for gender, age, BMI, dyslipidemia, FBG and ethnicity

A - agalactosylation, M - monogalactosylation, D- digalactosylation, G - galactosylation (total), S - s

healthy controls for each of the four ethnic groups and the pooling of the four ethnic groups.

KGZ			TJK			
Mann-Whitney test		Logistic regression	Mann-Whitney test	Logistic regression	Mann-Whitney test	
Z	P values	<i>P-adjusted</i>	Z	P values	<i>P-adjusted</i>	Z
-1.02	3.09E-01	6.21E-01	-3.64	2.70E-04	9.44E-01	-4.41
-0.38	7.03E-01	7.79E-01	-0.30	7.67E-01	4.22E-01	-3.76
-0.88	3.77E-01	7.99E-01	-3.34	8.34E-04	7.48E-01	-4.32
-0.72	4.71E-01	9.89E-01	-3.22	1.27E-03	8.97E-01	-4.33
-0.96	3.38E-01	2.42E-01	-3.72	1.96E-04	9.33E-01	-2.64
-0.55	5.83E-01	5.50E-01	-1.86	6.24E-02	4.24E-01	-0.56
0.00	1.00E+00	5.64E-01	-0.27	7.91E-01	4.05E-01	-1.16
-1.12	2.62E-01	3.25E-01	-3.59	3.34E-04	8.90E-01	-5.45
-0.27	7.91E-01	4.06E-01	-1.01	3.14E-01	7.36E-01	-5.32
-1.30	1.92E-01	3.59E-01	-3.44	5.80E-04	8.45E-01	-5.17
-1.35	1.75E-01	3.16E-01	-2.84	4.48E-03	9.67E-01	-5.47
-1.76	7.88E-02	4.28E-02	-3.00	2.66E-03	9.02E-01	-3.23
0.00	1.00E+00	6.24E-01	-1.23	2.18E-01	7.71E-01	-2.03
-0.76	4.44E-01	1.85E-01	-0.34	7.33E-01	1.49E-01	-0.46
-0.45	6.51E-01	9.73E-01	-2.61	9.09E-03	8.10E-01	-7.23
-0.02	9.80E-01	9.44E-01	-1.62	1.06E-01	6.99E-01	-4.82
-0.64	5.24E-01	4.30E-01	-2.25	2.42E-02	5.48E-01	-6.31
-0.20	8.42E-01	5.02E-01	-0.90	3.70E-01	8.07E-01	-8.00
-0.32	7.53E-01	8.97E-01	-3.13	1.74E-03	8.07E-01	-1.70
-0.68	4.98E-01	7.22E-01	-2.69	7.14E-03	7.92E-01	-0.09
-0.02	9.88E-01	7.90E-01	-0.44	6.58E-01	5.70E-01	-3.72

glycosylation, B - bisecting GlcNAc, F - fucosylation. Number represents specific IgG subclass.

Pooling of 4 ethnic groups		
Witney test	Logistic regression	
<i>P</i> values	<i>P-adjusted</i>	<i>FDR</i>
1.03E-05	1.30E-03	6.07E-03
1.73E-04	1.50E-02	3.50E-02
1.53E-05	2.14E-02	4.28E-02
1.48E-05	6.00E-03	1.87E-02
8.34E-03	1.04E-02	2.91E-02
5.75E-01	4.74E-01	6.98E-01
2.44E-01	3.44E-01	5.35E-01
4.94E-08	5.00E-04	4.67E-03
1.02E-07	1.00E-03	5.60E-03
2.30E-07	2.30E-03	9.20E-03
4.42E-08	4.00E-04	5.60E-03
1.22E-03	5.10E-03	1.79E-02
4.20E-02	1.25E-01	2.32E-01
6.47E-01	2.29E-01	3.76E-01
4.95E-13	8.00E-04	5.60E-03
1.45E-06	1.88E-02	4.05E-02
2.71E-10	1.25E-02	3.18E-02
1.24E-15	1.00E-04	2.80E-03
8.95E-02	8.42E-01	1.12E+00
9.31E-01	2.14E-01	3.74E-01
1.96E-04	6.83E-01	9.56E-01