

3-13-2020

## Comparison between high- and low-intensity eccentric cycling of equal mechanical work for muscle damage and the repeated bout effect

Georgios Mavropalias  
*Edith Cowan University*

Tomoko Koeda

Oliver R. Barley  
*Edith Cowan University*

Wayne C. K. Poon  
*Edith Cowan University*

Aiden J. Fisher

*See next page for additional authors*

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworkspost2013>



Part of the [Medicine and Health Sciences Commons](#)

---

[10.1007/s00421-020-04341-5](https://doi.org/10.1007/s00421-020-04341-5)

This is a post-peer-review, pre-copyedit version of an article published in *European Journal of Applied Physiology*. Mavropalias, G., Koeda, T., Barley, O. R., Poon, W. C., Fisher, A. J., Blazevich, A. J., & Nosaka, K. (2020). Comparison between high-and low-intensity eccentric cycling of equal mechanical work for muscle damage and the repeated bout effect. *European Journal of Applied Physiology*, 120, 1015-1025.

<https://doi.org/10.1007/s00421-020-04341-5>

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/8125>

---

## Authors

Georgios Mavropalias, Tomoko Koeda, Oliver R. Barley, Wayne C. K. Poon, Aiden J. Fisher, Anthony J. Blazeovich, and Kazunori Nosaka

1  
2  
3 **COMPARISON BETWEEN HIGH- AND LOW-INTENSITY**  
4 **ECCENTRIC CYCLING OF EQUAL MECHANICAL WORK FOR**  
5 **MUSCLE DAMAGE AND THE REPEATED BOUT EFFECT**  
6

7 Georgios Mavropalias<sup>1</sup>, Tomoko Koeda<sup>2</sup>, Oliver R. Barley<sup>1</sup>, Wayne C.K. Poon<sup>1</sup>,  
8 Aiden J. Fisher<sup>3</sup>, Anthony J. Blazevich<sup>1</sup>, Kazunori Nosaka<sup>1</sup>  
9

10 <sup>1</sup>School of Medical and Health Sciences, Edith Cowan University, Australia

11 <sup>2</sup>Faculty of Rehabilitation Sciences, Nagoya Gakuin University, Japan

12 <sup>3</sup>School of Science, Edith Cowan University, Australia  
13

14 Running head: High- vs low-intensity eccentric cycling and muscle damage  
15  
16

17 \*Corresponding author:

18 **Georgios Mavropalias**

19 School of Medical and Health Sciences

20 Edith Cowan University

21 270 Joondalup Drive, Joondalup, WA, 6027, Australia

22 Email: georgios.mavropalias@gmail.com

23 Phone: +61 411 89 69 02

24 ORCID: 0000-0001-7753-5693  
25

## ABSTRACT

**Purpose:** We compared high- and low-intensity eccentric cycling (ECC) with the same mechanical work for changes in muscle function and muscle soreness, and examined the changes after subsequent high-intensity ECC.

**Methods:** Twenty men performed either high-intensity ECC (1 min x 5 at 20% of peak power output: PPO) for two bouts separated by 2 weeks (H-H, n=11), or low-intensity (4 min x 5 at 5% PPO) for the first and high-intensity ECC for the second bout (L-H, n=9). Changes in indirect muscle damage markers were compared between groups and bouts.

**Results:** At 24 h after the first bout, both groups showed similar decreases in maximal isometric (70° knee angle,  $-10.6 \pm 11.8\%$ ) and isokinetic ( $-11.0 \pm 8.2\%$ ) contraction torque of the knee extensors (KE), squat ( $-7.7 \pm 10.4\%$ ) and counter-movement jump ( $-5.9 \pm 8.4\%$ ) heights ( $P < 0.05$ ). Changes in KE torque and jump height were smaller after the second than first bout for both groups ( $P < 0.05$ ). Increases in plasma creatine kinase activity were small, and no significant changes in vastus lateralis or intermedius thickness nor ultrasound echo-intensity were observed. KE soreness with palpation was greater ( $P < 0.01$ ) in H-H (peak:  $4.2 \pm 1.0$ ) than L-H ( $1.4 \pm 0.6$ ) after the first bout, but greater in L-H ( $3.6 \pm 0.9$ ) than H-H ( $1.5 \pm 0.5$ ) after the second bout. This was also found for muscle soreness with squat, KE stretch and gluteal palpation.

**Conclusion:** The high- and low-intensity ECC with matched mechanical work induced similar decreases in muscle function, but DOMS was greater after high-intensity ECC, which may be due to greater extracellular matrix damage and inflammation.

**Keywords:** lengthening muscle action; maximal voluntary isometric torque; squat jump; countermovement jump; delayed onset muscle soreness; creatine kinase activity; rate of force development

52

## ABBREVIATIONS

53

CK	Creatine kinase
CMJ	Counter-movement jump
DOMS	Delayed-onset muscle soreness
H-H	High-High group
L-H	Low-High group
MVCC	Maximal voluntary concentric contraction
MVIC	Maximal voluntary isometric contraction
PPO	Peak power output
RBE	Repeated bout effect
RFD	Rate of force development
SJ	Squat jump

54

55

## INTRODUCTION

Lengthening muscle actions (eccentric actions) are performed when activated muscles are lengthened by an external force. It has been well documented that exercises consisting of unaccustomed eccentric actions induce muscle damage (Clarkson et al. 1992). Eccentric exercise-induced muscle damage may be indicated by histological alterations in myofilament structure such as Z-line streaming (Fridén et al. 1983; Armstrong 1990) and associated with delayed-onset muscle soreness (DOMS), prolonged decreases in muscle function, increased ultrasound echo-intensity, muscle swelling and increases in intramuscular proteins in the blood such as creatine kinase (CK) and myoglobin (Clarkson et al. 1992; Nosaka and Clarkson 1995). Repeating the same exercise within several weeks after the initial bout results in smaller changes in these markers when compared with the initial bout, which is referred to as the repeated bout effect (RBE) (Nosaka and Clarkson 1995; Hortobágyi et al. 1998; McHugh 2003).

While many factors affect the magnitude of muscle damage and RBE induced by eccentric exercise, exercise intensity appears to be a key factor; i.e., higher intensity eccentric actions induce greater damage than lower intensity ones (Nosaka and Newton 2002; Chen et al. 2007, 2012; Hasenoehrl et al. 2017; Tsuchiya et al. 2019). For example, Nosaka et al. (2002) compared the effects of 30 eccentric actions of the elbow flexors that were performed maximally to those performed at 50% of maximal voluntary isometric contraction (MVIC) torque by young men, and reported that changes in MVIC torque, muscle soreness and plasma CK activity were significantly greater following the maximal exercise. This was also observed in other studies in which maximal and submaximal intensities were compared after elbow flexor eccentric exercise (Chen et al. 2007, 2012; Hasenoehrl et al. 2017; Tsuchiya et al. 2019). However, the exercise mechanical work was not matched between conditions in these studies,

so it is not explicitly clear whether the intensity or the mechanical work was the critical factor influencing the magnitude of damage and DOMS.

To the best of our knowledge only three studies have matched exercise mechanical work to examine the effect of eccentric action intensity on muscle damage. Gauche et al. (2009) compared the effects of 15 eccentric actions of the elbow flexors at 80% of maximal concentric torque to 30 eccentric actions at 40%, and found no difference in MVIC torque changes after the exercises. Similarly, Hill et al. (2019) compared 36 eccentric actions of the elbow flexors at 80% of maximal eccentric torque to 72 eccentric actions at 40%, and also found no difference in the changes in MVIC torque after exercise. In the knee extensors, Paschalis (2005) compared the effects of 120 maximal eccentric actions to a condition where submaximal (50% of maximal eccentric torque) eccentric actions were performed until equal mechanical work was accomplished. The results revealed no difference in DOMS between conditions, but did show significantly greater changes in MVIC torque and plasma CK activity after the maximal intensity exercise. When considered together, the current evidence indicates a lack of difference in muscle damage between higher- and lower-intensity eccentric exercises when exercise intensity is submaximal (Gauche et al. 2009; Hill et al. 2019), but that damage may be greater after work-matched maximal-intensity exercise than work-matched submaximal-intensity exercise (Paschalis et al. 2005). Nonetheless, more studies are required to explicitly test this hypothesis.

It should be noted that the total mechanical work of exercise imposed in previous studies was small when compared to eccentric cycling that involves the completion of a large number of eccentric actions at a relatively low intensity. Although eccentric cycling is generally performed at intensities less than 30% of maximal knee extension capacity (Peñailillo et al. 2015b), it induces moderate muscle soreness and function decreases already by 30 min of cycling in which ~1800 eccentric actions are performed (Yu et al. 2002; Peñailillo et al.

2013). It therefore appears that eccentric cycling is a good exercise model in which to study the effect of eccentric exercise intensity on muscle damage when the exercise bout consists of a larger number of submaximal eccentric actions, such as in hiking, long distance running, and ball sports. It is also ideal that the total mechanical work of exercise ( $\text{workload} \times \text{cadence} \times \text{time}$ ) can be easily matched between high- and low-intensity protocols. Using this paradigm, a recent study reported greater muscle soreness, function loss and plasma CK activity after high-intensity eccentric cycling (193 W) when compared to lower-intensity (112 W) eccentric cycling in older adults (González-Bartholin et al. 2019). However, the study included four eccentric cycling preconditioning (i.e. familiarisation) sessions, and the same participants performed both exercise intensity conditions. Therefore, the muscle damage profiles would have been strongly affected by the repeated bout and/or preconditioning effects (Nosaka and Clarkson 1995; Chen et al. 2012). Consequently, it is of interest to compare the muscle damage responses to high- and low-intensity eccentric cycling without familiarisation or preconditioning, especially given that previous studies showed smaller changes in, and faster recovery of, indirect markers of muscle damage after the second than the first eccentric cycling bout (Peñailillo et al. 2013, 2015b; Valladares-Ide et al. 2019).

Given the above, the present study compared high- and low-intensity eccentric cycling performed with the same total mechanical work for changes in indirect markers of muscle damage and DOMS. It was hypothesised that changes in muscle function and soreness would be similar between the two eccentric cycling conditions. The present study also included a second bout of eccentric cycling that was performed 14 days after the first bout, in which high-intensity eccentric cycling was always performed to determine whether low-intensity eccentric cycling in the first bout could confer a protective effect against damage and soreness from a subsequent high-intensity bout. It was hypothesised that the magnitude of changes in markers



of muscle damage and soreness would be smaller after the second than the first bout regardless of the intensity used in the first bout.

## METHODS

### Participants and study design

The sample size of the present study was estimated using G\*Power (Version 3.1.9.2, Universitat Kiel, Germany) based on the study by Paschalis et al. (2005), who compared high- and low-intensity eccentric knee extensions, showing a greater decrease in MVIC torque after the high- than low-intensity eccentric exercise with an effect size of 1.34. With a power level of 0.8, and an alpha level of 0.05, it was found that a total of 18 participants were required, thus at least 9 participants were recruited for each group. Ethical approval from the Edith Cowan University Human Research Ethics Committee was sought before the study commencement.

All participants were healthy young men who had not performed eccentric exercise in the past 6 months and reported no history of orthopaedic lower limb injuries or any neurological disorders. Each participant signed a written informed consent form and answered medical questionnaires before participating in the study. The participants were randomly allocated to one of two groups, H-H and L-H, based on their peak power output during a maximal 10-s isokinetic concentric cycling bout at 60 rpm (PPO) on a cycle ergometer. The participants in H-H performed the same high-intensity eccentric cycling exercise in two bouts separated by 14 days, whereas L-H performed low-intensity eccentric cycling exercise in the first bout and high-intensity eccentric cycling exercise of equal mechanical work in the second bout at 14 days later. As shown in Table 1, no significant between-group differences were found for any physical or physiological characteristics at baseline.

The dependent variables, in the order of measurement in the study, included plasma CK activity, knee extensor ultrasound echo intensity and muscle thickness, muscle soreness, squat

(SJ) and counter-movement jump (CMJ), peak power output (PPO), knee extensor maximal voluntary isometric contraction torque (MVIC) at 70° and 20° of knee flexion (0° = full extension), rate of force development (RFD) during MVIC at 70°, and knee extensor isokinetic maximal voluntary concentric torque at 270°/s, 180°/s and 90°/s (MVCC<sub>270</sub>, MVCC<sub>180</sub>, MVCC<sub>90</sub>). These were measured before, immediately after, and 24, 48 and 72 h after each exercise bout. In order to warm up for the performance tests, the participants completed a 2-min concentric cycling warm-up at 50 W before the jump tests. Five days before the initial bout, the participants completed a familiarisation session to experience all measurements shown above, but they did not perform the eccentric cycling to avoid a possible protective effect against muscle damage.

### **Eccentric cycling**

Using an eccentric cycle ergometer (Grucox Eccentric Trainer, Grucox, South Africa), participants completed a 2-min concentric cycling warm-up at 50 W after an appropriate seat position was determined by the investigator in relation to their leg length. In the first bout, participants in the H-H performed high-intensity eccentric cycling at 20% PPO for 5 sets of 1 min with a 1-min rest between sets, and those in L-H performed low-intensity eccentric cycling at 5% of PPO for 5 sets of 4 min with a 1-min rest between sets, in order to achieve an equal mechanical work. In all cycling sessions, the cadence was set at 60 rpm. These protocols were chosen based on our pilot studies showing that the same mechanical work was achieved with a distinct difference in intensity in a similar exercise fashion (interval exercise). In the second session that was performed 14 days after the first bout, the same PPO value was used, however H-H repeated the same exercise whereas L-H performed high-intensity eccentric cycling with mechanical work equal to that achieved in the low-intensity eccentric cycling session. This was

accomplished by decreasing the duration of each set from 5 min to 1 min but increasing the intensity from 5% to 20% PPO.

Each participant was given 1 min of cycling familiarisation immediately before both the initial and secondary exercise bouts, during which eccentric cycling was performed from 30 rpm reaching 60 rpm at the end of the minute for a target power output. The participants were instructed to resist the pedals by using the lower limbs to perform smooth backward rotations while maintaining their target power, which was shown on a computer screen set in front of the ergometer. The pedal straps were removed from the ergometer to ensure the participants only pushed against the pedals and did not pull when the pedals were moving away from them to generate power using their knee and hip flexors.

Heart rate during cycling was continuously recorded by a heart rate monitor (Polar S810i, Polar Electro, Finland). Rate of perceived effort was assessed at the end of each set using a modified version of Borg's category-ratio scale (Borg 1998), as it has been reported that measuring effort during eccentric cycling is more accurate than exertion (Peñailillo et al. 2018). At the end of each set, the participants were asked to rate their physical effort required by their knee extensors to maintain the target power.

#### **Maximal voluntary isometric (MVIC), concentric (MVCC) contraction torque and rate of force development (RFD)**

An isokinetic dynamometer (System 3, Biodex Medical Systems, USA) was used for the measurements of MVIC and MVCC and RFD. During measurements, visual feedback of torque was displayed on a computer screen using software connected to the dynamometer (LabVIEW, National Instruments, Australia). MVIC torque was measured from the right knee extensors at knee joint angles of 70° and 20° (MVIC<sub>70</sub> and MVIC<sub>20</sub>, 0° = full knee extension) after several submaximal contractions as warm-up. Three 4-s maximal voluntary isometric

knee extensions were performed at each joint angle, separated by 1 min of passive rest. Participants were advised not to perform any countermovement motion before the muscle action, and were instructed to “exert force as fast and then as hard as possible” (Maffiuletti et al. 2016). The highest torque value was used for further analysis.

RFD has been identified as a more sensitive measure of exercise-induced muscle damage than MVIC (Peñailillo et al. 2015a). RFD at 70° knee angle was assessed in the time periods of 0-100 ms (RFD<sub>0-100</sub>) and 100-200 ms (RFD<sub>100-200</sub>), where the onset of force production was at 0 s. The onset of force production was identified as the point at which torque data exceeded the baseline by > 7.5 Nm (Peñailillo et al. 2015a). RFD was not measured within shorter time intervals because the data were not of sufficient reliability (e.g. Maffiuletti et al., 2016). If the torque decreased more than 5 Nm below baseline immediately before torque onset, a countermovement considered to have occurred, and the trial was not considered for analysis. The contractions with the highest RFD values were selected for further analysis.

Maximal concentric knee extensor torque (MVCC) was also measured on the isokinetic dynamometer at the angular velocities of 270°/s (MVCC<sub>270</sub>), 180°/s (MVCC<sub>180</sub>) and 90°/s (MVCC<sub>90</sub>), in this order. Each contraction started at a knee angle of 90° before extending the knee to 20° before flexing again to 90° as fast and as hard as possible for four consecutive contractions. The three trials (sets) were separated by 1 min of passive rest. The highest torque value was used for further analysis.

### **Concentric cycling sprint peak power output (PPO)**

Participants performed a PPO test on the cycle ergometer as described above. Power data during the sprint were recorded at a frequency of 40 Hz, and the maximal power achieved was calculated from the extracted data from the ergometer.

## **Squat jump (SJ) and counter-movement jump (CMJ)**

SJ and CMJ tests were performed on a force platform (9290AD, Kistler Instruments, Switzerland). SJ was performed from a self-selected knee joint angle, and CMJ was performed from an upright (standing) position. The participant was asked to keep their hands on their hips throughout the jump, to land with their toes first followed by their heel, and to not bend their knees while in air. They were instructed to maintain the same technique for the jump tests, and to jump as high as possible. Each participant performed 3 jumps, and the peak jump height was selected for further analysis.

## **Muscle soreness**

The magnitude of knee extensor soreness was assessed by palpation, stretch and movement of the knee extensors using a custom 0 (no soreness) – 10 (maximal soreness) scale. Each participant laid supine on a massage table and the investigator palpated the proximal, middle and distal points of vastus medialis, vastus lateralis, and rectus femoris while the middle gluteus maximus was palpated with the participant in the prone position. The average values of the nine knee extensor palpation points were used for subsequent statistical analysis. The investigator then brought the right lower limb to full knee flexion to assess the muscle soreness during stretch. Participants were then asked to sit down to and stand up from a chair slowly (~3 s for each direction) for the movement-based assessment of muscle soreness.

## **Muscle ultrasound echo-intensity and thickness**

Ultrasound images of vastus lateralis and intermedius were taken to measure muscle thickness and echo intensity using a real-time B-mode ultrasound apparatus (Aloka SSD-alpha10, Aloka Co., Japan). The ultrasound settings including focus, contrast and gain were kept the same between measurements. An ultrasound probe (60-mm, linear array) was placed on a marked

site at the middle point between the lateral epicondyle and greater trochanter of the right leg, while the participant was lying on a massage table. Three transverse images of each muscle were obtained and then transferred to a computer for the assessment of muscle thickness and mean echo intensity of a greyscale histogram (0: black, 256: white) using an image analysis software (ImageJ, National Institute of Health, USA). The average value of the three images was used for subsequent statistical analysis.

#### **Plasma CK activity**

A venous blood sample was obtained at baseline, and 24, 48 and 72 h post-exercise, before any other testing was conducted. Two 6-mL blood samples were collected in EDTA tubes from an antecubital vein. Immediately after sampling, a 30- $\mu$ L whole blood sample was used for plasma CK activity analysis using a Reflotron (Roche Diagnosis, Germany).

#### **Statistical analyses**

A two-way (group  $\times$  time) repeated measures analysis of variance (ANOVA) was used to compare between groups for changes in the dependent variables over time, and between bouts for each group. The data were assessed for assumptions of normality by analysing the standardized residuals using a Shapiro-Wilk test, and for sphericity by Mauchly's sphericity test. Ordinal data (muscle soreness and perceived effort values) were rank-transformed before analysis (Wobbrock et al. 2011). In the case of a significant interaction effect, a Holm's sequential Bonferroni procedure was performed to identify possible differences between groups for each time point. The significance level was set at  $p \leq 0.05$ . All statistical testing was performed using SPSS v25 (IBM, USA). Data are presented as mean  $\pm$  standard deviation (SD).

## RESULTS

### **Eccentric cycling intensity, mechanical work and effort**

Average eccentric cycling intensity was  $154.0 \pm 25.0$  W for the first and second bouts in H-H, and was  $36.0 \pm 4.9$  W in the first bout and  $143.4 \pm 22.6$  W in the second bout in L-H. The total mechanical work (in joules, calculated by power  $\times$  seconds) performed in the first bout was  $46,200.0 \pm 7,715.3$  J in H-H and  $43,140.0 \pm 5,839.3$  J in L-H without a significant ( $P = 0.340$ ) difference between groups. During the second bout, the total mechanical work for H-H and L-H were  $46,200.0 \pm 7,715.3$  J and  $43,006.7 \pm 6,791.5$  J, respectively, without a significant difference between groups ( $P = 0.322$ ) or between bouts in either group ( $P = 0.944$ ).

Average heart rate was greater in H-H ( $130 \pm 17$  bpm) than L-H ( $108 \pm 19$  bpm) in the first bout ( $P < 0.001$ ), and in L-H ( $134 \pm 18$  bpm) than H-H ( $120 \pm 12$  bpm) in the second bout ( $P < 0.01$ ); a significant bout  $\times$  time interaction effect was observed for heart rate in L-H ( $P < 0.001$ ). Average perceived effort was greater in H-H ( $5.1 \pm 2.2$ ) than L-H ( $3.0 \pm 0.9$ ) in the first bout ( $P < 0.05$ ), and was also greater in L-H ( $6.4 \pm 0.5$ ) than H-H ( $4.2 \pm 0.4$ ) in the second bout ( $P < 0.05$ ); a significant bout  $\times$  time interaction effect was observed for L-H ( $P < 0.001$ ).

### **Muscle function**

As shown in Figure 1, most muscle function measures showed significant changes over time, but no significant differences between H-H and L-H were observed for the changes after the first or second bouts. MVIC<sub>70</sub> torque was reduced immediately ( $-13.5 \pm 8.6\%$ ) and 24 h after the first bout ( $-10.6 \pm 11.8\%$ ) for both groups, and this was similar after the second bout ( $-9.5 \pm 12.3\%$ ,  $-9.1 \pm 15.8\%$ , respectively) (Figure 1A), and this was also the case for MVIC<sub>20</sub> torque. Similarly, no significant differences between groups were observed for the changes in MVCC<sub>90</sub> torque after the first (e.g., 24 h post-exercise:  $-11.0 \pm 8.2\%$ ) and second ( $-7.8 \pm 14.5\%$ ) bouts (Figure 1B), which was also found for MVCC<sub>180</sub> and MVCC<sub>270</sub> torques. Between bouts, H-H

showed a significant interaction effect ( $P < 0.05$ ) only for MVCC<sub>90</sub> torque, but no significant difference between bouts was found for L-H, or for any other measure.

Regarding RFD, no significant between-group differences were observed for changes in RFD<sub>0-100</sub> (Figure 1C) or RFD<sub>100-200</sub> (Figure 1D) after the first or second bouts. No significant differences in the changes were evident between bouts for each group. No significant differences were found for the first bout in H-H and the second bout in L-H in which both groups performed the high-intensity eccentric cycling for the first time.

SJ decreased significantly from baseline at 24 h post-cycling for both groups ( $-7.7 \pm 10.4\%$ ) and CMJ decreased from baseline at 24 ( $-5.9 \pm 8.4\%$ ), 48 ( $-6.2 \pm 11.1\%$ ) and 72 h after cycling ( $-5.2 \pm 10.5\%$ ), without a significant difference between groups for the first bout. For the second bout, a significant group  $\times$  time interaction effect was observed whereby the reductions in jump height from baseline were greater in L-H than H-H. When comparing the first and second bouts for each group, changes were significantly smaller for the H-H group in SJ and CMJ, and for the L-H group in CMJ.

### **Muscle soreness**

Changes in muscle soreness during palpation were similar between the nine different knee extensor sites, thus the average of the nine sites is shown in Figure 2. Muscle soreness increased at 24 – 72 h after the first and second bouts in both groups ( $P < 0.001$ ), and a significant difference between groups was evident. After the first bout, peak muscle soreness was greater in H-H ( $4.2 \pm 1.0$  on 1 – 10 scale) than L-H ( $1.4 \pm 0.6$ ), but after the second bout it was greater in L-H ( $3.6 \pm 0.9$ ) than H-H ( $1.5 \pm 0.5$ ). When comparing the first and second bouts for each group, a significant interaction effect was found for both groups such that the increases in muscle soreness were smaller after the second bout in H-H but greater after the second bout in L-H, which was smaller when compared with the first bout in H-H. Similar changes were



observed for the gluteal muscle soreness with palpation (1<sup>st</sup> bout; peak H-H:  $4.2 \pm 3.0$ , L-H:  $0.9 \pm 1.9$ , 2<sup>nd</sup> bout; peak H-H:  $1.2 \pm 1.8$ , L-H:  $2.4 \pm 2.6$ ), muscle soreness assessed during a squat (1<sup>st</sup> bout; H-H:  $3.8 \pm 0.1$ , L-H:  $1.0 \pm 0.1$ , 2<sup>nd</sup> bout; H-H:  $1.0 \pm 0.1$ , L-H:  $2.4 \pm 0.2$ ) and stretching of the knee extensors (1<sup>st</sup> bout; H-H:  $4.9 \pm 2.8$ , L-H:  $2.0 \pm 1.4$ , 2<sup>nd</sup> bout; H-H:  $1.9 \pm 1.8$ , L-H:  $3.6 \pm 1.7$ ).

### **Plasma CK activity**

As shown in Figure 3, no significant group  $\times$  time interaction was evident for the changes in plasma CK activity after the first and second bouts, although significant increases were found at 24 h post-exercise in the first bout (average peak CK activity:  $248.2 \pm 272.5$  IU/L), and 24 – 48 h post-exercise in the second bout for both groups. When comparing between bouts for each group, the changes were not statistically different for either group.

### **Muscle ultrasound echo-intensity and thickness**

As shown in Figure 4, no significant changes in the variables were observed after either bout for either group in muscle thickness and echo intensity, and no group  $\times$  time interactions were evident.

## **DISCUSSION**

This was the first study to compare between high- and low-intensity eccentric cycling of the same total mechanical work for changes in muscle function, muscle soreness and other indirect muscle damage markers after the initial exercise bout, and to examine the changes after a second bout of high-intensity eccentric cycling performed 14 days later. The primary finding was that changes in both muscle function and plasma CK activity were similar between the high- and low-intensity eccentric cycling bouts, which was consistent with the hypothesis that

muscle damage evoked by eccentric exercise would not be related to exercise intensity when the exercise was performed submaximally. Nonetheless, muscle soreness was greater after high- than low-intensity cycling, indicating a different effect of exercise intensity on muscle soreness than other muscle damage markers. An additional and unexpected finding of the study was that changes in muscle function and plasma CK activity were similar between the first and second (repeated) bouts of ECC, although a decreased soreness response was observed in H-H whilst greater soreness was observed in the second (high-intensity) bout than the first (low-intensity) bout in L-H.

The present study used two groups to compare two intensities of eccentric cycling, with the average intensities being 154 W for the high-intensity and 36 W for the low-intensity groups. However, while there was a 4-fold difference in intensity, the total work was similar between the groups, as planned. The average heart rate was approximately 20 bpm higher during the high- than low-intensity eccentric cycling, and subjective effort was approximately 2 points greater during high- than low-intensity cycling. Thus, given that eccentric exercise intensity in the current cohort could easily be greater than 500 W if performed maximally (Lipski et al. 2018), even the “high-intensity” eccentric cycling group performed the exercise at a relatively low intensity.

There were no differences observed in muscle function after high- versus low-intensity eccentric cycling (Figure 1), which was consistent with findings of previous studies comparing work-matched submaximal high- and low-intensity eccentric elbow flexor exercise (Gauche et al. 2009; Hill et al. 2019). No previous study has compared the muscle damaging effects between different submaximal intensities of eccentric exercise. However, Paschalis et al. (2005) compared maximal and 50% of maximal eccentric knee extensor exercise matched for mechanical work, and reported a decrease in MVIC torque of 14% in the maximal exercise condition only. Paulsen et al. (2010) and Raastad et al. (2010) reported a 47% reduction in

maximal concentric knee extensor torque after 300 (30 sets of 10) maximal eccentric (isokinetic) knee extensions, which was still 13% below baseline at 7 days post-exercise. It is interesting to note that the number of eccentric actions in the high-intensity cycling in the present study was also 300, but the magnitude of muscle strength decrease was much smaller ( $< 15\%$ ), and most muscle strength measures returned to baseline by 3 days post-exercise. Although the exercise mode was very different, this suggests that the intensity of eccentric actions is a factor influencing the magnitude of muscle function decrement when maximal exercise is performed.

The magnitude of the decreases in all muscle function measures (Figure 1) and changes in plasma CK activity (Figure 3) were similar between the high- and low-intensity eccentric cycling, although the intensity differed by 4-fold. These findings suggest that exercise volume (i.e. number of eccentric actions) is a more important factor affecting the magnitude of decrease in muscle function than exercise intensity when the intensity is submaximal. This has been shown in the previous studies (Gauche et al. 2009; Hill et al. 2019) in which work-matched high- and low-intensity eccentric exercise of the elbow flexors were compared. Taken together, the results of the present study supported the hypothesis that there would be no difference in the magnitude of decrease in muscle function between higher- and lower-intensity submaximal eccentric exercise bouts when the eccentric action intensity was submaximal and the total mechanical work was matched.

It is of interest that muscle soreness developed more after the high- than low-intensity eccentric cycling, and peak muscle soreness in H-H ( $4.2 \pm 1.0$  out of 10) was three times of that of L-H ( $1.4 \pm 0.6$ ). These results contradict the findings of other studies comparing non-work-matched maximal and low-intensity eccentric elbow flexor exercise (Nosaka and Newton 2002) or work-matched maximal and lower-intensity eccentric knee extensor exercise (Paschalis et al. 2005), which did not show significant differences between the two intensities

for DOMS. However, the results of the present study are consistent with previous studies examining non-work-matched submaximal eccentric elbow flexor exercise which reported that higher intensities evoked greater DOMS (Chen et al. 2007, 2012). Although DOMS does not appear to be severe after eccentric cycling, DOMS could be an issue for the prescription of eccentric exercise. Moreover, DOMS was greater after high- than low-intensity eccentric cycling in the second bout (Figure 2). It appears that the magnitude of DOMS is more closely associated with exercise intensity than the mechanical work, even when exercise mechanical work is matched. These results also highlight that the level of DOMS is not necessarily associated with the level of functional loss or other indirect markers of muscle damage (e.g., CK activity in the blood), as shown previously (Nosaka et al. 2002; Paschalis et al. 2005).

It has been documented that DOMS is more associated with damage and inflammation to the muscular connective tissue (i.e. endomysium, perimysium and epimysium) than the muscle fibres (Crameri et al. 2007; Lau et al. 2015; Sorensen et al. 2018). Moreover, muscle fibre strain, rather than force, appears to be the main determinant of the magnitude of histological damage following eccentric exercise (Lieber and Fridén 1993; Warren et al. 1993). Higher-intensity eccentric actions could impose greater mechanical strain on muscle fibres, and thus connective tissue disruption may be increased. In fact, Paulsen et al. (2010) reported the accumulation of inflammatory cells in the endomysium after 300 maximal eccentric knee extensions that induced only a small magnitude of muscle fibre damage (less than 1.4% of whole muscle fibres were infiltrated by the inflammatory cells). It seems possible that the intensity of eccentric actions is a mediator of the magnitude of muscle fibre strain, which could be the primary contributor to the development of DOMS due to connective tissue disruption. To explicitly test this hypothesis, future studies are needed to examine the relationship between DOMS, muscle fibre strain and connective tissue damage induced by eccentric actions.

Previous studies have reported recovery of MVIC torque to baseline by 1 day post-exercise, a peak muscle soreness reduction of 84%, and no significant increase in plasma CK activity after the second bout of eccentric cycling performed at 60% PPO than after the first bout that was performed 2 weeks before (Peñailillo et al. 2013). In that study, the strength loss lasted for 3 days, muscle soreness peaked at 25 mm in a 100-mm visual analogue scale, and peak plasma CK activity was 246.5 IU/L, while in the present study changes in many of the muscle function measures were not statistically different between bouts. Also, although changes in MVCC<sub>90</sub>, SJ, CMJ and muscle soreness were significantly smaller after the second than the first bout in H-H, the magnitude of the differences between bouts observed were not as large as those reported by Peñailillo et al. (2013). Moreover, no changes in muscle function, except CMJ (Figure 1), were observed in L-H between the first and second bouts. Cumulatively, indicators of muscle damage and muscle soreness in the second bout of exercise in the present study were less reduced than those in other studies, suggesting that the lower-intensities of exercise used in the present study induced a smaller RBE.

Previous studies have concluded that a greater RBE is produced when muscle damage in the initial bout is greater (Chen et al. 2007, 2012). As discussed above, changes in muscle damage markers after the first eccentric cycling bout were small in the present study. It seems possible that the relatively small RBE observed in the present study was due to the relatively minor muscle damage produced by the first bout. In fact, there may be a lack of the repeated bout effect when submaximal eccentric exercise causes only a small amount of muscle damage after the first bout (Falvo et al. 2009). Therefore, a ‘ceiling effect’ might explain the relative lack of RBE in the present study.

While the muscle soreness was greater in the second than the first bout in L-H, it was still significantly smaller than the muscle soreness in the first bout of H-H (Figure 2). This indicates that a moderate RBE for muscle soreness was induced by performing low-intensity

eccentric exercise, which is consistent with previous reports of lower-intensity eccentric exercise conferring a protective effect on higher-intensity eccentric exercise (Chen et al. 2012, 2013). As discussed above, DOMS being more associated with connective tissue damage and inflammation, and it being induced more in high- than low-intensity eccentric cycling, probably explains the greater muscle soreness after the second than the first bout in L-H. Several mechanisms of the RBE have been proposed, with extracellular matrix remodelling being considered as a likely player (Hyldahl et al. 2017). It would be interesting to investigate further whether the mechanisms underpinning the RBE differ for muscle soreness than for other symptoms of muscle damage (e.g., loss of muscle function), and then to compare changes in connective tissue damage markers between bouts.

Relative joint power contributions during the eccentric cycling were not assessed in the present study. It is therefore not known whether muscles crossing other joints may have performed significant eccentric work, and that this might have been different between groups. However, given that intensities were submaximal in both conditions, we assumed that the joint contributions would not differ significantly. Future studies should control for joint contributions to ensure that they do not differ significantly between intensities. Moreover, in the present study, we assessed muscle damage through indirect markers only. It would be of interest to examine histological responses in the muscle fibres and extracellular matrix that could potentially provide a clearer understanding of the effects of different intensities of eccentric exercise on the RBE.

The current findings have a number of practical implications. First, the magnitude of muscle damage was minor after eccentric cycling, but even low-intensity eccentric cycling could result in prolonged decreases in muscle function. Previous studies reported increases in muscle function and muscle volume after eccentric cycling training (LaStayo et al. 2000). Thus, eccentric cycling may be suitable for exercise prescription for all populations including older

adults and frail individuals, but potential muscle damage induced by eccentric cycling should be cognisant. Second, DOMS was more significant after the high-intensity exercise, so individuals may “feel” greater pain in the days after exercise if higher initial intensities are performed. Given this finding, it should be noted that muscle pain in the days after exercise may not provide a good indication of the level of muscle damage. Third, regardless of whether the bout was preceded by a high- or low-intensity initial bout, and despite there being no detectable differences in muscle damage markers between conditions, less DOMS was evoked by a subsequent high-intensity bout when the initial bout was performed at a high intensity. Thus, a single bout of lower-intensity eccentric cycling may not be sufficient to minimise DOMS after subsequent bouts. We recommend that intensity should be increased over several sessions in populations who require protection from soreness after training; this is an important consideration for exercise adherence (LaStayo et al. 2014).

In conclusion, muscle damage was not severe even after an unaccustomed bout of high-intensity eccentric cycling, and was not different between high- and low-intensity bouts when the exercise was submaximal and mechanical work matched. Thus, mechanical work, rather than intensity, is probably a more significant factor influencing muscle damage under these conditions. Nonetheless, muscle soreness was greater after high- than low-intensity cycling; under these conditions DOMS does not seem to be a predictor of muscle damage. DOMS was greater when low-intensity rather than high-intensity exercise was performed in the initial bout. So the repeated bout effect for DOMS was intensity-dependent during the submaximal cycling exercise even though exercise mechanical work was matched.

499

## **AUTHOR CONTRIBUTIONS**

500 GM, AB and KN conceived and designed the study, and GM conducted the experiments. GM,  
501 OB, AF, WP, and TK collected and analysed the data. GM drafted the manuscript with AB and  
502 KN. All authors read, edited and approved the manuscript.

503

504

## **FUNDING**

505 No funding was received for this work.

506

507

## **COMPLIANCE WITH ETHICAL STANDARDS**

508 The authors declare no conflict of interest. All participants signed an informed consent form  
509 prior to partaking in the study.

510



- 512 Armstrong RB (1990) Initial events in exercise-induced muscular injury. *Med Sci Sports Exerc*  
 513 22:429–435. <https://doi.org/10.1249/00005768-199008000-00002>
- 514 Borg G (1998) Borg's perceived exertion and pain scales. Human Kinetics, Champaign, IL
- 515 Chen TC, Nosaka K, Sacco P (2007) Intensity of eccentric exercise, shift of optimum angle,  
 516 and the magnitude of repeated-bout effect. *J Appl Physiol* 102:992–999.  
 517 <https://doi.org/10.1152/japplphysiol.00425.2006>
- 518 Chen TC, Tseng W-C, Huang G-L, et al (2013) Low-intensity eccentric contractions attenuate  
 519 muscle damage induced by subsequent maximal eccentric exercise of the knee  
 520 extensors in the elderly. *Eur J Appl Physiol* 113:1005–1015.  
 521 <https://doi.org/10.1007/s00421-012-2517-3>
- 522 Chen TC-C, Chen H-L, Pearce AJ, Nosaka K (2012) Attenuation of eccentric exercise-induced  
 523 muscle damage by preconditioning exercises. *Med Sci Sports Exerc* 44:2090–2098.  
 524 <https://doi.org/10.1249/MSS.0b013e31825f69f3>
- 525 Clarkson PM, Nosaka K, Braun B (1992) Muscle function after exercise-induced muscle  
 526 damage and rapid adaptation. *Med Sci Sports Exerc* 24:512–520.  
 527 <https://doi.org/10.1249/00005768-199205000-00004>
- 528 Crameri RM, Aagaard P, Qvortrup K, et al (2007) Myofibre damage in human skeletal muscle:  
 529 effects of electrical stimulation versus voluntary contraction. *J Physiol* 583:365–380.  
 530 <https://doi.org/10.1113/jphysiol.2007.128827>
- 531 Falvo MJ, Schilling BK, Bloomer RJ, Smith WA (2009) Repeated bout effect is absent in  
 532 resistance trained men: An electromyographic analysis. *J Electromyogr Kinesiol*  
 533 19:e529–e535. <https://doi.org/10.1016/j.jelekin.2008.10.010>
- 534 Fridén J, Sjöström M, Ekblom B (1983) Myofibrillar damage following intense eccentric  
 535 exercise in man. *Int J Sports Med* 4:170–176. <https://doi.org/10.1055/s-2008-1026030>
- 536 Gauche E, Couturier A, Lepers R, et al (2009) Neuromuscular fatigue following high versus  
 537 low-intensity eccentric exercise of biceps brachii muscle. *J Electromyogr Kinesiol*  
 538 19:e481–e486. <https://doi.org/10.1016/j.jelekin.2009.01.006>
- 539 González-Bartholin R, Mackay K, Valladares D, et al (2019) Changes in oxidative stress,  
 540 inflammation and muscle damage markers following eccentric versus concentric  
 541 cycling in older adults. *Eur J Appl Physiol* 119:2301–2312.  
 542 <https://doi.org/10.1007/s00421-019-04213-7>
- 543 Hasenoehrl T, Wessner B, Tschan H, et al (2017) Eccentric resistance training intensity may  
 544 affect the severity of exercise induced muscle damage. *J Sports Med Phys Fitness*  
 545 57:1195–1204. <https://doi.org/10.23736/S0022-4707.16.06476-8>
- 546 Hill EC, Housh TJ, Smith CM, et al (2019) High- vs. low-intensity fatiguing eccentric exercise  
 547 on muscle thickness, strength, and blood flow. *J Strength Cond Res Publish Ahead of*  
 548 *Print*: <https://doi.org/10.1519/JSC.0000000000002632>

549 Hortobágyi T, Houmard J, Fraser D, et al (1998) Normal forces and myofibrillar disruption  
550 after repeated eccentric exercise. *J Appl Physiol* 84:492–498.  
551 <https://doi.org/10.1152/jappl.1998.84.2.492>

552 Hyldahl RD, Chen TC, Nosaka K (2017) Mechanisms and mediators of the skeletal muscle  
553 repeated bout effect. *Exerc Sport Sci Rev* 45:24–33.  
554 <https://doi.org/10.1249/JES.0000000000000095>

555 LaStayo P, Marcus R, Dibble L, et al (2014) Eccentric exercise in rehabilitation: safety,  
556 feasibility, and application. *J Appl Physiol* 116:1426–1434.  
557 <https://doi.org/10.1152/jappphysiol.00008.2013>

558 LaStayo PC, Pierotti DJ, Pifer J, et al (2000) Eccentric ergometry: increases in locomotor  
559 muscle size and strength at low training intensities. *Am J Physiol Regul Integr Comp*  
560 *Physiol* 278:R1282–1288. <https://doi.org/10.1152/ajpregu.2000.278.5.R1282>

561 Lau WY, Blazeovich AJ, Newton MJ, et al (2015) Changes in electrical pain threshold of fascia  
562 and muscle after initial and secondary bouts of elbow flexor eccentric exercise. *Eur J*  
563 *Appl Physiol* 115:959–968. <https://doi.org/10.1007/s00421-014-3077-5>

564 Lieber RL, Fridén J (1993) Muscle damage is not a function of muscle force but active muscle  
565 strain. *J Appl Physiol* 74:520–526. <https://doi.org/10.1152/jappl.1993.74.2.520>

566 Lipski M, Abbiss CR, Nosaka K (2018) Cardio-pulmonary responses to incremental eccentric  
567 and concentric cycling tests to task failure. *Eur J Appl Physiol* 118:947–957.  
568 <https://doi.org/10.1007/s00421-018-3826-y>

569 Maffiuletti NA, Aagaard P, Blazeovich AJ, et al (2016) Rate of force development:  
570 physiological and methodological considerations. *Eur J Appl Physiol* 116:1091–1116.  
571 <https://doi.org/10.1007/s00421-016-3346-6>

572 McHugh MP (2003) Recent advances in the understanding of the repeated bout effect: the  
573 protective effect against muscle damage from a single bout of eccentric exercise. *Scand*  
574 *J Med Sci Sports* 13:88–97. <https://doi.org/10.1034/j.1600-0838.2003.02477.x>

575 Nosaka K, Clarkson PM (1995) Muscle damage following repeated bouts of high force  
576 eccentric exercise. *Med Sci Sports Exerc* 27:1263–1269.  
577 <https://doi.org/10.1249/00005768-199509000-00005>

578 Nosaka K, Newton M (2002) Difference in the magnitude of muscle damage between maximal  
579 and submaximal eccentric loading. *J Strength Cond Res* 16:202–208.  
580 <https://doi.org/10.1519/00124278-200205000-00006>

581 Nosaka K, Newton M, Sacco P (2002) Delayed-onset muscle soreness does not reflect the  
582 magnitude of eccentric exercise-induced muscle damage. *Scand J Med Sci Sports*  
583 12:337–346. <https://doi.org/10.1034/j.1600-0838.2002.10178.x>

584 Paschalis V, Koutedakis Y, Jamurtas AZ, et al (2005) Equal volumes of high and low intensity  
585 of eccentric exercise in relation to muscle damage and performance. *J Strength Cond*  
586 *Res* 19:184–188. <https://doi.org/10.1519/R-14763.1>

587 Paulsen G, Cramer R, Benestad HB, et al (2010) Time course of leukocyte accumulation in  
588 human muscle after eccentric exercise. *Med Sci Sports Exerc* 42:75–85.  
589 <https://doi.org/10.1249/MSS.0b013e3181ac7adb>

590 Peñailillo L, Blazeovich A, Numazawa H, Nosaka K (2013) Metabolic and muscle damage  
591 profiles of concentric versus repeated eccentric cycling. *Med Sci Sports Exerc*  
592 45:1773–1781. <https://doi.org/10.1249/MSS.0b013e31828f8a73>

593 Peñailillo L, Blazeovich A, Numazawa H, Nosaka K (2015a) Rate of force development as a  
594 measure of muscle damage. *Scand J Med Sci Sports* 25:417–427.  
595 <https://doi.org/10.1111/sms.12241>

596 Peñailillo L, Blazeovich AJ, Nosaka K (2015b) Muscle fascicle behavior during eccentric  
597 cycling and its relation to muscle soreness. *Med Sci Sports Exerc* 47:708–717.  
598 <https://doi.org/10.1249/MSS.0000000000000473>

599 Peñailillo L, Mackay K, Abbiss CR (2018) Rating of perceived exertion during concentric and  
600 eccentric cycling: are we measuring effort or exertion? *Int J Sports Physiol Perform*  
601 13:517–523. <https://doi.org/10.1123/ijsp.2017-0171>

602 Raastad T, Owe SG, Paulsen G, et al (2010) Changes in calpain activity, muscle structure, and  
603 function after eccentric exercise. *Med Sci Sports Exerc* 42:86–95.  
604 <https://doi.org/10.1249/MSS.0b013e3181ac7afa>

605 Sorensen JR, Skousen C, Holland A, et al (2018) Acute extracellular matrix, inflammatory and  
606 MAPK response to lengthening contractions in elderly human skeletal muscle. *Exp*  
607 *Gerontol* 106:28–38. <https://doi.org/10.1016/j.exger.2018.02.013>

608 Tsuchiya Y, Ueda H, Ochi E (2019) Muscular recruitment is associated with muscular function  
609 and swelling following eccentric contractions of human elbow flexors. *J Sports Med*  
610 *Phys Fitness* 59:1097–1101. <https://doi.org/10.23736/s0022-4707.18.09102-8>

611 Valladares-Ide D, Peñailillo L, Collao N, et al (2019) Activation of protein synthesis,  
612 regeneration and MAPK signaling pathways following repeated bouts of eccentric  
613 cycling. *Am J Physiol Endocrinol Metab*. <https://doi.org/10.1152/ajpendo.00216.2019>

614 Warren GL, Hayes DA, Lowe DA, Armstrong RB (1993) Mechanical factors in the initiation  
615 of eccentric contraction-induced injury in rat soleus muscle. *J Physiol* 464:457–475.  
616 <https://doi.org/10.1113/jphysiol.1993.sp019645>

617 Wobbrock JO, Findlater L, Gergle D, Higgins JJ (2011) The aligned rank transform for  
618 nonparametric factorial analyses using only ANOVA procedures. In: CHI 2011 - 29th  
619 Annual CHI Conference on Human Factors in Computing Systems, Conference  
620 Proceedings and Extended Abstracts. Vancouver, BC, Canada, pp 143–146

621 Yu J-G, Malm C, Thornell L-E (2002) Eccentric contractions leading to DOMS do not cause  
622 loss of desmin nor fibre necrosis in human muscle. *Histochem Cell Biol* 118:29–34.  
623 <https://doi.org/10.1007/s00418-002-0423-1>

624

625

## TABLES

626

627 Table 1: Participant characteristics (mean  $\pm$  SD) for each group (H-H, L-H) before the first  
 628 eccentric cycling exercise bout. P-values are based on independent samples t-tests for  
 629 comparison between groups.

	H-H (n=11)	L-H (n = 9)	P
Age (years)	25.3 $\pm$ 3.6	24.7 $\pm$ 5.9	0.781
Height (cm)	181.2 $\pm$ 7.2	178.6 $\pm$ 8.6	0.467
Body mass (kg)	83.3 $\pm$ 11.4	75.7 $\pm$ 10.2	0.139
BMI (kg/m <sup>2</sup> )	25.4 $\pm$ 3.2	23.7 $\pm$ 2.7	0.233
PPO (W)	778.5 $\pm$ 120.3	702.9 $\pm$ 105.1	0.157
MVIC <sub>70</sub> (Nm)	269.6 $\pm$ 79.8	251.4 $\pm$ 61.3	0.582
MVIC <sub>20</sub> (Nm)	105.5 $\pm$ 30.9	96.0 $\pm$ 20.2	0.435
MVCC <sub>270</sub> (Nm)	139.4 $\pm$ 25.4	123.0 $\pm$ 26.1	0.172
MVCC <sub>180</sub> (Nm)	160.8 $\pm$ 31.3	143.0 $\pm$ 29.7	0.211
MVCC <sub>90</sub> (Nm)	200.0 $\pm$ 37.9	182.8 $\pm$ 43.3	0.383
SJ (cm)	35.0 $\pm$ 4.2	35.5 $\pm$ 6.7	0.851
CMJ (cm)	38.4 $\pm$ 5.6	37.6 $\pm$ 6.6	0.782
CK activity (IU/L)	99.4 $\pm$ 50.2	113.1 $\pm$ 92.6	0.679

630 BMI: body mass index; PPO: peak power output during a maximal 10-s isokinetic concentric  
 631 cycling effort at 60 rpm; MVIC: maximal voluntary isometric contraction torque at knee joint  
 632 angles of 70° (MVIC<sub>70</sub>) and 20° (MVIC<sub>20</sub>); MVCC: maximal voluntary isokinetic concentric  
 633 contraction torque at angular velocities of 270°/s (MVCC<sub>270</sub>), 180°/s (MVCC<sub>180</sub>) and 90°/s  
 634 (MVCC<sub>90</sub>); SJ: squat jump; CMJ: counter-movement jump height; CK: creatine kinase.

635

636

## FIGURE LEGENDS







**Figure 1.** Maximal voluntary isometric contraction torque at 70° (A; mean  $\pm$  SD), isokinetic (90°/s) concentric contraction torque (B), RFD 0-100 ms (C), RFD 100-200 ms (D), squat jump power (E), and countermovement jump power (F) before (Pre), immediately after (Post), and 24-72 h after the first and second exercise bouts for H-H (●) and L-H (○). \*: significant ( $P < 0.05$ ) difference from the baseline for both groups. ANOVA results for interaction (bout  $\times$  time) effect are shown in the legend for each bout. Results of comparisons between groups and between bouts are shown on the top of each figure.

**Figure 2.** Muscle soreness upon palpation (average value of the nine knee extensor sites  $\pm$ SD) before (Pre), immediately after (Post), and 24-72 hours after the first and second exercise bouts for H-H (●) and L-H (○). \*: significant ( $P < 0.05$ ) difference from the baseline for both groups. #: significant ( $P < 0.05$ ) difference between groups. ANOVA results for interaction (bout  $\times$  time) are shown in the legend for each bout. Results of comparisons between groups and between bouts are shown on the top of the figure.


**Figure 3.** Plasma creatine kinase (CK) activity (mean  $\pm$  SD) before (Pre), and 24-72 hours after the first and second exercise bouts for the H-H (●) and L-H (○) groups. \*: significant ( $P < 0.05$ ) difference from the baseline for both groups. ANOVA results for interaction (bout  $\times$  time) are shown in the legend for each bout. Results of comparisons between groups and between bouts are shown on the top of the figure.

**Figure 4.** Ultrasound echo intensity (top) and muscle thickness (bottom) before (Pre), immediately after (Post), and 24-72 hours after the first and second exercise bouts (mean  $\pm$  SD)

662 for the H-H (●) and L-H (○) groups. ANOVA results for interaction (bout × time) are shown  
663 in the legend for each bout. Results of comparisons between groups and between bouts are  
664 shown on the top of each figure.




H-H: 1st vs 2nd ( $P < 0.01$ ), L-H: 1st vs 2nd ( $P < 0.01$ ),  
H-H 1st vs L-H 2nd ( $P < 0.01$ ), L-H 1st vs H-H 2nd (n.s.)






H-H: 1st vs 2nd (n.s.), L-H: 1st vs 2nd (n.s.),  
H-H 1st vs L-H 2nd (n.s.), L-H 1st vs H-H 2nd (n.s.)



H-H: 1st vs 2nd (n.s.), L-H: 1st vs 2nd (n.s.),  
H-H 1st vs L-H 2nd (n.s.), L-H 1st vs H-H 2nd (n.s.)



H-H: 1st vs 2nd (n.s.), L-H: 1st vs 2nd (n.s.),  
H-H 1st vs L-H 2nd (n.s.), L-H 1st vs H-H 2nd (n.s.)

