

1-1-2020

A rapid absorbance-based growth assay to screen the toxicity of oligomer A β 42 and protect against cell death in yeast

Prashant Bharadwaj
Edith Cowan University

Ralph Martins
Edith Cowan University

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworkspost2013>



Part of the [Medicine and Health Sciences Commons](#)

10.4103/1673-5374.280318

Bharadwaj, P., & Martins, R. (2020). A rapid absorbance-based growth assay to screen the toxicity of oligomer A β 42 and protect against cell death in yeast. *Neural Regeneration Research*, 15(10), Article 1931. <https://doi.org/10.4103/1673-5374.280318>

10.4103/1673-5374.280318

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworkspost2013/8136>

A rapid absorbance-based growth assay to screen the toxicity of oligomer A β ₄₂ and protect against cell death in yeast

Prashant Bharadwaj^{1,2,*}, Ralph Martins^{1,3,4}

1 Centre of Excellence for Alzheimer's Disease Research and Care, School of Medical and Health Sciences, Edith Cowan University, Western Australia, Australia

2 School of Pharmacy and Biomedical Sciences, Curtin Health and Innovation Research Institute (CHIRI), Faculty of Health Sciences, Curtin University, Western Australia, Australia

3 School of Psychiatry and Clinical Neuroscience, University of Western Australia, Western Australia, Australia

4 School of Biomedical Science, Macquarie University, Sydney, NSW, Australia

Funding: This study was supported by the National Health and Medical Research Council-Australian Research Council dementia research development fellowship (APP1107109) to PB.

Abstract

Multiple lines of evidence show that soluble oligomer forms of amyloid β protein (A β ₄₂) are the most neurotoxic species in the brain and correlates with the degree of neuronal loss and cognitive deficit in Alzheimer's disease. Although many studies have used mammalian cells to investigate oligomer A β ₄₂ toxicity, the use of more simple eukaryotic cellular systems offers advantages for large-scale screening studies. We have previously established and validated budding yeast, *Saccharomyces cerevisiae* to be a simple and a robust model to study the toxicity of A β . Using colony counting based methods, oligomeric A β ₄₂ was shown to induce dose-dependent cell death in yeast. We have adapted this method for high throughput screening by developing an absorbance-based growth assay. We further validated the assay with treatments previously shown to protect oligomer A β ₄₂ induced cell death in mammalian and yeast cells. This assay offers a platform for studying underlying mechanisms of oligomer A β ₄₂ induced cell death using gene deletion/overexpression libraries and developing novel agents that alleviate A β ₄₂ induced cell death.

Key Words: Alzheimer's disease; amyloid toxicity; autophagy; A β ₄₂ oligomer; high-throughput screening; latrepirdine; neuroprotection; yeast model

Chinese Library Classification No. R446; R364; R741

*Correspondence to:

Prashant Bharadwaj, PhD,
p.bharadwaj@ecu.edu.au.

orcid:

0000-0003-4361-9906
(Prashant Bharadwaj)

doi:10.4103/1673-5374.280318

Received: September 24, 2019

Peer review started: September 30, 2019

Accepted: December 13, 2019

Published online: April 3, 2020

Introduction

Alzheimer's disease is an incurable neurodegenerative disorder featuring progressive loss of memory and ability to perform daily tasks ultimately leading to death. The disease is characterized by deposition of intracellular neurofibrillary tangles and extracellular amyloid plaques within the brain. The cerebral amyloid plaques are insoluble protein deposits associated with dystrophic neurites, inflammatory processes and a dense core comprising primarily of aggregated fibrillar forms of A β protein (Fraser et al., 1993). The amyloid β protein (A β) is a 40/42-amino acid product of amyloid precursor protein and is found in the brains and cerebrospinal fluid of both healthy individuals and people with AD (reviewed in (Selkoe and Hardy, 2016)). Numerous isoforms of A β have been detected in the brain, however the AD brain exhibits markedly increased levels of the longer aggregate-prone isoform, A β ₄₂ [reviewed in Bharadwaj et al. (2009)]. Soluble oligomeric form of A β ₄₂ correlates with cognitive loss in AD as compared to larger fibrillar A β aggregates and is considered the most neurotoxic species (McLean et al., 1999; Dahlgren et al., 2002).

In vitro and *in vivo* studies have shown that A β oligomers induce neuronal synaptic loss and degeneration and also reduce viability of non-neuronal mammalian cells and non-mamma-

lian cells, including yeast (Bharadwaj et al., 2008; Treusch et al., 2011), suggesting a common mechanism for oligomer A β ₄₂ induced toxicity shared between different cell types. Yeasts are eukaryotic cells and offer greater advantage over other disease models mainly due to the ease of manipulation, availability of powerful genomic and proteomic tools and high-throughput screening techniques. In addition to genetic similarity, yeast cells show similarity in numerous metabolic and signaling pathways with humans, especially those involved in ageing and cellular death. Yeast models have been used extensively to study misfolded or aggregated proteins involved in late-onset neurodegenerative disorders such as Huntington polyglutamine repeats, α -synuclein, superoxide dismutase 1 and A β ₄₂ (Bharadwaj et al., 2010). Due to their robust nature and ability to effectively coordinate the activities of protective stress response pathways to toxic proteins, they are ideally suited for investigating disease relevant pathways and for testing of small molecules that protect against cell death.

Similar to neuronal cells (Dahlgren et al., 2002), we have previously shown that oligomer A β ₄₂ is significantly more toxic than fibrillar A β ₄₂ and induced dose-dependent cell death in yeast (Bharadwaj et al., 2008). This yeast model for A β ₄₂ toxicity have been implemented for investigating the neuroprotective effects of whey derived peptides (Bharad-

waj et al., 2013), functional characterisation of A β fusion proteins (Caine et al., 2011) and in studying the role of autophagy in protection against A β_{42} toxicity (Bharadwaj et al., 2012). These assays have been based on colony counting which is time-consuming to set-up and unsuitable for screening of gene deletion/overexpression libraries and small molecule collections. To increase feasibility for such broad-based screening methods, here we have developed a growth based high-throughput assay based on our established colony-count method to measure oligomer A β_{42} induced cell death. This novel assay was further validated by assessing known protective agents that promote survival against oligomer A β_{42} mediated cell death.

Materials and Methods

Yeast strain

Wild-type *Saccharomyces cerevisiae* strain BY4743 [genotype: *MATa/α his3Δ1/his3Δ1 leu2Δ0/leu2Δ0 LYS2/lys2Δ0 met15Δ0/MET15 ura3Δ0/ura3Δ0*] (Dr. Gabriel Perrone University of New South Wales) was used for this study. Yeast cells were grown on YEPD (1% yeast extract; 2% peptone; 2% dextrose), YNB (yeast nutrient base) synthetic complete or nitrogen starvation media, YNB (-N) for toxicity experiments. Yeast cells were stored as glycerol stocks at -80°C and in media plates at 4°C .

Reagents

Autophagy modulating compounds including Latrepirdine, Rapamycin and SMER28 were used as controls in this study. Latrepirdine (Dimebolin dihydrochloride, molecular weight (MW): 392.37) was purchased from Biotrend AG, Zurich, Switzerland. Latrepirdine was dissolved into sterile double distilled water to 30 mM concentration. Rapamycin and SMER28 were purchased from Sigma (Missouri, USA) and solutions were prepared in DMSO (Dimethyl sulfoxide). Freshly prepared working stocks (1 mM) prepared from -80°C frozen stocks were used for all experiments. Synthetic human A β_{42} and A β_{16} were purchased from ERI Laboratory (Connecticut). Recombinant human insulin (I2643, MW: 5800 Da) was purchased from Sigma.

A β preparation

A β peptides were prepared according to Bharadwaj et al. (2012). A β peptide (0.5 mg) was dissolved in 500 μL hexafluoroisopropanol solution and incubated overnight at room temperature. The hexafluoroisopropanol was then evaporated using vacuum and the dried peptide films were stored in a -80°C freezer. A β peptide films were dissolved in 200 μL sterile double-distilled water and prepared prior to use. The peptide solution was vortexed followed by sonication for 5 minutes on ice and then centrifuged for 10 minutes at $14,000 \times g$. The resulting supernatant containing soluble A β oligomers and was used for yeast toxicity assays following overnight incubation at room temperature to allow oligomerisation. Insulin (MW: 4.8 kDa) was freshly prepared in $1 \times \text{PBS}$ and truncated A β_{16} (MW: 1.6 kDa) was prepared in a similar fashion as A β_{42} for toxicity assays.

A β toxicity assay in yeast: colony forming unit assay

Wild-type *Saccharomyces cerevisiae* (strain, BY4743) was used for A β toxicity experiments. Yeast cells were stored on YEPD agar plates at 4°C . A single colony was inoculated in YEPD liquid media and incubated by shaking at 30°C overnight. The overnight yeast culture was re-inoculated in fresh YEPD liquid media and incubated with shaking and grown for 3–4 hours at 30°C to reach exponential phase growth. For inducing autophagy, cells were washed in sterile water and pre-incubated with latrepirdine (5, 10, and 20 μM), rapamycin (50, 100, 150, and 200 nM), SMER28 (small molecule enhancer of rapamycin, 10, 20, and 50 μM) or incubated in nitrogen starvation media (YNB, -N). These cultures were diluted to $\sim 5 \times 10^3$ yeast cells/mL in sterile distilled water. Cells were then dispensed into 96-well microtitre plates for peptide treatments. Vehicle or peptide preparations were added to the diluted cell suspension to required concentrations. The final volume in each well was made up to $\sim 100 \mu\text{L}$ and then the microtitre plate was sealed with a gas permeable membrane (optional) and incubated at 30°C by shaking for time periods as indicated. The cell suspensions were then plated on YEPD agar plates and incubated at 30°C for 2 days to measure the number of colony-forming units (CFU). Cell viability following peptide treatment was calculated from the CFU count and represented as percent change compared to vehicle treatment.

Absorbance-based growth assay

In this study, we adapted the colony forming unit assay to an absorbance-based growth assay using a 96-well plate format to increase the feasibility of this yeast model in large scale screening techniques. Cells were treated with peptides and/or autophagy modulating reagents as described above. Exponentially growing yeast cells were treated with increasing doses of oligomeric A β_{42} for 24 hours, and then the cell suspensions were supplemented with yeast growth media (YEPD, 50 μL) and the growth was measured after 16-hour incubation at 30°C by absorbance measurement (optical density, OD600 nm) using a spectrophotometer FLUOstar Optima plate reader (BMG Labtech, Ortenberg, Germany). The cell density measured as absorbance values was represented as percent cell viability. Cell viability in peptide treated samples was evaluated from the absorbance values from similar time points as the untreated and represented as percent change from vehicle treated.

Statistical analysis

For the colony count assay cell viability was determined by counting colonies across the various treatments as described previously (Bharadwaj et al., 2008). For the growth assay, the mean absorbance values between treatment replicates were used to determine cell survival across the various peptide treatments used. All comparisons were made using two-tailed Student's *t*-test (equal sample size, unequal variances) using GraphPad Prism software version 8 (GraphPad Software Inc., La Jolla, CA, USA).

Results

Oligomer A β_{42} toxicity

Initially, we compared the colony forming unit assay and the absorbance-based growth assay to measure oligomer A β_{42} toxicity (Figure 1). Both assays revealed a dose dependent toxicity profile, however colony counting showed greater sensitivity at lower A β_{42} concentrations (0.5–2 μ M) (Figure 1B). Both assays were comparable at higher A β_{42} concentrations (4–10 μ M) (Figure 1B). We also compared the toxicity of oligomer A β_{42} as the positive control and non-aggregating C-terminal truncated A β_{16} peptide (MW 1760 Da) and insulin peptide (MW 5800 Da) were used as the negative controls (Figure 2). Oligomer A β_{42} showed a dose-dependent loss in cell viability, whereas A β_{16} and insulin treatment did not alter cell viability at similar concentrations tested (Figure 2). It is noted that cell viability analysis using the growth-based assay showed 50–80% survival at 2 μ M, 40–60% at 4 μ M and 10% at 10 μ M in Figure 1B, whereas Figure 2 showed 75–85% survival at 2 μ M, 50% at 4 μ M and 10% at 10 μ M.

Inhibitors of oligomer A β_{42} toxicity

Wild type yeast was incubated in nitrogen depleted media (Figure 3A) or treated with latrepirdine (5–20 μ M) (Figure 3B), rapamycin (50–200 nM) (Figure 3C), or SMER28 (10–50 μ M) to induce autophagy followed by oligomer A β_{42} treatment (2–10 μ M) (Figure 3D). A dose dependent loss in viability was observed in cells treated with oligomeric A β_{42} . Low concentrations of A β_{42} (2 μ M) showed ~20% cell death and higher concentrations ranging 4–10 μ M showed 50–80% loss in viability. Cell viability following A β_{42} treatment was significantly higher in nitrogen starved, latrepirdine, rapamycin or SMER28 treated cells compared to vehicle treated (Figure 3). Notably, only nitrogen starvation showed significant protection (a 40% increase in viability) against the highest A β_{42} concentration tested (10 μ M). Whereas, latrepirdine, rapamycin and SMER28 increased cell survival (20–60% increase in viability) only at lower A β_{42} concentrations (2–5 μ M). The incubation period with the autophagy enhancers was found to be critical for inducing protection against oligomer A β_{42} toxicity. Shorter incubation times (6 hours, 30°C) were used for latrepirdine, rapamycin and nitrogen starvation. However, longer incubations (16 hours, 30°C) were required for SMER28 to promote cell viability with A β_{42} treatment (Figure 3). The results showed significant differences in the levels of protection against oligomer A β_{42} toxicity conferred by physiological (nitrogen starvation) and chemical inducers (latrepirdine, rapamycin and SMER28) of autophagy.

Discussion

A novel high-throughput assay for measuring oligomer A β_{42} toxicity

Cell based high-throughput assays are indispensable for analysis of signaling pathways and discovery of small-molecule modulators of disease processes. The budding yeast, *Saccharomyces cerevisiae* has been broadly employed for mechanism-of-action studies and drug target identification.

Existing high-throughput viability assays such as colony count and spotting assay are based on clonogenic analysis (Bharadwaj et al., 2010), which measures the effectiveness of specific agents on the survival and proliferation of yeast cells in agar media. Compared to such clonogenic assays, absorbance-based viability assays offer a simple and more sensitive technique for screening gene deletion/overexpression libraries and compound repositories.

In our previous work, we developed a novel assay for measuring toxicity of oligomer A β_{42} peptide, a key protein associated with pathogenesis and disease development in AD (Bharadwaj et al., 2008, 2012). Similar to neuronal cells, oligomer A β_{42} was found to be more toxic and induced dose dependent cell death in yeast compared to fibrillar A β_{42} . The original method used a colony count based assay to measure oligomer A β_{42} toxicity (Bharadwaj et al., 2008). In this study, we have developed and validated an absorbance based high-throughput assay to measure toxicity of oligomer A β_{42} . It is noted that cell viability analysis using the growth-based assay showed slight variability in Figures 1B and 2. It is likely that the differences in the standard deviation is due to the number of replicates conducted for experiments in Figures 1B ($n = 8$) and 2 ($n = 12$). However, the overall level of toxicity observed in Figures 1B and 2 are consistent. Notably, both the colony count and the absorbance-based growth assay in yeast showed significant loss in cell viability at oligomer A β_{42} concentrations (≥ 1 μ M) required to induce toxicity in neuronal cells (Dahlgren et al., 2002). This yeast assay offers a robust and high-throughput cell-based platform for analysis of disease relevant pathways and screening compounds that can alleviate oligomer A β_{42} induced cell death.

Previous studies have developed high-throughput yeast assays to measure A β_{42} induced toxicity by targeting it to the secretory pathway (Treusch et al., 2011) or used strains expressing A β_{42} fused to the essential functional domain of Sup35, a prion protein, to measure A β_{42} oligomerization (Park et al., 2011). Some of the advantages of our yeast model compared to the studies of Park et al. and Treusch et al. include 1) ability to test toxicity of different A β peptides and its various isoforms and 2) suitability for genetic screening as the assay uses structurally characterized synthetic A β_{42} preparations and eliminate the effects of gene deletion/overexpression on A β_{42} production. Also, the use of an absorbance-based growth assay increases its suitability for screening gene deletion/overexpression and drug libraries as compared to the use of fluorescent/colorimetric dyes which has limitations due to likelihood of toxicity in certain mutants, interference with cell metabolism and fluorescent/colorimetric assay read out. Overall, data from the absorbance-based assay indicated its suitability for large scale, high-throughput screening methods and offers a robust cell-based platform for analysis of disease relevant pathways and screening compounds that can alleviate oligomer A β_{42} induced cell death.

Induction of autophagy protects against oligomer A β_{42} toxicity

A wide range of inhibitors of A β_{42} aggregation, including




Figure 1 Comparative analysis of oligomer Aβ₄₂ toxicity in yeast by colony count and the absorbance-based high-throughput assay. (A) Exponentially growing yeast cells in YEPD (1% yeast extract; 2% peptone; 2% dextrose) media were serially diluted in sterile double distilled water and treated with oligomer Aβ₄₂ peptide for 24 hours. Following treatment, the cell suspensions were supplemented with fresh YEPD media and the growth/cell density was monitored for 16 hours at 30°C by absorbance measurement (optical density, OD₆₀₀). (B) The absorbance values measured at 16 hours are represented as percent cell viability for each treatment and directly compared to those values obtained with the established colony count assay (Bharadwaj et al., 2008). Although less sensitive at low concentrations (1–2 μM) of oligomer Aβ₄₂ treatment compared to the colony count assay, the high-throughput assay shows dose-dependent loss of viability with oligomer Aβ₄₂ treatment, compared to vehicle treatment. Statistical significance between treatments was assessed using two-tailed Student’s *t*-test (*n* = 8, Aβ₄₂ treated vs. untreated, **P* < 0.05, ***P* < 0.01). (C) Images showing the colony count and the high-throughput assay. Aβ: Amyloid β protein.




Figure 2 Toxicity of oligomer Aβ₄₂, Aβ₁₆ and insulin peptides. Exponentially growing yeast cells in YEPD (1% yeast extract; 2% peptone; 2% dextrose) media were serially diluted in sterile double distilled water and treated with oligomer Aβ₄₂, Aβ₁₆ and insulin peptide preparations for 24 hours. Oligomer Aβ₄₂ treatment showed dose dependent loss of viability, whereas Aβ₁₆ and insulin treatment were significantly less toxic as compared to vehicle treated. Statistical significance between treatments was assessed using two-tailed Student’s *t*-test (*n* = 12, Aβ₄₂ treated vs. untreated, **P* < 0.05, ***P* < 0.01). Aβ: Amyloid β protein.

peptide and small molecules of natural and synthetic origin have been shown to regulate toxicity of Aβ₄₂ (Mangialasche et al., 2010). An important caveat in designing such inhibitors is to target the toxic form of Aβ₄₂ and identify compounds with low cytotoxicity. In addition to inhibiting the formation or disaggregating the Aβ₄₂ toxic oligomer structures, there is growing interest towards inducing effective cellular responses to Aβ₄₂ induced toxicity and cell death, and thereby suppress the downstream apoptotic cascades and necrosis in the AD brain. A cellular pathway that can confer such protection is autophagy (Nixon, 2013). Autophagy is a vital intracellular catabolic pathway that plays an important role in the removal of aggregated or misfolded proteins in neurodegenerative diseases. Modulation of autophagy can enhance protective responses to toxicity, and suppress apoptotic cascades or necrosis leading to neurodegeneration in the brain (Mputhia et al., 2019).

Substantial benefits have been observed with autophagy-inducing agents in various neurodegenerative disease models (Nixon, 2013). A widely used inducer of autophagy, rapamycin is a selective inhibitor of mTOR (mammalian tar-




Figure 3 Induction of autophagy protects against oligomer A β_{42} toxicity in yeast.

Exponentially growing yeast cells were incubated in (A) nitrogen depleted media or (B) YNB (yeast nutrient base)-synthetic complete media containing autophagy modulating compounds, i.e., latrepirdine (5, 10 and 20 μ M) or (C) rapamycin (50, 100, 150 and 200 nM) and (D) SMER28 (10, 20 and 50 μ M)) at 30°C to induce autophagy followed by oligomer A β treatment. Loss of cell viability from the different treatments was then measured using the absorbance-based A β_{42} toxicity assay. Cell viability following oligomer A β_{42} treatment was significantly higher in nitrogen starved or latrepirdine treated wild type cells compared to vehicle treated. Statistical significance between treatments was assessed using two-tailed Student's *t*-test (Different A β_{42} treated groups were compared. In the A β_{42} treated groups, control cells and cells treated with autophagy modulators were compared. **P* < 0.05, ***P* < 0.01). Only nitrogen starvation showed significant protection (40% increase in viability) against 10 μ M A β_{42} . Latrepirdine, rapamycin and SMER28 increased cell survival (20–40% increase in viability) only at 2–5 μ M A β_{42} . A β : Amyloid β protein.

get of rapamycin) (Huang et al., 2004). Rapamycin has been shown to reduce A β induced neurodegeneration and improve cognition in transgenic mouse models (Spilman et al., 2010). An mTOR-independent autophagy inducer, SMER28 (small-molecule enhancer of rapamycin) (Huang et al., 2004) has shown to reduce the levels of A β and alleviate toxicity in Huntington disease model (Tian et al., 2011). Recently, we showed that enhanced autophagic activity mediated by rapamycin and a pro-neurogenic compound, latrepirdine protected yeast cells against oligomer A β_{42} toxicity (Bharadwaj et al., 2012). We also showed that latrepirdine enhanced mTOR- and Atg5-dependent autophagy and improved cognition in AD mouse model (Steele et al., 2012). Here, we validated the use of the absorbance-based A β_{42} toxicity assay by determining the protective ability of different classes of autophagy enhancers. The results showed significant differences in the levels of protection against oligomer A β_{42} toxicity conferred by physiological (nitrogen starvation) and chemical inducers (latrepirdine, rapamycin and SMER28) of autophagy. These findings indicate how autophagy can be modulated distinctly by different cellular pathways of induction and how they respond to increasing levels of toxicity and cellular damage caused by oligomeric A β_{42} .

Previous studies favour a strategy of enhancing the efficacy of autophagy by preventing or reversing the impairments of the specific processes that are disrupted. Several compounds with autophagy-related functions as the primary or secondary drug target have been trialled for several neurodegenerative diseases including AD (Nixon, 2013). Although autophagy activation is a promising intervention, there are some caveats to consider (Sarkar, 2013; Mputhia et al., 2019). In diseases like AD, where there is a pre-existing impairment in autophagy, the success of any autophagy-based intervention may depend on whether lysosomal clearance is functional (Nixon, 2013; Mputhia et al., 2019; Tripathi et al., 2019). In addition, effects of autophagy activation may also vary significantly depending on the physiological state of the cell, especially during proteotoxic stress. In fact, some studies suggest that autophagy activation may be harmful in ageing conditions with pre-existing pathology (Carosi and Sargeant, 2019; Tripathi et al., 2019).

In summary, our previous work (Bharadwaj et al., 2012) together with the current findings provides evidence for autophagy as a preventative treatment against oligomer A β_{42} mediated cell death. In addition, here we have established the potential use of the high-throughput yeast assay as a bi-

ological screen to dissect disease-relevant pathways and network of genes that mediate protection against A β_{42} -induced cell death. Although yeast is an excellent screening model for inhibitors of A β_{42} toxicity, it has limitations due to lack of specialized neuronal structures and functions. However, such cell-based systems are necessary for drug discovery and genetic screening, where large, diverse small molecule collections can be tested for their therapeutic efficacy in AD. Such biological assays can cast a much wider net in the discovery of potential leads and even identify undesirable targets of drugs as compared to target-by-target approaches.

Author contributions: Conceptualization, methodology, data acquisition, original draft preparation: PB; writing and review and funding acquisition: RM, PB. Both authors approved the final version of the paper.

Conflicts of interest: The authors have no conflicts of interest to declare.

Financial support: This study was supported by the National Health and Medical Research Council-Australian Research Council dementia research development fellowship (APP1107109) to PB.

Copyright license agreement: The Copyright License Agreement has been signed by both authors before publication.

Data sharing statement: Datasets analyzed during the current study are available from the corresponding author on reasonable request.

Plagiarism check: Checked twice by iThenticate.

Peer review: Externally peer reviewed.

Open access statement: This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non-Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

References

- Bharadwaj P, Martins R, Macreadie I (2010) Yeast as a model for studying Alzheimer's disease. *FEMS Yeast Res* 10:961-969.
- Bharadwaj P, Waddington L, Varghese J, Macreadie IG (2008) A new method to measure cellular toxicity of non-fibrillar and fibrillar Alzheimer's A β using yeast. *J Alzheimers Dis* 13:147-150.
- Bharadwaj P, Head R, Martins R, Raussens V, Sarroukh R, Jegasothy H, Waddington L, Bennett L (2013) Modulation of amyloid-beta 1-42 structure and toxicity by proline-rich whey peptides. *Food Funct* 4:92-103.
- Bharadwaj PR, Dubey AK, Masters CL, Martins RN, Macreadie IG (2009) A β aggregation and possible implications in Alzheimer's disease pathogenesis. *J Cell Mol Med* 13:412-421.
- Bharadwaj PR, Verdile G, Barr RK, Gupta V, Steele JW, Lachenmayer ML, Yue Z, Ehrlich ME, Petsko G, Ju S, Ringe D, Sankovich SE, Caine JM, Macreadie IG, Gandy S, Martins RN (2012) Latrepirdine (Dimebon) enhances autophagy and reduces intracellular GFP-A β_{42} levels in yeast. *J Alzheimers Dis* 32:949-967.
- Caine JM, Bharadwaj PR, Sankovich SE, Ciccotosto GD, Streltsov VA, Varghese J (2011) Oligomerization and toxicity of A β fusion proteins. *Biochem Biophys Res Commun* 409:477-482.
- Carosi JM, Sargeant TJ (2019) Rapamycin and Alzheimer disease: a double-edged sword? *Autophagy* 15:1460-1462.
- Dahlgren KN, Manelli AM, Stine WB Jr, Baker LK, Krafft GA, LaDu MJ (2002) Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *J Biol Chem* 277:32046-32053.
- Fraser PE, Levesque L, McLachlan DR (1993) Biochemistry of Alzheimer's disease amyloid plaques. *Clin Biochem* 26:339-349.
- Huang J, Zhu H, Haggarty SJ, Spring DR, Hwang H, Jin F, Snyder M, Schreiber SL (2004) Finding new components of the target of rapamycin (TOR) signaling network through chemical genetics and proteome chips. *Proc Natl Acad Sci U S A* 101:16594-16599.
- Mangialasche F, Solomon A, Winblad B, Mecocci P, Kivipelto M (2010) Alzheimer's disease: clinical trials and drug development. *Lancet Neurol* 9:702-716.
- McLean CA, Cherny RA, Fraser FW, Fuller SJ, Smith MJ, Beyreuther K, Bush AI, Masters CL (1999) Soluble pool of A β amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease. *Ann Neurol* 46:860-866.
- Mputhia Z, Hone E, Tripathi T, Sargeant T, Martins R, Bharadwaj P (2019) Autophagy modulation as a treatment of amyloid diseases. *Molecules* doi: 10.3390/molecules24183372.
- Nixon RA (2013) The role of autophagy in neurodegenerative disease. *Nat Med* 19:983-997.
- Park SK, Pegan SD, Mesecar AD, Jungbauer LM, LaDu MJ, Liebman SW (2011) Development and validation of a yeast high-throughput screen for inhibitors of A β_{42} oligomerization. *Dis Model Mech* 4:822-831.
- Sarkar S (2013) Regulation of autophagy by mTOR-dependent and mTOR-independent pathways: autophagy dysfunction in neurodegenerative diseases and therapeutic application of autophagy enhancers. *Biochem Soc Trans* 41:1103-1130.
- Selkoe DJ, Hardy J (2016) The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med* 8:595-608.
- Spilman P, Podlutska N, Hart MJ, Debnath J, Gorostiza O, Bredesen D, Richardson A, Strong R, Galvan V (2010) Inhibition of mTOR by rapamycin abolishes cognitive deficits and reduces amyloid-beta levels in a mouse model of Alzheimer's disease. *PLoS One* 5:e9979.
- Steele JW, Lachenmayer ML, Ju S, Stock A, Liken J, Kim SH, Delgado LM, Alfaro IE, Bernales S, Verdile G, Bharadwaj P, Gupta V, Barr R, Friss A, Dolios G, Wang R, Ringe D, Fraser P, Westaway D, St George-Hyslop PH, et al. (2012) Latrepirdine improves cognition and arrests progression of neuropathology in an Alzheimer's mouse model. *Mol Psychiatry* 18:889-897.
- Tian Y, Bustos V, Flajolet M, Greengard P (2011) A small-molecule enhancer of autophagy decreases levels of A β and APP-CTF via Atg5-dependent autophagy pathway. *FASEB J* 25:1934-1942.
- Treusch S, Hamamichi S, Goodman JL, Matlack KE, Chung CY, Baru V, Shulman JM, Parrado A, Bevis BJ, Valastyan JS, Han H, Lindhagen-Persson M, Reiman EM, Evans DA, Bennett DA, Olofsson A, DeJager PL, Tanzi RE, Caldwell KA, Caldwell GA, et al. (2011) Functional links between A β toxicity, endocytic trafficking, and Alzheimer's disease risk factors in yeast. *Science* 334:1241-1245.
- Tripathi T, Kalita P, Martins R, Bharadwaj P (2019) Autophagy promotes memory formation. *ACS Chem Neurosci* 10:3337-3339.

C-Editors: Zhao M, Li CH; T-Editor: Jia Y

© 2020. This article is published under
(<http://creativecommons.org/licenses/by-nc-sa/3.0/>)(the “License”).
Notwithstanding the ProQuest Terms and Conditions, you may use this
content in accordance with the terms of the License.